N. III 8
17

47 76 1/2

74
3
222

1:6

2:2
6
4

WINGATE, E.
C

Thomas Bell

53382/A

35952

Mr. *Wingate's* *Arithmetick*:
CONTAINING
A PLAIN and FAMILIAR
METHOD
For Attaining the
Knowledge *and* Practice
OF
Common Arithmetick.

Compoſed by *EDMUND WINGATE*
of *Grays-Inn*, Eſq;

And upon his Requeſt Enlarged in his Life-Time, alſo
ſince his Deceaſe carefully Reviſed and much Improv'd,
as will appear by the Preface and Table of Contents.
By *JOHN KERSEY*, late Teacher of the Mathematicks.

The Eleventh Edition.

With A new Supplement, Of *Eaſie Contractions*
in the neceſſary *Parts* of *Arithmetick*, uſeful
Tables of Intereſt and *Flemiſh Exchanges*, as
alſo *Practical Menſuration.*

By *GEORGE SHELLEY*, Writing-
Maſter at the *Hand and Pen* in *Warwick-Lane*,
near St. *Pauls*.

LONDON:
Printed for *J. Philips*, at the *King's-Arms*;
J. Taylor, at the *Ship*; and *J. Knapton*, at
the *Crown*, in St. *Paul's* Church-yard. 1704.

85952

WELLCOME LIBRARY INSTITUTE

TO THE
RIGHT HONOURABLE
THOMAS
Earl of *Arundel* and *Surrey*,
Earl Marſhal of
ENGLAND, &c.

Right Honourable,

THE good *Affection you bear to all kind of* Learning, *and in particular to the* Mathematicks, *makes me adventure to preſent your Lordſhip with this Tractate of* Arithmetick, *becauſe that Art, compared with other Mathematical Sciences, is as the* Primum Mobile, *in reſpect of the*

the othor inferior Orbs: For as the Poets uſed in times paſt to ſay of Venus, Sine Cerere & Baccho Friget Venus, *ſo may I alſo confidently averr of them, without* Arithmetick *they are Poor, ond without Motion. Preſuming therefore that your Lordſhip, loving the Art, cannot diſaffect the Artiſt, nor his intention to do good in that kind, I am bold to ſhelter this Treatiſe nnder your Lordſhip's Protection, humbly intreating your gracious Acceptation, and earneſtly deſiring for ever to remain*

Your Honours, in all

Service affectionately

devoted,

EDM. WINGATE.

THE PREFACE OF *JOHN KERSEY.*

ABOUT the Year 1629. our Learned Country-Man *Edmund Wingate*, Eſquire, Publiſhed a Treatiſe of *Arithmetick* divided into two *Books*, the one entituled *Natural Arithmetick*, and the other *Artificial Arithmetick*; and in regard his principal deſign in that Treatiſe was, to remove the difficulties which ordinarily ariſe in the practice of *Common-Arithmetick*, by the help of artificial or borrowed Numbers, called *Logarithms*, (whoſe proper work is to perform *Multiplication* by *Addition*; *Diviſion* by *Subtraction*, &c.)

he did then in his ſaid firſt Book omit divers pieces of *Common* or *Practical Arithmetick*, which, for the perfect and univerſal underſtanding thereof, were neceſſary to have been inſerted: But after the firſt Impreſſion of both thoſe Books was ſpent, our ſaid Author being importuned to take care of the ſecond Edition, he promiſed his Aſſiſtance therein; yet his other neceſſary Employments not permitting him to purſue his ſaid Purpoſe, he was pleaſed to impart his Thoughts concerning the ſame unto me, together with his requeſt, that I would puruſe the ſaid firſt Book, and ſupply it with ſuch pieces of *Practical Arithmetick*, which for the reaſons aforeſaid were wanting in the firſt Edition.

In purſuance of which requeſt, I have contributed my Talent towards perfecting this Tractate, upon our Author's Foundation; partly in his Life-time, to his good liking, and partly ſince his Deceaſe, in ſeveral Editions committed to my Care to be prepared for the Preſs; wherein I have uſed my beſt endeavours, as well to preſerve this Book as a Monument of our ſaid Author's worth, as alſo to make it a compleat Store-houſe of *Common Arithmetick*;

from whence the ingenious may be furniſh'd with the excellencies of that Art, in reference both to common Affairs, as alſo to the practical Parts of the Mathematicks. And in order to thoſe ends I have made theſe following Alterations and Additions; namely,

Firſt, For the Eaſe and Benefit of ſuch Learners, who deſire only ſo much Skill in Arithmetick, as is uſeful in Accompts, Trade, and ſuch like ordinary Employments; the Doctrine of whole Numbers, (which in the firſt Edition was intermingled with Definitions and Rules concerning broken Numbers, commonly called Fractions) is now entirely handled apart. And to the end the full knowledge of *Practical Arithmetick* in whole Numbers might more clearly appear, I have explained divers of the old Rules in the firſt five Chapters, and framed anew the Rules of *Diviſion*, *Reduction*, and the *Golden Rule*, in the ſixth, ſeventh, eighth, and ninth Chapters; ſo that now Arithmetick in whole Numbers is plainly and fully handled before any entrance be made into the craggy Paths of Fractions, at the ſight whereof ſome Learners are ſo diſcouraged, that they make a ſtand, and cry out, *non plus ultra*, there's no progreſs farther

Secondly, To aſſiſt ſuch young Students as deſire to lay a good Foundation for the attaining of a general Knowledge in the Mathematicks, I have in a familiar Method delivered the entire Doctrine of Fractions, both Vulgar and Decimal, which was omitted in the firſt Edition; and have alſo newly framed the Extraction of the Square and Cube Roots, in a method which by Experience is found to be much eaſier than that commonly uſed heretofore, and is exactly ſuitable to the Conſtruction or Compoſition of Square and Cube Numbers.

Laſtly, I have added an *Appendix*, which is furniſhed with variety of choice and delightful knowledge in Numbers, both Practical and Theoretical. In all which performances, I have earneſtly aimed at Truth, Perſpicuity, and exact Correction, both of the Text and Numbers; ſo that I hope this Book is now ſupplied with all things neceſſary to the full Knowledge and Practice of *Common Arithmetick*, the uſefulneſs whereof is ſo generally known, that there will be no need of Arguments to excite any one that deſires his own or the Publick Good, to be acquainted with ſo excellent an Art.

But if the more curious Artiſt, after he is well exercis'd in vulgar Arithmetick, deſires farther inſpection into the Myſteries of Numbers, his beſt Guide is the admirable Art called *Algebra*; the Elements whereof I have expounded at large in a Treatiſe lately Publiſh'd.

JOHN KERSEY.

The Table of Contents.

Where thoſe Chapters of Mr. *Wingate*'s, that have been altered and framed anew by *John Kerſey*, are diſtinguiſhed by this mark, ☞, and thoſe Chapters that have been entirely Compoſed by the ſaid *J. K.* may be diſcovered by this Aſterisk *.

The Doctrine of whole Numbers is contained in the firſt 15 Chapters, the Titles whereof are theſe following.

	Chap.	Pag
☞ *Concerning the Notation of Numbers.*	1	1
Concerning Engliſh Moneys, Weights, Meaſures, &c.	2	8
Addition.	3	16
Subtraction.	4	23
Multiplication.	5	31
☞ *Diviſion.*	6	38
☞ *Reduction.*	7	58
☞ *The Rule of Three direct.*	8	69
☞ *The Rule of Three inverſe.*	9	82
☞ *The double Rule of Three direct.*	10	87
☞ *The double Rule of Three inverſe.*	11	94
The Rule of Three compound of five Numbers.	12	98
The Rule of Fellowſhip.	13	102
The Rule of Alligation.	14	108
The Rule of Falſe.	15	125

The Doctrine of Fractions both Vulgar & Decimal, is contained in the 16th Chapter next following.

Concerning Vulgar Fractions.

	Chap.	Pag
* *Notation.*	16	133
* *Reduction.*	17	137
* *Addition.*	18	151
* *Subtraction.*	19	154
* *Multiplication.*	20	160
* *Diviſion.*	21	164

Concerning Decimal Fractions.

* *Notation.*	22	167
* *Reduction.*	23	173
* *Addition.*	24	211
* *Subtraction.*	25	214
* *Multiplication.*	26	215
* *Diviſion.*	27	226

In Vulgar and Decimal Fractions.

* *The Rule of Three Direct.*	28	240
* *The Rule of Three Inverſe.*	29	245
* *The double Rule of Three.*	30	247
* *The Rule of Falſe.*	31	250

The Extraction of Roots is contained in the two Chapters next following.

☞ *The Extraction of the Square Root*	32	257
☞ *The Extraction of the Cube Root.*	33	270

The Relation of Numbers in Quantity and Quality is contained in the two following Chapters.

The Relation of Numbers in Quantity.	34	290
The Relation of Numbers in Quality, whereof Arithmetical and Geometrical Proportion.	35	295

The Contents of the Appendix.

	Cha.	Pag.
* *Concerning Contractions in the Rule of Three.*	1	309
* *Rules of Practice by* Aliquot *parts*	2	315
* *Exchanges of Coins, Weights and Meaſures.*	3	339
* *Practical Queſtions about Tare, Tret, Loſs, Gain, Barter, Factorſhip, and meaſuring of Tapeſtry.*	4	360
* *Intereſt of Money, with Tables to value Annuities,* &c. *at any rate* per centum *from* 4l. *to* 12l. *and the manner of making thoſe Tables.*	5	368
* *A Demonſtration of the Rule of Three*	6	440
* *A Demonſtration of the double Rule of Fellowſhip.*	7	443
* *A Demonſtration of the Rule of Alligation, with its uſe in the Compoſition of Medicines*	8	446
* *A Demonſtration of the Rule of Falſe.*	9	461
* *A Collection of ſubtil Queſtions to exerciſe all the parts of Vulgar Arithmetick; to which alſo are added various practical Queſtions about the menſuration of Superficial Figures and Solids, with the Gauging of Veſſels.*	10	475
* *Sports and Paſtimes.*	11	528

A
TREATISE
OF
Common Arithmetick.

The Firſt Book.

CHAP. I.

Concerning Notation of Numbers.

I. ARITHMETICK is the Art of Accompting by Number. As Magninitude or Greatneſs is the Subject of *Geometry*, ſo Multitude or Number is that of *Arithmetick*.

Number. II. Number is that by which every thing is numbred; or that which anſwers this queſtion, how many? (unleſs it be anſwered by nothing:) So if it be asked how many days are in a week, the anſwer is ſeven, which is called number.

The Characters by which number is expreſſed.

III. The Notes or Characters, by which Number is ordinarily expreſſed, are theſe; 1 one, 2 two, 3 three, 4 four, 5 five, 6 ſix, 7 ſeven, 8 eight, 9 nine, 0 nothing.

IV. Theſe Notes or Characters are either ſignificant figures, or a Cypher.

V. The ſignificant figures are the firſt nine; *viz.* 1,2,3,4,5,6,7,8,9,. The firſt whereof is more particularly called an Unit, or Unity, and the reſt are ſaid to be compoſed of Unities: ſo 2 is compoſed of two Unities, 3 of three Unities, *&c.*

VI. The Cypher is the laſt, which though of it ſelf it ſignifies nothing, yet being annexed after any of the reſt, it encreaſeth their value: As will appear in the following Rules.

VII. Arithmetick hath two parts, Notation and Numeration.

VIII. Notation teacheth how to expreſs, read, or declare, the ſignification or value of any number written; and alſo to write down any Number propounded, with proper Characters in their due places.

The Places or Degree of any number.

IX. A number is ſaid to have ſo many places or degrees, as there are Characters in the number; *viz.* when divers figures, whether they be intermixt with a Cypher or Cyphers or not, are placed together like letters in a word, without any point, comma, line, or other note of diſtinction interpoſed, all thoſe Characters make but one number, which conſiſts of ſo many places as there are Characters ſo placed to one another: ſo this number 205 conſiſts of 3 places, and this 30600 of five places, &c.

X. Notation conſiſts in the knowledge of two things; viz. the Order of places, and the Value of every place in any number.

The Order of places in any number.

XI. The Order of the places is from the right hand towards the left: So in this number 465, the figure 5 ſtandeth in the firſt place, 6 in the ſecond, and 4 in the third; likewiſe in this number 7560, a Cypher ſtands in the firſt place, 6 in the ſecond, 5 in the third, and 7 in the fourth.

The Value of places in any number.

XII. The firſt place of a Number, (which as before is the outermoſt towards the right hand) is called the place of Units or Unities; in which place any figure ſignifieth its own Simple value: ſo in this number 465, the figure 5 ſtanding in the firſt place ſignifieth five Unities, or five.

XIII. The ſecond place of a number is called the place of Tens; in which place any figure ſignifieth ſo many Tens as the figure containeth Unities: ſo in this number 465, the figure 5 in the firſt place ſignifieth ſimply 5, but the figure 6 in the ſecond place ſignifieth ſix tens, or ſixty.

XIV. The third place of a number is called the place of Hundreds: in which place any figure ſignifieth ſo many Hundreds as there are Unities contain'd in the figure: So in this number 465, the figure 4 in the third place ſignifieth four Hundreds: wherefore if it be required to read or pronounce this number 465, you are to begin on the left hand,

and according to the aforeſaid rules to pronounce it thus, four hundred ſixty five, likewiſe this number 315 is to be pronounced thus, three hundred and fifteen: and this number 205, two hundred and five; alſo this number 500, five hundred. Whence it is manifeſt, that although a Cypher of it ſelf ſignifies nothing, yet beiug placed on the right hand of a figure it encreaſeth the value thereof, by advancing ſuch figure to an higher place than that wherein it would be ſeated, if the Cypher were abſent.

The true reading or pronouncing the value of any number written, as alſo the writing down any number propounded, depends principaly upon a right underſtanding of the three firſt places beforementioned, and therefore I ſhall adviſe the Learner to be well exercis'd therein, before he proceeds to the following Rules.

XV. The fourth place of a number is called the place of Thouſands (that is, any number of Thouſands under ten thouſand;) the fifth place tens of thouſand; the ſixth place hundreds of thouſands; the ſeventh place Millions; (a Million being ten hundred thouſand;) the eighth place tens of Millions; the ninth place hundred of Millions; the tenth place thouſands of Millions; the eleventh place tens of thouſans of Millions; the twelfth place hundred of thouſands of Millions: And in that order you may conceive places to be continued infinitely from the right hand towards the left, each following place being ten times the value of the next preceeding place; but to give names to them would be both a troubleſome and unneceſſary task.

XVI. *From the Rules aforegoing, an eaſie way may be collected to read or expreſs the value of a Number propounded, Viz.* Let it be required to read or pronounce this number 521426341. *Firſt*, Diſtinguiſh by a Comma, or Point, every three places, beginning at the right hand, and proceeding towards the left, ſo will the aforeſaid number be diſtinguiſhed into parts, which may be called *Periods*, and ſtand thus, 521, 426, 341. Where you may note the *firſt period* towards the right hand to conſiſt of theſe figures 341, the *ſecond* of theſe 426, and the *third* of theſe 521. *Secondly*, read or pronounce the figures in every *Period* as if they ſtood apart from the reſt, ſo will the firſt *Period* be pronounced three hundred forty one, the ſecond four hundred twenty ſix, and the third five hundred twenty one. *Thirdly*, to every *Period* except the firſt towards the right hand, a peculiar *denomination* or *ſirname* is to be applyed, *Viz.* the *ſirname* of the ſecond *Period*, is *Thouſands*; of the third, *Millions*; of the fourth, *Thouſands* of *Millions*, *&c.* Therefore beginning to pronounce at the higheſt *Period*, which in this Example is the third, and giving every *Period* its due *ſirname*, the ſaid number will be pronounced thus, *Five hundred twenty one Millions, four hundred twenty ſix Thouſands, three hundred forty one.*

A brief way of Notation.

A Period.

Note, When a number is diſtinguiſhed into *Periods*, as before, the higheſt *Period* will not always compleatly conſiſt of three places, but ſometimes of one place, and ſometimes of two, neverthеleſs after ſuch *Period* is pronounced as if it ſtood apart, the due *ſirname* is to be annexed; ſo this number 3204689, after it is divided into *Periods* will ſtand thus, 3, 204, 689. and to be pronounced thus, *Three Millions, two hundred and four thouſands, ſix hundred eighty nine.*

XVII. The aforeſaid Rules for the right pronouncing or reading of a Number which is written down, being well underſtood, will ſufficiently inform the Reader how to write down any number propounded to be written.

The Table of Notation.

The order of Places.			The values of Places.
	&c.		&c.
Fourth Period,	*Twelfth place*	3	*Hundreds of Thouſand Millions.*
	Eleventh place	2	*Tens of Thouſand Millions.*
	Tenth place	1	*Thouſand Millions.*
Third Period,	*Ninth place*	9	*Hundreds of Millions.*
	Eighth place	8	*Tens of Millions.*
	Seventh place	7	*Millions.*
Second Period,	*Sixth place*	6	*Hundreds of Thouſands.*
	Fifth place	5	*Tens of Thouſands.*
	Fourth place	4	*Thouſands.*
Firſt Period,	*Third place*	3	*Hundreds.*
	Second place	2	*Tens.*
	Firſt place	1	*Unites.*

Notation of Numbers by Latin Letters.

1	I.	21	XXI.
2	II.	30	XXX.
3	III.	40	XL.
4	IIII. or thus IV.	49	XLIX.
5	V.	50	L.
6	VI.	59	LVIIII. or thus LIX.
7	VII.	60	LX.
8	VIII. or thus IIX.	89	LXXXIX.
9	VIIII. or thus IX.	100	C.
10	X.	200	CC.
11	XI.	300	CCC.
12	XII.	400	CCCC.
18	XVIII. or thus IIXX.	500	D. or thus IↃ.
19	XVIIII. or thus XIX.	600	DC. or thus IↃC.
20	XX.	700	DCC. or thus IↃCC.

1000	CIↃ. or thus M.
2000	CIↃ. CIↃ.
3000	CIↃ. CIↃ. CIↃ.
5000	IↃↃ.
10000	CCIↃↃ.

50000	IↃↃↃ.
100000	CCCIↃↃↃ. or thus CM.
500000	IↃↃↃↃ.
1000000	CCCCI. ↃↃↃↃ.
1703	MDCCIII.

CHAP. II.

Concerning Engliſh Money, Weights, Meaſures, &c.

I. *THe things expreſſed by Numbers are principally Money, Weight, Meaſure, Time, and things accompted by the dozen:* Of the three firſt of theſe, there are infinite kinds and varieties according to the diverſity of the ſeveral *Common-wealths* in which they are uſed, all which here to produce were both endleſs and needleſs: wherefore we intend here to treat only of ſuch *Money, Weights, Meaſures,* &c. as are uſed in this *Nation*, being indeed only neceſſary for our preſent purpoſe,

Of Engliſh Money.

II. *The leaſt piece of Money uſed in* England *is a Farthing, from whence this following Table is produced.*

1 *Farthing*	make	1 *Farthing.*
4 *Farthings*		1 *Penny.*
12 *Pence*		1 *Shilling.*
20 *Shillings*		1 *Pound.*

Engliſh (or ſterling) Money is ordinarily written down with Figures after this manner.

l.	s.	d.	f.
34	13	05	2
09	05	10	1
69	00	06	3
00	12	11	0
00	00	07	2

Th

The firſt Rank of the ſaid Numbers ſignifies thirty four pounds, thirteen ſhillings, five pence, two farthings: the ſecond Rank expreſſeth nine pounds, five ſhillings, ten pence, one farthing: the third Rank, ſixty nine pounds, no ſhillings, ſix pence, three farthings, *&c.*

III. The ſmalleſt Weight uſed in *England* is a grain, that is, the weight of a grain of Wheat well dried and gathered out of the middle of the ear, whereof thirty two make another weight, called a Peny-weight, and twenty Peny-weight, make an Ounce Troy.

Vide State. de compoſitione ponderum 51 *Hen.* 3.

Here obſerve, That by the *Statutes* quoted in the Margent, the weight of two and thirty grains of Wheat make a peny-weight, which weight being once diſcovered by two and thirty ſuch grains, the ſaid peny-weight (being the twentieth part of an ounce Troy) is uſually ſubdivided into four and twenty parts only, called alſo *Grains*, as appears by the enſuing Table.

31 *Ed.* 1 *v. Raſt. weights.* 7 & 8. 12 *Hen.* 7. 5.

A Table of Troy Weights.

Troy weight

24 *Grains of Wheat*	make	24 *Artificial Grains.*
24 *Grains*		1 *Peny weight.*
20 *Peny Weight*		1 *Ounce.*
12 *Ounces*		1 *Pound Troy.*

Troy Weight is ordinarily written down with Figures after this manner.

lb.	*oz.*	*p.w.*	*gr.*
17	05	13	13
00	11	07	06
00	00	05	20

The firſt rank of the ſaid numbers expreſſeth ſeventeen pounds, five ounces, thirteen peny weight, thirteen Grains of *Troy weight*: the ſecond rank, no pounds, eleven ounces, ſeven peny weight, ſix grains: and the third, no pounds, no ounces, five peny weight, and twenty grains.

Malynes lex Mercat. p. 49. *Malynes* ib. pag. 252.

Now this *Troy weight* ſerveth only to weigh Bread, Gold, Silver, and Electuaries. And here obſerve alſo by the way, that *Troy weight* regulateth and preſcribeth a form how to keep the Money of *England* at a certain *Standard*. For about two hundred years before the Conqueſt, *Osbright* a Saxon, being then King of *England*, cauſed an ounce *Troy* of Silver to be divided into 20 pieces, at the ſame time called Pence; and ſo an ounce of Silver at that time was worth no more than twenty pence, or one ſhilling eight pence, which continued at the ſame value until the time of *Henry* the ſixth, (who in regard of the enhancing of Money in Foreign Parts) valued the ſame at thirty pence, ſo that then there were accordingly thirty pieces made out of the Ounce, and the old pieces went then for three half-pence, until the time of *Edward* the Fourth, who valued the Ounce at forty pence, and then the old pieces went for two pence apiece. After this, *Henry* the Eight valued the Ounce of ſterling Silver at forty five pence, which value continued until Queen *Elizabeth*'s Time, who valued the ſame Old pence at Three-pence the piece, ſo that all Three-pences coined by the ſame Queen weighed but a peny weight, and every Six-pence two peny weight; and ſo in like manner the Shilling and other pieces accordingly; which made the ounce *Troy* of Silver, to be valued at ſixty pence or five ſhillings, as it now remains at this day without alteration.

IV. The weights uſed by Apothecaries are derived from a pound *Troy*, which is ſubdivided as in the following Table:

Apothecaries weights.

A Table of Apothecaries Weights.

℔	*A Pound Troy*	is equal unto	12 *Ounces.*
℥	*An Ounce*		8 *Drams.*
ʒ	*A Dram*		3 *Scruples.*
℈	*A Scruple*		20 *Grains.*

So that if you were to expreſs in Figures twelve pounds, ten ounces, five drams, two Scruples, and ſixteen grains: alſo three pounds, five ounces, ſeven drams, one ſcruple, and two grains, the ordinary way to write them down is briefly thus:

℔	℥	ʒ	℈	*gr.*
12	10	5	2	16
03	05	7	1	02

V. Beſides *Troy* weight before mentioned, there is another kind of weight uſed in *England*, called Avoirdupois weight, a pound whereof is equal unto 14 Ounces, twelve peny weight *Troy*. This Avoirdupois weight ſerveth to weigh all kind of Grocery-ware, as alſo Butter, Cheeſe, Fleſh, Tallow, Wax and every other thing, which beareth the name of *Garbel*, and whereof iſſueth a refuſe or waſte.

Malynes ib. pag. 49.

VI. Avoirdupois weight is either greater or leſs.

VII. The greater is, when one hundred and twelve pounds Avoirdupois are conſidered as one intire weight

Avoirdupois greater weight.

commonly called an Hundred weight, and then ſuch Hundred weight is ſubdivided firſt into four quarters, and each quarter into eight and twenty Pounds: again, each Pound into four quarters, or (if you will be more exact) into 16 Ounces, and if you pleaſe each Ounce into four Quarters. But ordinarily a Pound is the leaſt quantity that is taken notice of in Avoirdupois groſs weights.

A Table of Avoirdupois greater Weight.

28 *Pounds*	make	*A Quarter of* 112 *lb.*
4 *Quarters*		*An Hundred weight or* 112 *lb.*
20 *Hundred*		*One Tun.*

So that if you were to expreſs by Figures eight Hundred, three Quarters, and five Pounds; likewiſe, ſeven Hundred, one Quarter, and ſeventeen Pounds; the ordinary way to write them down is briefly thus,

C.	*q.*	*lb.*
8	3	5
7	1	17

Avoirdupois leſſer weight. *VIII.* The leſſer Avoirdupois weight is, when a Pound is the higheſt name or Integer, each Pound being ſubdivided into ſixteen Ounces, and each Ounce again into ſixteen Drams, and if you pleaſe each Dram into four Quarters, as by the ſubſequent Table is manifeſt.

A Table of Avoirdupois leſſer Weight.

4 *Quarters of a Dram*	make	1 *Dram.*
16 *Drams*		1 *Ounce.*
16 *Ounces*		1 *Pound.*

So that if you were to expreſs by figures eighteen Pounds, twelve Ounces, five Drams, and three quarters of a Dram; likewiſe five Pounds, no Ounces, twelve Drams, and one quarter of a Dram, the ordinary way to write them down is briefly thus,

℔. oʒ. ʒ q.
18 —— 12 —— 05 —— 3
5 —— 00 —— 12 —— 1

IX. The meaſures uſed in *England* are either of Capacity or Length.

X. The meaſures of Capacity are thoſe which are produced from Weight, and they are either Liquid or Dry.

XI. The Liquid meaſures are thoſe, in which all kind of Liquid ſubſtances are meaſured, and they are expreſſed in the Table following.

Liquid Meaſure.

A Table of Liquid Meaſures.

1 *Pound of Wheat Troy weight*	make	1 *Pint.*
2 *Pints*		1 *Quart.*
2 *Quarts*		1 *Pottle.*
2 *Pottles*		1 *Gallon.*
8 *Gallons*		1 *Firkin of Ale, Soap, Herring.*
9 *Gallons*		1 *Firkin of Beer.*
10 *Gallons and an half*		1 *Firkin of Salmon or Eels.*
2 *Firkins*		1 *Kilderkin.*
2 *Kilderkins*		1 *Barrel.*
42 *Gallons*		1 *Tierce of Wine.*
63 *Gallons*		1 *Hogſhead.*
2 *Hogſheads*		1 *Pipe or But.*
2 *Pipes or Buts*		1 *Tun of Wine.*

Dry Meaſures. *XII.* Dry meaſures are thoſe, in which all kind of dry ſubſtances are meted, as Grain, Sea-coal, Salt, and the like; their Table is this that follows:

A Table of Dry Meaſures.

1 *Pint*	make	1 *Pint.*
2 *Pints*		1 *Quart.*
2 *Quarts*		1 *Pottle.*
2 *Pottles*		1 *Gallon.*
2 *Gallons*		1 *Peck.*
4 *Pecks*		1 *Buſhel land meaſure.*
5 *Pecks*		1 *Buſhel water meaſure*
8 *Buſhels*		1 *Quarter.*
4 *Quarters*		1 *Chaldron.*
5 *Quarters*		1 *Wey.*
2 *Weys*		1 *Laſt.*

Long Meaſures. *XIII.* Long Meaſures are expreſs'd in this Table following.

3 *Barly Corns in length*	make	1 *Inch.*
12 *Inches*		1 *Foot.*
3 *Foot or* 16 *Nails*		1 *Yard.*
3 *Foot nine Inches*		1 *Ell.*
6 *Foot or* 2 *Yards*		1 *Fathom.*
5 *Yards and an half*		1 *Pole or Perch.*
40 *Poles or Perches*		1 *Furlong.*
8 *Furlongs*		1 *Engliſh Mile.*

ALSO

16	*Nails or*	4	*Quarters*	1 *Yard.*
20		5		1 *Ell Engliſh.*
12		3		1 *Flemiſh Ell or Auln.*

Note, That a Yard is uſually ſubdivided into 4 Quarters, and each Quarter into 4 Nails.

And each Ell into 4 Quarters, but each Quarter of an Ell contains 5 Nails.

XIV. Su-

Land Meaſures.

XIV. Superficial or ſquare Meaſures of Land, are ſuch as are expreſs'd in the Table following.

40 *Square Poles or Perches*	make	1 *Rood or quarter of an Acre.*
4 *Roods*	make	1 *Acre.*
640 *Acres*	make	1 *Mile.*

So that if you would expreſs by Figures theſe quantities of Land, *viz.* Thirty ſix Acres, three Roods, twenty Perches: alſo ſeven Acres, no Roods, thirty two Perches; the ordinary way to write them down is thus,

A.	*R.*	*P.*
36	3	20.
7	0	32.

Time.

XV. A Table of Time is this that follows.

1 *Minute*	make	1 *Minute.*
60 *Minutes*	make	1 *Hour.*
24 *Hours*	make	1 *Day natural.*
7 *Days*	make	1 *Week.*
4 *Weeks*	make	1 *Month of twenty eight days.*
13 *Months* 1 *Day, and* 6 *Hours*	make	1 *Year very near.*

But in ordinary computations of time, the whole Year conſiſting of three hundred ſixty five Days, is divided either into twelve equal Parts or Months; each Month then containing thirty Days and ten Hours: or elſe into twelve unequal *Kalendar* Months, according to the antient Verſe:

Thirty Days hath September, April, June, and November:
February hath twenty eight alone, each of the reſt hath thirty one.

Note, That every *Leap-year* (which happeneth once in four Years) containeth three hundred ſixty ſix Days, and in ſuch Year *February* containeth twenty nine Days.

Of things accounted by the dozen.

XVI. Of things accounted by the dozen, a Groſs is the Integer conſiſting of twelve dozen, each dozen containing again twelve particulars: ſo that if you would expreſs in Figures, ſeven Groſs, four Dozen, and five Particulars; alſo four Dozen and eight Particulars, they may be briefly written thus.

G. *D.* *P.*
7——04——05
0——04——08

CHAP. III.

Addition of whole Numbers.

I. COncerning Notation of Numbers, and how thereby the quantities of things are uſually expreſt, a full Declaration hath been made in the preceeding Chapters: Numeration enſueth, which comprehends all manner of operations by Numbers.

II. In Numeration, the four primary or fundamental operations (commonly called Species) are theſe, Addition, Subtraction, Multiplication, and Diviſion.

III. Addition is that by which divers Numbers are added together, to the end that their ſum, aggregate, or total, may be diſcovered.

IV. In Addition, place the Numbers given, one above another in ſuch ſort, that like places or degrees in each number may ſtand in the ſame rank: that is, Units above Units, Tens above Tens, Hundreds above Hundreds, *&c.* So theſe numbers 1213 and 462 being given to be added together, you are to order them as you ſee in the margent.

Addition of numbers of one denomination.

1213
462

V. Having thus placed the Numbers, and drawn a line under them, add them together, beginning with the Units firſt, and ſaying thus, 2 and 3 make 5, which write under the Rank of Units, then proceed to the ſecond Rank and ſay, 6 and 1 make 7, which write under the ſecond Rank (being the place of tens) again 4 and 2 make 6 which write under the third Rank. Laſtly, write down 1 being all that ſtands in the fourth Rank, ſo the ſum of the ſaid given Numbers is found to be 1675, and the operation will ſtand as in the Margent.

1213
462
———
1675

In like manner the numbers 2315, 7423, and 141, being given to be added together, their ſum will be found to be 9879, and the operation thereof will ſtand as you ſee in the Example.

2315
7423
141
———
9879

VI. When the ſum of the Figures of any of the Ranks amounts unto ten, or any number of tens without any exceſs, write down a Cypher under that Rank; but when the ſum of any Rank exceeds ten, or any number of tens, write down the exceſs under ſuch Rank, and for every ten contained in the ſum of any Rank, reſerve an Unit or 1 in your mind, and add ſuch Unit or Units to the Figures of the next Rank towards the left hand; ſo the numbers 4937, 9878, and 394 being given to be added together, the operation will be thus, *viz.* beginning with the rank of Units, I ſay 4, 8 and 7 make 19, wherefore I write down 9, the exceſs above ten, and carrying 1 in mind inſtead of the ten contained in the ſaid 19, I ſay 1 and 9 (9 being the lowermoſt figure of the ſecond rank) make 10, which added to 7 and 3, the other figures of the ſame rank, the whole ſum of them is 20, wherefore ſetting down a Cypher under the line in that rank (becauſe the exceſs above the two tens is nothing) I carry 2 to the third rank, and ſay 2 and 3 (3 being the lowermoſt figure of the third rank) make 5, which being added to 8 and 9 (the other figures of the ſame rank) the ſum of them is 22, wherefore writing down 2 (being the exceſs above the two tens) under the line in the third rank, I carry 2 in mind (becauſe there were two tens in 22) to the fourth rank, and ſay 2 and 9 make 11, which added to 4 makes 15, this 15 becauſe it is the ſum of the laſt rank I write totally down under the line, on the left hand of the Figures before ſubſcribed; ſo the ſum of the three Numbers given is found to be 15209, as in the Example.

$$\begin{array}{r} 4937 \\ 9878 \\ 394 \\ \hline 15209 \end{array}$$

Addition of Numbers of divers Denominations.

VII. When numbers given to be added, do expreſs things of divers Denominations; firſt write them down orderly (according to the Examples in Chap. 2.) then after a line is drawn under them all, begin to add the numbers of the leaſt Denomination, and if the ſum of them amounts to one Integer, or many Integers of the next greater Denomination, with ſome exceſs of the leſs Denomination, write down that exceſs, or a Cypher when there is no exceſs, under the line, ſo as it may ſtand under the leaſt Denomination, and keep the ſaid Integer or Integers in mind, to be added to thoſe of the next greater Denomination, on the left hand: But when the ſum of the numbers of the leaſt Denomination amounts not to one Integer of the next greater Denomination, ſet down the ſum it ſelf under the line; then add the Integer or Integers kept in mind (when any happens) to the numbers of the next greater Denomination on the left hand, and proceed to add them, as alſo thoſe of every greater Denomination, in like manner as above is directed, untill you come to the numbers of the greateſt (or higheſt) Denomination, which are to be added according to the fore-going Rules *V.* and *VI.* of this Chapter. So theſe ſeveral ſums 24*l.*——13*s.*——5*d.*——3*f.* Alſo 12*l.*——0*s.*——8*d.* and 5*l.*——18*s.*——2*f.* being propounded to be added, their total ſum is 42*l.*——12*s.*——2*d.*——1*f.* For having written them down orderly according to the ſecond Rule of the Second Chapter, and drawn a line underneath, I begin with the Farthings firſt, and ſay, two Farthings and three Farthings make five Farthings, that is, one Peny with a Farthing over and above; wherefore ſetting down 1 under the Denomination of Farthings, I

l.	*s.*	*d.*	*f.*
24—	13—	05—	3
12—	00—	08—	0
05—	18—	00—	2
42—	12—	02—	1

carry one Penny to the denomination of Pence, then I ſay, 1, 8; and five pence make 14 Pence, which contain one Shilling and two Pence, wherefore writing two under the denomination of Pence; I likewiſe carry 1 Shilling to the denomination of Shillings: Then adding the ſaid 1 Shilling unto 18 Shillings and 13 Shillings, the ſum will be found 1 Pound and 12 Shillings, wherefore ſetting down 12 under the denomination of Shillings, I carry 1 Pound in mind unto the denomination of Pounds ſaying, 1 Pound in mind, together with 5, 2, and 4 Pounds which ſtand in the firſt rank of Pounds, make 12 Pounds, wherefore (according to the ſixth Rule of this Chapter) I write 2, the exceſs above 10, underneath the ſaid firſt rank of Pounds, and carry 1 in mind for the ſaid 10 to the ſecond rank of Pounds, then ſaying in like manner; 1 in mind, together with 1 and 2 which ſtand in the ſecond rank of Pounds make 4, which I write underneath the line, that done, I find the total of the three ſums propounded to be 42*l*—12*s.*—and 2*d.*—1*f.*

In like manner 3 *lb.*--05 *oz.*--19 *p.w.*--15 *gr.* Alſo 2 *lb.*—0 *oz.*—3 *p.w.*—7 *gr.* Alſo 0 *lb.*—10 *oz.*—6 *p.w.*—0 *gr.* And 0 *lb.*—9 *oz.*—0 *p.w.*—17 *gr.* being given to be added together, their ſum will be found 7 *lb.*—01 *oz*—9 *p.w.*—15 *gr.* and the work will ſtand thus,

lb.	*oz.*	*p.w.*	*gr.*
03	05	19	15
02	00	03	07
00	10	06	00
00	09	00	17
07	01	09	15

Note, In adding together the Numbers in the laſt Example, it muſt be remembred that twenty four Grains make one Penny-weight; twenty Penny-weight one Ounce; and twelve Ounces one Pound *Troy* (as before declared in the third Rule of the ſecond Chapter;) And then you are to proceed according to Rule VII. of this Chapter.

More Examples of Rule VII. are theſe following, which preſuppoſe the Learner to be well exercis'd in the Tables of Chap. 2. that he may readily know what Integers are to be carried from every leſſer Denomination to the next greater.

Addition of Engliſh Money.

lb.	*s.*	*d.*	*f.*
230	17	10	1
175	12	11	3
052	05	00	0
009	00	08	1
506	13	00	2
974	19	06	3

l.	*s.*	*d.*
0	13	05
0	17	08
0	00	10
0	10	03
0	15	06
2	17	08

Addition of Troy Weight.

lb.	*oz.*	*p. w.*	*gr.*
23	07	16	13
17	10	15	07
325	06	19	20
49	11	07	12
417	00	19	04

oz.	*p. w.*	*gr.*
536	13	16
208	11	10
063	10	05
099	00	12
907	15	19

Addition of Avoirdupois Weight.

C.	*q.*	*lb.*	*lb.*	*oz.*	*dr.*
235	3	13	14	13	12
[illegible]76	1	17	05	10	14
628	2	15	12	00	06
412	0	10	06	09	05
1852	3	27	39	02	05

Addition of Meaſures of Length.

yards.	*q.*	*nails.*	*Ells.*	*q.*	*nails.*
26	3	2	15	3	2
13	1	3	16	1	3
12	0	1	09	0	1
29	1	1	12	2	1
81	2	3	53	3	2

Addition of Superficial Meaſures of Land.

Acres.	*Roods.*	*Per.*	*A.*	*R.*	*P.*
136	3	13	240	2	17
513	1	26	500	3	13
212	2	10	249	1	36
517	0	00	006	0	10
1379	3	09	996	3	36

CHAP. IV.

Subtraction of whole Numbers.

I. SUbtraction is that by which one number is taken out of another, to the end that the remainder or difference, between the two numbers given may be known.

II. The Number out of which the Subtraction is to be made, muſt be greater, or at leaſt, equal with the other. As you may Subtract, 4347 or 9478 out of 9478, ſo can you not ſubtract 9478 out of 4347.

Subtraction of numbers of one denomination.

III. In ſubtraction rank the two given numbers one under the other as in Addition, with this caution, that the number placed uppermoſt may exceed, or at leaſt be equal unto the other: So if the number 4347 be given to be ſubtracted from 9478, I order them as in the Margin: then proceeding to the Subtraction, I ſay 7 taken out of 8, there remains 1, which I place in the ſame rank under the Line. In like manner 4 being taken out of 7, the remainder is 3, which likewiſe I ſet under the line in the next rank; again taking 3 from 4, the remainder is 1, which I likewiſe place under the third rank; laſtly, ſubtracting 4 from 9, there will remain 5; which I ſubſcribe under the fourth rank; ſo the whole operation being finiſhed, I find, that if 4347 be taken out of 9478, the remainder is 5131, or (which is the ſame) the difference between the numbers 9478 and 4347 is 5131, as in the Example:

$$\begin{array}{r}9478\\ \underline{4347}\\ 5131\end{array}$$

In like manner if 106 be ſubtracted from 2856 the remainder will be found 2750; for after the numbers are orderly ranked, I begin at the place of Units, and ſay, 6 from 6, there remains nothing; wherefore I ſubſcribe 0, then proceeding to the ſecond rank, I ſay, if 0 (or nothing) be taken from 5, there will remain 5, which I alſo ſubſcribe under the line; again 1 from 8, there remains 7: laſtly, 0 from 2, there remains 2. See the work in the Margin.

2856
106
——
2750

IV. When any of the figures of the number given to be ſubtracted is greater than the upper figure out of which it is to be ſubtracted, you muſt borrow 10 of the next rank towards the left hand, and add the ſaid 10 to the ſaid upper figure, then from the ſum of ſuch Addition ſubtract the lower figure and ſet down the remainder: In this caſe the figure of the next rank which is to be ſubtracted; muſt be eſteemed an unite greater than it is; wherefore keeping one in your mind add it to the next figure of the number given to be ſubtracted, and deducting all out of the figure above it, proceed in like ſort till you have finiſhed the whole operation. *Example,* Let it be required to ſubtract 374 out of 8023. Having ranked them as before, I ſay 4 out of 3, that cannot be, wherefore borrowing 10 of the next rank, and adding the ſame to the ſaid 3, I ſay 4 out of 13, there remains 9; then writing 9 under the line, and carrying 1 in my mind, I ſay 1 and 7 make 8, 8 out of 2 that cannot be, but 8 out of 12 (12, becauſe 10 being borrowed, and added to 2, makes 12) there remains 4, which I ſubſcribe under the line

8023
374
——
7649

line; again 1 in my mind being added to 3 makes 4, 4 out of nothing, that cannot be, but 4 out of 10 there remains 6, which I likewiſe ſubſcribe under the line; laſtly 1 in mind being taken out of 8 there remains 7. Thus you ſee that the remainder after 374 is ſubtracted from 8023 is 7649. Note diligently, that as often as 10 is borrowed, 1 muſt be kept in mind to be added to the figure ſtanding in the next place of the lower number, and the ſum of ſuch Addition muſt be ſubtracted from the upper place; but if it happen that there is no figure in the next place of the lower number, then the 1 in mind muſt be ſubtracted from the upper place, (as in the laſt rank of the laſt Example.) *Another Example.* Let it be required to ſubtract 92 from 62801. Having placed the greater number uppermoſt, and the leſſer orderly underneath, I begin at the place of Units, and ſay, 2 from 1 I cannot take, but borrowing 10, and adding it to the ſaid 1, I ſay 2 from 11 there remains 9, which I ſubſcribe under the line; then I proceed and ſay, 1 in mind with 9, makes 10, 10 out of 0 I cannot take, but borrowing 10, I ſay 10 out of 10 and there remains 0, wherefore I ſubſcribe 0 under the line; again, 1 in mind out of 8, there remains 7; then becauſe there are no more Figures in the lower number, I ſay 0 out of 2 there remains 2; laſtly, 0 out of 6 there remains 6; therefore I conclude that 62801 exceeds 92 by 62709.

$$\begin{array}{r} 62801 \\ 92 \\ \hline 62709 \end{array}$$

Subtraction of numbers of divers denominations.

V. If the numbers propounded have divers denominations, place them as before, and beginning with the leaſt denomination firſt, ſubtract the lower number from the upper when it may be ſubtracted, and place the remainder underneath; but if it happen that the lower number cannot be taken out of the upper, you muſt borrow an Integer of the next greater denomination on the left hand; which Integer, after it is converted into the ſame denomination with the ſaid upper number, muſt be added to it: Then from the ſum of ſuch Addition, you are to ſubtract the lower number, and write down the remainder, keeping 1 (that is the Integer borrowed) in your mind, to be added to the next place of the number given to be ſubtracted, as before: So 90*l.*——14*s.*——10*d.*——3*f.* being ſubtracted from 124*l.*——11*s.*——7*d.*——1*f.* the remainder is 33*l.*——16*s.*——8*d.*——2*f.* For beginning with the Farthings, I ſay, 3 Farthings out of 1 Farthing I cannot take, wherefore borrowing 1 Penny (that is an Integer of the next greater denomination) and having converted this Penny into 4 Farthings, I add them to the aforeſaid 1 Farthing; ſo the ſum is 5 Farthings, out of which ſubtracting 3 Farthings, there remains 2 Farthings, which I place underneath the denomination of Farthings; then I proceed to the next denomination, and ſay 1 Penny which I borrowed and 10*d.* make 11*d.* this 11*d.* out of 7*d.* I cannot take, wherefore borrowing 1 Shilling or 12*d.* and adding 12*d.* to the ſaid 7*d.* the ſum is 19*d.* from which I ſubtract the ſaid 11*d.* ſo there remains 8*d.* which I ſubſcribe under the denomination of Pence; again 1 Shilling which I borrowed being added to 14*s.* makes

l.	*s.*	*d.*	*f.*
124	11	07	1
90	14	10	3
33	6	08	2

makes 15*s.* which I cannot ſubtract out of 11*s.* and therefore I borrow 1 Pound or 20*s.* which being added to the ſaid 11*s.* makes 31*s.*from which ſubtracting 15*s.* there remains 16*s.* which I ſubſcribe under the denomination of Shillings; then carriyng 1 Pound which I borrowed to the lower place of Pounds, I ſay 1 in mind with 0 makes 1, which taken out of 4, there remains 3; again 9 out of 2, I cannot take, but 9 out of 12 (10 being borrowed and added to the ſaid 2, according to the fourth Rule of this Chapter) and there remains 3. Laſtly, 1 (for the 10 that was borrowed) being taken out of 1, there remains nothing; and ſo at laſt I find,that if A. being indebted to B. in 124——11*s.*——7*d.*——1*f.* hath paid in part thereof 90*l.*——14*s.*——10*d.*——3*f.* there remains yet undiſcharged 33*l.*——16*s.*——8*d.*——2*f.*

Subtraction of many numbers from one number.

VI. When many numbers are given to be ſubtracted from a number propounded, you muſt firſt add thoſe numbers together,according to theRules of the third Chapter, and then the ſum found is to be ſubtracted from the number firſt propounded. *Example*, A. being indebted unto B. in 3240*l.* paid thereof at one time 700*l.* at a ſecond payment 1236. and at a third 305*l.* the Queſtion is how much of the debt remained undiſcharged? Firſt, I add together the ſeveral ſums paid, and find the total to be 2241*l.* this I ſubtract from 3240*l.* ſo there remains 999*l.* undiſcharged as you ſee by the operation in the Margin.

l.	
3240	*The Debt*
700	
1236	*Payments*
305	
2241	*Total paid*
999	*reſt unpaid*

	l.	*s.*	*d.*
The Debt	500	00	00
Payments	340	12	06
	13	18	03
	17	16	10
Paid in all	372	07	07
Reſt unpaid	127	12	05

Another Example of like nature. A. being indebted to B. in 500*l.* paid in part thereof at one Payment. 340*l.*——12*s.*——06*d.* at a ſecond Payment 13*l.*——18*s.*——3*d.* at a third 17*l.*————16*s.*——10*d.* the Queſtion is how much was in arrear? Here if the operation be proſecuted as before it will appear that there was 127*l.*——12*s.*——05. unpaid: See the work in the Margin.

The proof of Addition and Subtraction.

VII. Addition is proved by Subtraction, and Subtraction by Addition. For having added divers numbers together, if you ſubtract one of them out of the ſum, the remainder muſt be equal to all the reſt, as you may obſerve by the Example following, *viz.* ſuppoſing

236	
——	
452	934
29	236
217	698
934	
698	

theſe 4 numbers are given to be added *viz.* 236, 452, 29, 217. and that their ſum is found to be 934 (by the Rules of the third Chapter) it is required to prove whether the ſaid ſum be true or not; to perform this, I draw a line under the uppermoſt number 236, to ſeparate it from the reſt, and ſeek the ſum of all the numbers given, except that uppermoſt, which ſum I find to be 698. Then I ſubtract the ſaid uppermoſt number 236 from 934, (the total ſum of all the numbers firſt found) and becauſe the remainder 698 is the ſame with the ſum of all the numbers, excluding the uppermoſt, I conclude that the ſum of all the numbers firſt found was truly computed.

In like manner is Subtraction proved by Addition, for if you add the remainder, and the number given to be ſubtracted together, the ſum muſt be equal to the number out of which the Subtraction is made, ſo if 4347 be ſubtracted from 9478 the remainder is 5131, for if 5131 be added to 4347, the ſum is 9478, which is the ſame with the number out of which the Subtraction was made. Again, if a Servant receive 24*l.*——13*s.*——7*d.* and lay out or diſburſe 19*l.*——15*s.*——08*d.* there muſt remain in his hands——4*l.*——17*s.*——11*d.* for this being added to 19*l.*——15*s.*————08*d.* which was the Mony he expended, the ſum will be equal to 24*l.*——13*s.*——07*d.* (being the Money wherewith he was firſt charged.)

Example 1.

out of	9478
ſubtr.	4347
Reſt	5131
Proof	9478

Example 2.

l.	*s.*	*d.*
24	13	07
19	15	08
04	17	11
24	13	07

More Examples of Subtraction are theſe that follow.

Subtraction of Engliſh Money.

	l.	*s.*	*d.*	*f.*
Rec.	3090	10	07	1
paid	0099	14	08	3
reſt	2990	15	10	2
proof	3090	10	07	1

l.	*s.*	*d.*	*f.*
24	00	00	0
05	17	11	3
18	02	00	1
24	00	00	0

Subtraction of Troy Weight.

	lb.	*oz.*	*p.w.*	*gr.*	*oz.*	*p.w.*	*gr.*
Boug.	352	10	13	15	205	13	19
Sold	019	11	16	18	118	16	20
Reſt	332	10	16	21	86	16	23
Proof	352	10	13	15	205	13	19

Subtraction of Avoirdupois Weight.

	C.	*q.*	*lb.*	*lb.*	*oz.*	*dr*
Bought	256	2	23	25	13	12
Sold	079	3	26	00	14	13
Reſt	176	2	25	24	14	15
Proof	256	2	23	25	13	12

Subtraction of Superficial Meaſures of Land.

	Acres,	*Roods,*	*Perch.*	*A.*	*R.*	*P.*
Bought	780	2	35	2040	1	20
Sold	090	3	36	919	3	30
Reſt	689	2	39	1120	1	30
Proof	780	2	35	2040	1	20

Queſtions to exerciſe Addition and Subtraction.

Qu. 1. Two Perſons, A. and B. owe ſeveral debts the leſſer debt being that of A. is 3045*l.* the difference of their debts is 104*l.* what is the debt of B? *Anſ.* 3149*l.*

Queſt. 2. Two Perſons A. and B. are of ſeveral Ages, the Age of the elder, being that of A. is 70, the difference of their Ages is 19, what is the Age of B? *Anſwer,* 51.

Queſt. 3. What number is that which being added to 168 maketh the ſum to be 205? *Anſ.* 37.

Queſt. 4. The ſum of two Numbers is 517, the leſſer is 40, what is the greater? *Anſ.* 477.

Queſt. 5. A certain Perſon born in the year of our Lord 1616, deſired to know his Age in the Year 1676, what was his Age? *Anſ.* 60.

Queſt. 6. The greater of two numbers is 130, their difference is 49, what is the leſſer number? *Anſ.* 81.

CHAP. V.

Multiplication of whole Numbers.

I. MUltiplication teacheth how by two Numbers given to find a third, which ſhall contain either of the numbers given ſo many times as the other contains 1 or unity.

II. Of the two numbers given in Multiplication, one (which you will) is called the Multiplicand and the other the Multiplicator, (or both are called factors.)

III. The number ſought, or ariſing by the multiplication of the two numbers given, is called the product, the Fact, or the Rectangle: ſo if 5 be given to be multiplied by 3 or 3 by 5, the product is 15, that is 3 times 5, or 5 times 3 makes 15: and here 5 may be called the Multiplicand, and 3 the Multiplicator, or 3 may be called the Multiplicand, and 5 the Multiplicator; and as 3 (one of the two numbers given) containeth 1 or unity thrice, ſo 15 the product containeth 5 (the other given number thrice; likewiſe as 5 (one of the given numbers) contains unity 5 times, ſo 15 (the product) contains 3 (the other given number) 5 times.

IV. Multiplication is either ſingle or compound.

Single Multiplication.

V. Single Multiplication, is, when the Multiplicand and Multiplicator conſiſt each of them of one onely figure as in the laſt Example. In like manner if you multiply 9 by 5, the product is 45, this is likewiſe ſingle Multiplication: now the ſeveral varieties of ſingle Multiplication are well expreſs'd in the Table following, uſually called *Pythagora*'s Table.

The Table of Multiplication.

1	2	3	4	5	6	7	8	9
2	4	6	8	10	12	14	16	18
3	6	9	12	15	18	21	24	27
4	8	12	16	20	24	28	32	36
5	10	15	20	25	30	35	40	45
6	12	18	24	30	36	42	48	54
7	14	21	28	35	42	49	56	63
8	16	24	32	40	48	56	64	72
9	18	27	36	45	54	63	72	81

The uſe of the *Table* is this, having one figure given to be multiplied by another to know the product of them, find the multiplicand in the top of the *Table.* and the multiplicator in the firſt Column thereof towards the left hand; this done, in the angle of poſition juſt againſt thoſe two figures you ſhall find the product. So 9 being given to be multiplied by 5. I find 9 in the top of the Table, and 5 in the firſt Column towards the left hand, then carrying my Eye from 5 in a right line equidiſtant to the upper ſide or top line of the Table, until I come to that ſquare which is directly under 9, I find 45, which is the product required. The particular varieties of this *Table* ought to be learned by heart, (that is, a Man muſt be able to give the Product of any ſingle multiplication, without the leaſt pauſe or ſtay) before he can readily work compound multiplication, as will further appear hereafter.

VI. Compound Multiplication, is when the multiplicator and multiplicand, either one or both, conſiſts of more figures than one.

Compound Mulplication.

VII. In compound Multiplication, when the numbers given do end with ſignificant figures, place them as in Addition and Subtraction. So 134 being given to be multiplied by 2, place them thus: Then proceeding to the multiplication ſay thus: 2 times 4 is 8, which write under the line in the rank of your multiplying figure; again, ſay 2 times 3 is 6, which likewiſe write under the line in the next rank: Laſtly, 2 times 1 is 2, which being likewiſe written down under the line in the next rank, the product is diſcovered to be 268, and the work will ſtand as in the Margin.

$$\begin{array}{r} 134 \\ 2 \\ \hline 268 \end{array}$$

VIII. When the Multiplicator conſiſts of more figures than one, as many figures as it hath, ſo many ſeveral products muſt be ſubſcribed under the line, which at laſt being added into one ſum, gives you the total product of all. So 1232 being given to be multiplied by 23, the operation thereof will ſtand thus, for 1232 being multiplied by 3, (according to the laſt Rule) the product is 3696. Again, 1232 being multiplied by 2, the product is 2464, which ſeveral products, after they are placed in their due order, (that is, the firſt figure ariſing in each product under his reſpective multiplying figure) and added together, produce 28336, the product required: In like manner 1321 being given to be multiplied by 123, the product is 162483, and the operation will ſtand as you ſee in the Margin.

```
 1232
   23
 ----
 3696
2464
-----
28336

  1321
   123
  ----
  3963
 2642
1321
------
162483
```

IX. When the product of any of the particular figures exceeds ten, place the exceſs under the line as before, and for every ten that it ſo exceeds, keep one in mind to be added to the next Rank.

Example, 3084 being given to be multiplied by 36, the work will ſtand thus; for 6 times 4 being 24, I write 4 under the line, and reſerve 2 in mind for the two tens; then I ſay 6 times 8 is 48, unto which if I add 2 kept in mind, the whole is 50, wherefore ſubſcribing 0 in the next rank under the line 0 (becauſe there is no exceſs of 50 above 5 tens) I reſerve 5 in mind for the 5 tens; again, I ſay 6 times

```
  3084
    36
 -----
 18504
 9252
------
111024
```

nothing is nothing, to which adding 5 that I kept in mind, the whole will be but 5, which I likewiſe ſubſcribe under the line in the next rank; again 6 times 3 is 18, which (in regard 3 is the laſt figure of the multiplicand) I write wholy down; ſo that the particular product ariſing from the multiplying figure 6 is 18504. In like manner proceeding with the multiplying figure 3, the particular product ariſing will be 9252. Laſtly, theſe ſeveral products being placed in due order, and added together (after the manner of the 8th Rule of this Chapter) will give 111024, which is the total product ariſing from the multiplication of 3084 by 36, and the operation will ſtand as in the Margin. After the ſame manner if 5073 be given to be multiplied by 256, the product will be found to be 1298688, and the operation will ſtand as you ſee in the example.

```
   5073
    256
  -----
  30438
 25365
10146
-------
1298688
```

X. When the two numbers given to be multiplied, do one or both of them end with a Cypher or Cyphers towards the right hand, multiply the ſignificant figures in both numbers, one by the other, neglecting ſuch Cyphers, and when the multiplication of the ſignificant figures is finiſhed, annex on the right hand of the number produced by the multiplication the Cypher or Cyphers with which one or both of the numbers firſt given did end, ſo will the whole give you the true product demanded: *Example*, 43100 being given to be multiplied by 15000 the product will be found for omitting the Cyphers which ſtand

```
    43100
    15000
---------
2155
431
---------
646500000
```

in the laſt places towards the right hand as well in the multiplicand as the multiplicator, I multiply the ſignificant figures 431, by the figures 15 (according to the former Rules,) ſo there will ariſe 6465, to which annexing on the right hand all the Cyphers before omitted, the true product will be 646500000: more Examples hereof are theſe following.

$$\begin{array}{rcr}
43125 & \qquad & 5108000 \\
1500 & & 125 \\
\hline
215625 & & 25540 \\
43125 & & 10216 \\
\cline{1-1}
64687500 & & 5108 \\
\cline{3-3}
 & & 638500000
\end{array}$$

XI. When in the multiplicator Cyphers are included between ſignificant figures, multiply by the ſaid ſignificant figures, neglecting ſuch Cyphers or Cypher, but obſerve diligently to ſet the particular products of the ſignificant figures in their due places according to the 8th Rule of this Chapter. So if 56324 be given to be multiplied by 20006, I firſt multiply the whole multiplicand 56324 by 6, and place the product orderly underneath the line, then paſſing over the three Cyphers, I multiply 56324 by 2 and place 8 (which is the firſt exceſs of this particular product) directly under the multiplying figure 2, and the reſt in their order, ſo at laſt the true product will be found 1126817944, and the work will ſtand as you ſee in the Example.

$$\begin{array}{r}
56324 \\
20006 \\
\hline
337944 \\
112648 \\
\hline
1126817944
\end{array}$$

More Examples hereof are theſe that follow.

$$\begin{array}{r} 3094 \\ 104 \\ \hline 12376 \\ 3094 \\ \hline 321776 \end{array} \qquad \begin{array}{r} 23765 \\ 10302 \\ \hline 47530 \\ 71295 \\ 23765 \\ \hline 244827030 \end{array}$$

Note, That one of the principal *cautions* to be obſerved in Multiplication, is the due placing of the particular products ariſing by each multiplying figure; and that may be performed either by taking care to place the firſt figure or Cypher which ariſeth in each Product under the reſpective multiplying figure; or at leaſt the firſt place ariſing in the ſecond Product muſt ſtand under the ſecond place of the firſt Product, and the firſt place of the third particular Product, under the third place of the firſt, *&c.*

XII. When a number is given to be multiplied by a number that conſiſts of 1 (or an unit) in the firſt place towards the left hand, and a Cypher or Cyphers on the right hand of ſuch an unit (ſuch are 10, 100, 1000, 1000, &c. the multiplication is performed by annexing the Cypher or Cyphers of the multiplicator at the end (to wit on the right hand) of the multiplicand; ſo if 326 be given to be multiplied by 10, the Product is 3260; if by 100, the Product is 32600; if by 1000, the Product is 326000; in like manner if 170 be multiplied by 10, the Product is 1700; if by 100, 17000, &c

XIII. When more numbers than two are given to be multiplied one by the other, that kind of Multiplication

Continual Multiplication.

is called *Continual*, and is thus performed, *viz.* firſt multiply any two of the numbers given one by the other, then multiply the Product by another of the numbers given, and this product by the fourth number given (if there be ſo many) and in that order till every one of the given numbers hath been made a Multiplicator, ſo the laſt product is the true product required. *Example*, if 4, 18, and 22 were given to be multiplied *continually*, firſt 18 multiplied by 4 produceth 72, which multiplied by 22 (the third number) produceth 1584, the laſt product or number required. See the work in the Margin. The proof of Multiplication is by Diviſion as will appear by the next Chapter.

```
  18
   4
  ——
  72 Prod. 1.
  22
 ———
 144
144
————
1584 Prod. 2.
```

CHAP. VI.

Diviſion by whole Numbers.

I. DIviſion is that by which we diſcover, how often one number is contained in another; or (which is the ſame) it ſheweth how to divide a number propounded into as many equal parts as you pleaſe.

II. In Diviſion there are always three remarkable Numbers which are commonly called by theſe Names, the *Dividend*, the *Diviſor*, and the *Quotient*.

III. The *Dividend* is the Number given to be divided into equal Parts.

IV. The Diviſor is the Number by which the Dividend is to be divided; that is, it is the number which declareth into how many equal parts the dividend muſt be divided.

V. The Quotient is the number ariſing from the diviſion, and ſheweth one of the equal parts required; ſo if 15 were given to be divided by 5, or into 5 equal parts, the number ariſing, or one of the equal parts will be 3, for 5 is found three times in 15: And here 15 is the *Dividend*, 5 the *Diviſor*, and 3 the *Quotient*.

Diviſion by a ſingle figure.

VI. Diviſion being the hardeſt leſſon in Arithmetick, muſt be heedfully intended by the Learner, for whoſe eaſe I ſhall uſe my utmoſt endeavours to make the way ſmooth by Rules and Examples, beginning with the eaſieſt firſt, which will be in that caſe when the Diviſor conſiſts of one figure only; for example, Let it be required to divide 192, by 8, or 192 pounds into 8 equal parts or ſhares; here 192 is the *Dividend*, 8 is the *Diviſor*, and the *Quotient* or one of the equal parts is ſought.

VII. Place a crooked line at each end of the *Dividend*, that on the left hand ſerving for the place of the *Diviſor*, and that on the right for the *Quotient*; then if the *Diviſor* be a ſingle figure, ſubſcribe a point under the firſt figure of the *Dividend* towards the left hand, if ſuch firſt figure be either equal unto, or greater than the *Diviſor*, but if ſuch firſt figure be leſs than the *Diviſor*, put a point under the next place of the *Dividend*; which number o diſtinguiſhed by the point may be alled the *Dividual*; ſo in the example

What the dividual is.

8) 192 (

given in the 6th Rule, 192 being the *Dividend* and 8 the *Diviſor*, I ſubſcribe a point under 9, not under 1, becauſe it is leſs than the *Diviſor*. This done the *Dividual*, or number whereof the queſtion muſt be asked, is 19.

VIII. Having thus prepared the numbers, ask how often the *Diviſor* is contained in the *Dividual*, and write the number which anſwers the queſtion in the *Quotient*; then multiply the *Diviſor* by the number placed in the *Quotient*, and ſubſcribe the product underneath the *Dividual*. Laſtly, having drawn a line under the product, ſubtract it from the *Dividual*, and ſubſcribe the remainder orderly underneath the line. So demanding how many times the *Diviſor* 8 is found in the *Dividual* 19, the anſwer is two times, wherefore I write 2 in the *Quotient*; then multiplying the *Diviſor* 8 by 2 (the number placed in the *Quotient*) the product is 16, which I ſubſcribe orderly under the *Dividual* 19; and after a line is drawn underneath the product 16, I ſubtract it from the *Dividual* 19, and place the remainder 3 underneath the line.

8) 192 (2
16·
―――
3

IX. Put another point under the next place of the *dividend* towards the right hand, and bring down the Figure or Cypher ſtanding in that place to the remainder; that is ſet it next after it, ſo the whole will be a new *Dividual*: Thus a point being placed under 2 which ſtands in the next place of the *Dividend*, I write 2 next after (to wi on the right hand of) the remainder 3, ſ is 32 a new *Dividual*, or number whereof the ſecond queſtion muſt be asked and the work will ſtand as you ſee in the Example

8) 192 (2
16·
―――
32

X. I

X. A new *Dividual* being ſet apart, renew the queſtion and proceed according to the 8th Rule of this Chapter. Thus demanding how often the *Diviſor* 8 is found in the *Dividual* 32, the anſwer is four times; wherefore I write 4 in the *Quotient*, then multiplying the *Diviſor* 8 by 4 (the figure laſt placed in the *Quotient*) the product is 32, which I ſubſcribe under the *Dividual* 32, and after a line is drawn underneath, I ſubtract the product 32 from the *Dividual* 32, and there being no remainder, I ſubſcribe 0 under the line, ſo the whole work being finiſh'd, the *Quotient* is found to be 24, and the operation ſtands as you ſee in the Example; wherefore I conclude, if 192 pounds be equally divided amongſt 8 Perſons, the ſhare of each Perſon will be 24 Pounds.

8)192(24
16
32
32
0

A ſecond Example. Let it be required to divide 936 Pounds into 9 equal parts; having diſtinguiſhed the firſt *Dividual* by a point, (according to the 7th Rule of this Chapter) I demand how often the *Diviſor* 9 is found in the *Dividual* 9, and finding it once contained in it, I write 1 in the *Quotient*; then multiplying the *Diviſor* 9 by 1, the product is 9, which I ſubſcribe under the *Dividual* 9, after this, a line being drawn under the product 9, I ſubtract it from the *Dividual* 9, and there being no remainder, I place a 0 underneath the line, as you ſee in the Example.

9)936(1
9
0

Again placing a point under 3 which ſtands in the next place of the *Dividend*, I tranſcribe the ſaid 3 next after the remainder 0 for a new *dividual*, then asking

9)936(10
9
03

how often the *Diviſor* 9 is contained in the *Dividual* 3, and not finding it once contained therein, I write 0 in the *Quotient*, and now becauſe the product which ought to ariſe from the Multiplication of the *Diviſor* by 0 (the Cypher laſt placed in the *Quotient*) amounts to 0, the *Dividual* 3, out of which that product ſhould have been ſubtracted, remains the ſame without alteration; wherefore after a point is ſubſcribed under ſix, the next place of the *Dividend*, I annex 6 to the *Dividual* 3, ſo there will be a new *Dividual*, to wit 36, then demanding how often the *Diviſor* 9 is found in the *Dividual* 36, the anſwer will be 4 times; wherefore I place 4 in the *Quotient*, and multiplying the *Diviſor* 9 by 4, the product is 36, which I ſubſcribe under, and ſubtract from the *dividual* 36, ſo the remainder is 0, thus the whole work being finiſht, the *Quotient* is found to be 104, as you ſee in the *Example*; wherefore I conclude, if 936*l.* be divided equally amongſt 9 Perſons, the ſhare of each will be 104*l.* In like manner if 296163 be divided by 7, the *Quotient* will be 42309.

```
9) 936 (104
   ...

    9
 -------
   036
    36
 -------
     0
```

The ſubſtance of diviſion by what method ſoever.

The whole work of Diviſion is briefly contained in this following Verſe.

Dic quot, multiplica, ſubduc, transferque ſecundum.

Or thus,

Firſt you muſt ask how oft, in Quotient anſwer make;
Then Multiply, Subtract, a new Dividual take.

A Compendious way of dividing by a ſingle figure.

XI. When in the Diviſion the *Diviſor* conſiſts of a ſingle Figure only, the *Quotient* may be written down, and all the operation performed in mind, without writing down any part thereof, ſo 82506 being given to be halfed or divided into two equal parts, the work will be thus, The *Diviſor* 2 is found in 8 four times; in 2 once; in 5 twice, and there will remain 1, which 1 being ſuppoſed to ſtand before (to wit, on the left hand of) the Cypher, makes 10, then I ſay 2 is found in 10, 5 times; and laſt of all in 6, 3 times; ſo that the true *Quotient* or one half of the given number 82506 is found to be 41253.

2)82506(41253

In like manner if 82506 be given to be divided by 3, or into 3 equal parts, the work will be thus, the *Diviſor* 3 is found in 8 twice, and there will remain 2, which 2 being ſuppoſed to ſtand before (to wit, on the left hand of) the following 2, makes 22; then I ſay 3 is found in 22, 7 times; in 15, 5 times, in 0 not at all, and laſtly in 6 twice; ſo that the true *Quotient* or one of the 3 equal parts required is 27502. After the ſame manner may diviſion be wrought by any ſingle figure, without much charge to the memory.

3)82506(27502

A note, concerning the remainder after the Diviſion is ended, if any happen.

Note, Here the *Learner* may ask what ſhall be done with the laſt remainder, if any happen, when the *Diviſion* is finiſhed? For a full anſwer to this, I refer the *Reader* to the Note in the fifth *Rule* of the ſeventh *Chapter*; yet I ſhall here propound an example where the ſaid caſe happens, *viz.* Let it be required to divide 351 by 8, or 351 *Pounds* equally amongſt 8 *Perſons*; now if the operation be proſecuted according to the former rules, the *Quotient* will be found to be 43, and after the *Diviſion*

7
8) 351 (43
15

is finiſh'd, there will remain 7, that is, each Perſon muſt have 43 Pounds, and there will be an overplus of 7 Pounds, which muſt be alſo divided equally among the 8 Perſons, but that cannot be done till the 7 Pounds be reduced into Shillings, and then thoſe Shillings muſt be divided by 8 to give every Perſon his due ſhare of the Shillings contained in the ſaid 7 Pounds; again, if there yet remain any ſurpluſage of Shillings, they muſt be reduced to Pence, which muſt alſo be divided by 8, to give every Perſon his due ſhare of Pence: ſo that when this queſtion is fully anſwered, each Perſons ſhare will appear to be 43*l.*——17*s.*——6*d.* But how the before-mentioned *Reduction* is performed, will be made manifeſt in the fifth Rule of the next Chapter.

Diviſion by two or more figures, the firſt and eaſieſt Method.

XII. When the Diviſor conſiſts of two, three or how many places ſoever, the operation is more difficult than the former, but depends upon the ſame grounds, and therefore the Learner being well vers'd in the preceding method of dividing by a ſingle figure, will the more readily underſtand theſe that follow, which are two, whereof the firſt is the eaſier, but the latter more expeditious, and that which indeed is principally to be aimed at: For an example of the former, let it be required to divide 4112772 by 708, or (which is the ſame) to divide 4112772 into 708 equal parts.

Firſt, A *Table* is to be made to ſhew at firſt ſight any *Multiple* or Product of the *Diviſor*, it being taken twice, thrice, or any number of times under ten, ſo having firſt written down the *Diviſor* it ſelf 708, and drawn a line on the right hand thereof, I place 1 on the right hand of the line directly againſt

The Deviſor.	708	1
Multiples of the Diviſor.	1416	2
	2124	3
	2832	4
	3540	5
	4248	6
	4956	7
	5664	8
	6372	9

againſt the *Diviſor*; then underneath the *Diviſor* 708 I ſubſcribe the double thereof, which is 1416, and place the figure 2 directly againſt the ſaid double, to wit, on the other ſide of the line. Again, adding 1416 (to wit the double, of the *Diviſor*) to the *Diviſor* it ſelf 708, the ſum is 2124 for the tripple of the *Diviſor*, this triple I ſubſcribe under the double, and place 3 on the other ſide of the line right againſt the triple. Again adding 2124 (the triple of the *Diviſor*) to the *Diviſor* 708, I find 2832 for the quadruple of the *Diviſor*, which quadruple I ſubſcribe under the triple, and proceeding in like manner, at laſt the table is finiſht, which readily ſhews the *Diviſor*, with the *duple, triple, quadruple, quintuple, ſextuple, ſeptuple, octuple, and noncuple* of the *Diviſor.*

Now for a proof of the ſaid Table, adding the laſt number thereof, to wit, 6372 (which was found to be nine times the *Diviſor*) to the *Diviſor* 708, I find the ſum to be 7080, which (by the 12th Rule of the fifth Chap.) is evident ten times the *Diviſor*; wherefore I conclude that the Table is true, in regard that the laſt number thereof is derived from all the ſuperior numbers.

The Table of Multiples or Products of the *Diviſor* being thus prepared, write down the *dividend* on the right hand of the *Diviſor*; then diſtinguiſh by a point ſo many of the foremoſt places of the *Dividend* towards the left hand, as are either equal in value (being conſider'd apart) to the *Diviſor*, or which

```
 708|1)4112772(5809
1416|2 3540···
    |  ----------
2124|3  5727
2832|4  5664
    |   ---------
3540|5    6372
4248|6    6372
    |   ---------
4956|7       0
5664|8
6372|9
```

which being greater, yet come neareſt to the value thereof, thus I ſubſcribe a point under 2, thereby ſetting apart 4112, being the feweſt of the foremoſt places which will contain the *Diviſor* 708, ſo is 4112 the *dividual* (or number whereof the firſt Queſtion muſt be asked;) then demanding how often the *Diviſor* 708 is contained in the *dividual* 4112, the anſwer will be found by the *Table* to be five times, for looking in the *Table* I cannot find the *dividual* exactly, but I ſee that 6 times the *Diviſor* is the next greater than the *dividual* 4112, and five times is the next leſſer; wherefore I write 5 in the *Quotient*, and the number in the *Table* which ſtands againſt 5, to wit, 3540 I ſubſcribe under the *Dividual* 4112, then having drawn a line underneath, I ſubtract 3540 (which is five times the *Diviſor*) from the *Dividual* 4112, and ſubſcribe the remainder 572 underneath the line; that done, I put a point under the next place of the *Dividend* towards the right hand, and becauſe the figure 7 ſtands in that place, I tranſcribe 7 next after the remainder 572, ſo there is 5727 for a new *Dividual.*

Then demanding how often the *Diviſor* 708 is contained in the *Dividual* 5727, the anſwer will be found by the *Table* to be 8 times, for looking in the *Table* I find that 9 times the *Diviſor* is the next greater, but 8 times is the next leſſer than the *Dividual*, wherefore I write 8 in the *Quotient*, and the number in the *Table*, which ſtands againſt 8, to wit, 5664 I ſubſcribe under, and ſubtract from the *Dividual* 5727,placing the remainder 63 underneath the line.

Again, I put a point under the next place of the *Dividend*, where I find the figure 7, and therefore tranſcribing 7 next after the remainder 63, the new *Dividual* will be 637, then demanding how often the *Diviſor* 708 is contained in the *Dividual* 637, and not finding it once contained therein, I write 0 in the *Quotient*, and ſince in this caſe (that is when a Cypher anſwers the Queſtion) the *Dividual* remains the ſame without alteration, the figure or Cypher ſtanding in the next place of the *Dividend* is to be tranſcribed after the *Dividual* for a new *Dividual*, ſo writing 2 next after 637, the new *Dividual* is 6372, wherefore demanding how often the *Diviſor* 708 is contain'd in 6372, I find by the *Table* it is contain'd in it 9 times, wherefore writing 9 in the *Quotient*, and placing the number which ſtands againſt 9 in the *Table*, to wit, 6372 under the *Dividual* 6372, and ſubtracting it from the *Dividual* there will remain 0. Wherefore I conclude if 4112772 be divided by 708, or into 708 equal parts, the true *Quotient* or one of the equal parts required is 5809.

In like manner if 20304 be divided by 188, that is into 188 equal parts, the *Quotient* ariſing, or one of thoſe equal parts will be 108,and the operation will ſtand as you ſee.

Diviſor.	188	1
Multiples of the Diviſor.	376	2
	564	3
	752	4
	940	5
	1128	6
	1316	7
	1504	8
	1692	9

```
) 20304 (108
   ...
  188
  ---
   1504
   1504
   ----
      0
```

The preceding method of *Diviſion* by the help of a *Table* of the *Multiples* or products of the *Diviſor*, as it is moſt eaſie, ſo in ſome Caſes namely, where the *Diviſor* is great, and a *Quotient* of many places is required, as in calculating *Tables* of *Intereſt*, *Aſtronomical Tables*, and ſuch like) it excels all other ways of *Diviſion*, both in reſpect of certainty and expedition, but for common practice it is too tedious,and therefore I ſhall proceed to the choiceſt practical method.

The latter and choiceſt practical Method of Diviſion, when the Diviſor conſiſts of many places.

XIII. *I now come to the laſt and principal method of Diviſion, when the Diviſor conſiſts of many places, which to ſuch as have the Table of Multiplication by heart will not be difficult*; For example, let 56304 be a number given to be divided by 184, that is, into 184 equal parts and the *Quotient* or one of the equal parts is required.

Firſt, diſtinguiſh by a point (as before) ſo many of the foremoſt places of the *dividend* towards the left hand, as are either equal in value (when they are conſider'd apart) to the *Diviſor*, or elſe, which being greater, yet come neareſt unto it, thus I ſubſcribe a point under the figure 3, thereby ſetting apart 563, being the feweſt of the foremoſt places which will contain the *Diviſor*; ſo is 563 the *dividual*, or number whereof the firſt queſtion muſt be asked. Having thus prepar'd the numbers, I demand how often the *Diviſor* 184 is contained in the *dividual* 563; and ſince to anſwer this queſtion and ſuch like, there is a neceſſity of tryal, it will be requiſite to ſhew how this tryal may fitly be made; firſt therefore

184) 56304(

fore compare the number of places in the *dividual* with the number of places in the *Diviſor*, and when the number of places is the ſame in both, let it be asked how often the firſt or extream figure of the *Diviſor* towards the left hand is contained in the firſt figure of the *dividual* towards the ſame hand; ſo here demanding how often 1 is contained in 5, the anſwer is 5 times; whence I infer that the *Diviſor* 184 is not contained oftner than 5 times in the *dividual* 563 (for 6 times 184 is manifeſtly greater than 563) but whether it be contained 5 times in it or not, examination muſt be made either by multiplying (in ſome by-place) the *Diviſor* 184 by the ſaid 5, and comparing the product with the *dividual*, 563; or elſe thus, ſaying 5 times 1 (to wit the 1 in the *Diviſor*) is contained in 5, to wit, the firſt figure of the *dividual* 563, 5 times, but then 8 the following figure of the *Diviſor*, cannot be found 5 times in 6, the following figure of the *dividend*, and conſequently the *Diviſor* 184 is not contained 5 times in the *dividual* 563; wherefore I make another tryal to ſee whether it may be contained 4 times in it or not, ſaying 4 times 1 is 4 which is found in 5, and there will remain 1, but then 4 times 8, which is 32, cannot be had in 16 (for the 1 before remaining being ſuppoſed to ſtand on the left hand of 6 maketh 16) hence I conclude again, that the *Diviſor* 184 is not contained 4 times in the *dividual* 563; wherefore I make another tryal to ſee whether it may be contained 3 times in it or not, ſaying 3 times 1 is 3, which is found in 5, and there will remain 2, again, 3 times 8 is 24, which is found in 26 (for the 2 before remaining being ſuppoſed to ſtand before the 6 in the *Dividual* makes 26)and there will remain 2:Laſtly,three times 4 is 12,which is likewiſe found in 23, (for the 2 remaining in the 26 being ſuppoſed to ſtand before the 3 in the *Dividual* makes 23)whereby I ſee that the *Diviſor* 184 is contained 3 times in the *Dividual* 563,wherefore I write 3 in the *Quotient*, and proceeding according to the 8th*Rule* of this*Chapter*, I multiply the *Diviſor* 184 by 3 (the figure placed in the *Quotient*) ſo the *Product*, is 552, which I ſubſcribe orderly underneath the *Dividual*563, then having drawn a line underneath the ſaid *Product*, I ſubtract it from the *Dividual*,and ſubſcribe the remainder which is 11 under the line.

```
184) 56304 (3
     552
    -----
      11
```

```
184) 56304 (306
       ...
     552
    -----
      1104
      1104
     -----
         0
```

Again according to the 9th *Rule* of this *Chapter*, I bring down 0 which ſtands in the next Place of the *Dividend*, to the remainder 11, ſo there is 110 for a new *Dividual*, then demanding how often the *Diviſor* 184 is found in the *Dividual* 110, and not finding it once contained in it, I write 0 in the *Quotient* (which is to be done as often as the Queſtion is anſwered by nothing;) now becauſe the *Product* ariſing from the multiplication of the *Diviſor* by 0 (the Cypher laſt placed in the *Quotient*) amounts to 0; the *Dividual* 110 out of which that *Product* ſhould be ſubtracted, remains the ſame without alteration; wherefore after a point is ſubſcribed under 4 the following place of the *Dividend*, I annex 4 to the laſt *Dividual* 110, and there will be a new *Dividual*,to wit,1104;and here the Queſtion at large is to know how often 184 is found in 1104:but to leſſen the trial, becauſe the *Dividual* conſiſts of one place more than is in the *Diviſor*, it muſt be asked how often the firſt figure of the *Diviſor* on the left hand is contained in the two foremoſt places of the *Dividual* towards the left hand, *viz.* I demand how often 1 is contained in 11, and although it may be had 11 times, yet I need never begin the trial above 9 times, therefore I make trial with 9, ſaying 9 times 1 is 9, which is found in 11, and there will remain 2; but then 9 times 8 which is 72 cannot be found in 20 (20 becauſe the 2 remaining being ſuppoſed to ſtand before 0 in the *Dividual* makes 20) therefore I make trial with 8 ſaying 8 times 1 is 8, which is found in 11, and there will remain 3, but then 8 times 8 cannot be had in 30 (30 becauſe the 3 remaining being ſuppoſed to ſtand before the 0 or Cypher makes 30) therefore I make trial with 7, ſaying 7 times 1 is 7, which is found in 11, and there will remain 4; but then 7 times 8 cannot be had in 40, therefore I make trial with 6, ſaying 6 times 1 is 6, which is found in 11, and there will remain 5; then 6 times 8 is 48, which is found in 50, and there will remain 2; laſtly, 6 times 4 is 24, which is found in 24, whereby at length I ſee that the *Diviſor* 184 is contained 6 times in the *Dividual* 1104, wherefore I write 6 in the *Quotient*, and proceeding according to the 8th *Rule* of this *Chapter*, I multiply the *Diviſor* 184 by 6 (the figure laſt placed in the *Quotient*) ſo the *Product* is 1104, which being ſubſcribed under and ſubtracted from the *Dividual* 1104, the remainder is 0, ſo at laſt I conclude that the *Quotient* ſought is 306.

Note, If the figure aſſumed for the *Quotient* holds good upon tryal, as aforeſaid, by two or three of the foremoſt places of the *dividual*, it will for the moſt part hold thoughout the *dividual*; but this muſt be a perpetual Rule, that whenſoever the *Product* of the multiplication of the *Diviſor* by the figure placed in the *Quotient* happens to be greater than the *dividual*, from which it ought to be ſubtracted, ſuch *Product* muſt be ſtruck out of the work, and a leſſer figure is to be placed in the Quotient.

For a ſecond *Example*, let it be required to divide 15114220 by 2987, or into 2987 equal parts.

Firſt, the *Diviſor* 2987 being greater than 1511, (to wit, the four foremoſt places of the Dividend) I ſet a point under 4, thereby ſetting apart 15114 for a *Dividual*; then becauſe the *Dividual* conſiſts of one place more than the *Diviſor*, I ask how often 2 (the firſt figure of the *Diviſor* to-towards the lefthand) is contained in 15 the two foremoſt places of the *dividual*) and finding the anſwer to be 7 times, I infer thence that the *diviſor* 2987 cannot be contained more than 7 times in the *dividual* 15114; but whether it will be contained 7 times in it or not, examination muſt be made, either by multiplying 2987 by 7 (in ſome by-place) and by comparing the Product with the *dividual* 15114, or elſe by the manner of tryal before delivered in the laſt Example: ſo at length it will be diſcovered that the *diviſor* 2987 will not be found above 5 times in the *dividual* 15114; wherefore (according to the 8th *Rule* of this *Chapter*) writing 5 in the Quotient, and multiplying 2987 by 5, I ſub-

2987) 15114220 (5
14935
179

I ſubſcribe the product of that Multiplication, which is 14935, under the *Dividual* 15114, then drawing a line underneath the ſaid Product, and ſubtracting it from the *Dividual* 15114, I ſubſcribe the remainder 179 under the line.

```
2987)15114220(50
      14935 .
      ———————
       1792
```

Again (according to the 9th *Rule* of this *Chapter*) I bring down 2, the next place of the Dividend, to the ſaid Remainder 179, ſo the new Dividual will be 1792; that done, asking how often the *Diviſor* 2987 is contained in the *Dividual* 1792, and not finding it once contained in it, I write o in the Quotient; and here becauſe the queſtion is anſwered by o, the next place of the *Dividend*, to wit 2, is to be brought down to the *Dvidual* 1792, ſo the new *Dividual* is 17922. Then renewing the queſtion, and proceeding as before, at length the Diviſion being finiſh'd, the Quotient will be found 5060 exactly, without any Remainder; but if any Remainder had hapned after the ſubtraction of the laſt Product it muſt have been proſecuted according to he note before given in the Example at the latter end of the 11th *Rule* of this *Chapter*.

```
2987)15114220(5060
     ....

     14935
      17922
      17922
     ———————
       00
```

In like manner if 120839550 be divided by 9999, or into 19999 equal Parts, the *Quotient*, or ne of thoſe equal parts, will be found 60450, and he work will ſtand as you ſee here.

19999) 1208939550 (60450
.....
119994
89995
79996
99995
99995
00

This latter method of Diviſion is to be preferr'd before any of the common ways of dividing by daſhing out of figures, where the ſteps of the Diviſion are ſo confounded (beſides the burden upon the memory, by a promiſcuous Multiplication and Diviſion) that if any error happen, it can hardly be corrected without beginning the work anew; But in the way before explained, the particular Multiplications, Subtractions, and Remainders, which belong to every figure of the Quotient, are ſo diſtinctly and clearly expreſt, that if an errour happen, the work may eaſily be reformed.

How the number of places in the Quotient may be diſcovered.

XIV. So often as the queſtion is repeated in Diviſion, ſo many places there muſt be in the quotient (which may be diſcovered by the number of Points placed under the *dividend)* and ſo many times is one and the ſame kind of operation repeated, the ſubſtance whereof is contained in the Verſe before-mentioned at the end of the tenth *Rule* of this *Chapter*.

A Compendious way of dividing by 10, 100, 1000, &c.

XV. When the *Diviſor* conſiſts of 1 or an unit in the extream place towards the left hand, and nothing but Cyphers towards the right, the diviſion is performed by cutting off with a line ſo many places of the *Dividend* towards the right hand as the *Diviſor* hath Cyphers; ſo the figures which ſtand on the left hand of the line, give the Quotient, and thoſe cut off to the right (if they be ſignificant figures) are to be proceeded with as a ſurpluſage or overplus remaining, according to the Note at the end of the eleventh *Rule* of this *Chapter*. So if 4720*l.* were given to be divided equally amongſt 10 perſons, the ſhare of each would be 472*l.* alſo if the ſaid 4720*l.* were to be divided equally amongſt 100 perſons, the ſhare of each would be 47 *l.* and there would be a ſurpluſage or remainder of 20 *l.* to be alſo ſubdivided amongſt them, after the ſaid 20 *l.* are converted into ſhillings according to the fifth *Rule* of the next *Chapter*. Laſtly, if the ſaid 4720*l.* were to be divided amongſt 1000 perſons, the ſhare of each would be 4*l.* and there would be a remainder of 720*l.* to be alſo divided as aforeſaid. See the form of the Work in the *Margin*.

10) 472|0 (472
100) 47|20 (47
1000) 4|720 (4

Another Compendium in Diviſion.

XVI. When the *Diviſor* conſiſts of any ſignificant figure or figures in the firſt or foremoſt place or places towards the left hand, and nothing but a Cypher or Cyphers towards the right, cut off by a line ſo many places of the *Dividend* towards the right hand as the *Diviſor* hath Cyphers towards the right; then divide the figures of the *Dividend*, which ſtand on the left hand of the line, by the figures in the *Diviſor* which remain, when the ſaid Cypher or Cyphers are omitted, remembring after the diviſion is finiſhed, to write down next after the laſt remainder the places of the *Dividend* which were firſt cut off: So if 36732 were given to be divided by 20, the Quotient will be 1836, and there will remain 12, *viz.* if you cut off one place from the *Dividend* towards the right hand (becauſe the *Diviſor* ends with one Cypher) and then divide the reſt, to wit, 3673 by 2 (according to the 11th *Rule* of this *Chapter*) there will ariſe in the Quotient 1836, and the laſt remainder after ſuch diviſion is finiſh'd, will be 1, unto which if 2 (the figure firſt cut off from the *Dividend*) be annexed, the total remainder is 12.

```
          1
2|0) 3673|2(1836
```

In like manner if 7456787, were given to be divided by 304000, the Quotient will be 24, and there will remain 160787; *viz.* If you cut off 3 places from the *Dividend* towards the right hand (3 places, becauſe the *Diviſor* ends with 3 Cyphers) and then divide 7456 by 304, there will ariſe in the Quotient 24, and the laſt remainder, after ſuch diviſion is finiſh'd, will be 160, unto which if 787 (the places firſt cut off from the *Dividend*) be annexed, the total remainder or ſurpluſage is 160787, which is to be proceeded with, as is directed in the Note at the latter end of the eleventh *Rule* of this *Chapter*.

```
304|000) 7456|787 (24
         608
         ----
         1376
         1216
         ----------
           160787
```

The Proof of Multiplication and Diviſion.

XVII. Diviſion and Multiplication do interchangeably prove one another; for in Diviſion if you multiply the *Diviſor* by the Quotient, the Product will be equal to the *Dividend*: So in the *Example* of the 13th *Rule* of this *Chapter*; if 184 the *Diviſor* be multiplied by 306 the *Quotient*, the Product is 56304, which is the ſame with the *Dividend*; but when, after the whole Diviſion is finiſhed, any figures remain of the laſt Subtraction, add them likewiſe to the Product: So in the laſt *Example* of the 16th *Rule* of this *Chapter*, the *Diviſor* 304000 being multipled by the *Quotient* 24, produceth 7296000, unto which if you add the number remaining, to wit, 160787, the ſum is 7456787, which is the ſame with the *Dividend*. Again in Multiplication, if the product be divided by the Multiplicator, the *Quotient* will give you the Multiplicand, or if the product be divided by the Multiplicand, the *Quotient* will give you the Multiplicator: So in the firſt Example of the 9th *Rule* of the laſt Chapter, if the product 111024 be divided by the Multiplicand 3084, the *Quotient* gives the Multiplicator 36.

There is alſo of Multiplication a *Common proof* argued from the Multiplicand, the Multiplicator and the Product by caſting away nines, but by that way of Proof (though rightly uſed) *a falſe* Product will be affirmed to be true: Example, if 3462 be multiplied by 786, the true Product is 2721132; but if I ſay 4953132 or 3153132 is the Product (or many others which may be given) the proof by nines will confirm them to be true Products, though they are falſe, as will be evident to ſuch as know the *Rule*, which I mention here only to ſet a brand upon it, that it may be avoided by all lovers of Truth.

CHAP. VII.

Reduction.

I. FOraſmuch as in *Money*, there are diverſities of kinds, *viz.* in *England*, *Pounds*, *Shillings*, *Pence*, and *Farthings*; alſo divers kinds of *Weights*, *Meaſures*, &c. as hath been fully declared in the ſecond Chapter; and becauſe it is often times required to find how many pieces of one kind of *Money* are equal in value to a given number of another (and ſo likewiſe of *Weights*, *Meaſures*, &c.) it will be convenient in this place to ſhew how that is performed, ſince thereby the *Rules* of *Multiplication* and *Diviſion* before delivered will be exercis'd This kind of operation is called *Reduction*.

II. Reduction is either deſcending or aſcending.

III. Reduction deſcending is, when ſome Integers of a Number of greater denomination being given, it is required to find how many Integers of a leſſer denomination are equal in value to that given number of the greater: As when it is required to find how many *ſhillings* are contained in 30*l.* Likewiſe how many *pence* in 320*s.* or how many *hours* in 365 *days*, &c.

IV. Reduction aſcending is, when ſome Integers of a number of leſſer denomination being given, it is required to find how many Integers of a greater denomination are equal in value to that given number of the leſſer: As when it is required to find how many *pence* are contained in 500 *farthings*: likewiſe how many *ſhillings* in 348 *pence*: or how many *days* in 864 *hours*: &c.

V. *Reduction* deſcending is performed by *Multiplication*, for if the given number of Integers of a greater denomination be multiplied by a number, which expreſſeth how many Integers of the leſſer are equal to one of the Integers given, the Product is the number of Integers of the leſſer denomination required.

Reduction deſcending is performed by Multiplication.

So 230*l.* of Engliſh Money will be reduced into 4600 *s.* for if 230 be multiplied by 20 (the number of *ſhillings* which are equal to 1 *pound)* the product is 4600; in like manner 4600*s.* will be reduced into 55200 *d.* for if 4600 be multiplied by 12(the number of *pence* contained in 1 *ſhilling*) the product is 55200. Alſo 55200 *pence* being multiplied by 4 becauſe 4 *farthings* make a *penny*) are reduced into 220800 *Farthings*, as by the operation in the *Margin* is evident.

$$\begin{array}{rl} 230 & \textit{pounds.} \\ 20 & \\ \hline 4600 & \textit{ſhillings.} \\ 12 & \\ \hline 92 & \\ 46 & \\ \hline 55200 & \textit{pence.} \\ 4 & \\ \hline 220800 & \textit{Farthings.} \end{array}$$

The like method is to be obſerved in *Weights*, *Meaſures*, &c. So 345 *Ounces Troy* are reduced into 6900 *peny weights*, and 6900 *penny weights* to 165600 *Grains*, as by the operation in the *Margin* you may ſee.

$$\begin{array}{rl} 345 & \textit{Ounces.} \\ 20 & \\ \hline 6900 & \textit{penny W.} \\ 24 & \\ \hline 276 & \\ 138 & \\ \hline 165600 & \textit{Grains.} \end{array}$$

Note, by this *Rule* the Learner is furniſhed with Skill to reſolve that caſe in Diviſion, when the *dividend* is *leſs* than the *diviſor* :

Compare this with the Note upon the laſt Example of the 11th Rule of the 6th Chapter.

Example, Let it be required to divide 7 *Pounds* of Engliſh Money equally amongſt 8 Perſons; here it is evident that the *Dividend* 7 is leſs than the *Diviſor* 8; that is, the number of *Pounds* is leſs than the number of Perſons, and conſequently each ſhare muſt be leſs than a Pound; ſo that in effect it is required to find how many *Shillings* and *Pence* belong to each Perſon for his ſhare: Firſt, therefore reduce the 7 *Pounds* into *Shillings*, which will be 140, theſe divided by 8 give 17 *Shillings* to each Perſon, and there will yet be a remainder of 4 *Shillings* to be alſo equally divided into 8 parts, but theſe 4 *Shillings* muſt be firſt reduced into *Pence*, which will be 48, then dividing 48 by 8, the *Quotient* will give 6 *Pence* more to every Perſon: ſo at laſt it appears that if 7 *Pounds* of Engliſh Money be equally divided into 8 parts, the entire *Quotient* (or one of the equal ſhares) will be 17 *Shillings* and 6 *Pence*.

In like manner, if 354 *Pounds* of Engliſh Money be given to be divided equally amongſt 125 Perſons, the ſhare of each will be found to be 2 *Pounds*, 16 *Shillings*, 7 *Pence*, 2 *Farthings*, and ſomewhat more, but the parts of a *Farthing* being of no moment (and not properly to be handled in this place) are neglected.

Compare theſe two Examples with the laſt Example of the eleventh Rule of the ſixth Chapter.

In *Reduction* deſcending, the Learner may receive help by the ſubſequent *Tables*.

1. *Of Engliſh Moneys.*

Pounds	*Multiplyed by*	20	*Produce*	*Shillings.*
Shillings.		12		*Pence.*
Pence		4		*Farthings.*

2. *Of Troy Weight.*

Pounds	*Multiplyed by*	12	*Produce*	*Ounces.*
Ounces		20		*Penny Weights.*
Penny Weight		24		*Grains.*

Alſo in *Apothecaries Weights.*

Ounces Troy	*Multiplyed by*	8	*Produce*	*Drams.*
Drams		3		*Scruples.*
Scruples		20		*Grains.*

3. *Of Averdupois Weights.*

Hundred W.	*Multip. by*	4	*Produce*	*Quarters.*
Quarters		28		*Pounds.*
Pounds		16		*Ounces.*
Ounces		16		*Drams.*

4. *Of Liquid Meaſures.*

Hogſheads	*Multip. by*	63	*Produce*	*Gallons.*
Gallons		2		*Pottles.*
Pottles		2		*Quarts.*
Quarts		2		*Pints.*

5. *Of Dry Meaſures.*

Quarters	*Multiplyed by*	8	*Produce*	*Buſhels.*
Buſhels		4		*Pecks.*
Pecks		2		*Gallons.*
Gallons		2		*Pottles.*
Pottles		2		*Quarts.*
Quarts.		2		*Pints.*

6. *Of Long Meaſures.*

Engliſh Miles	*Multiplied by*	8	*Produce*	*Furlongs.*
Furlongs		220		*Yards.*
Yards		3		*Feet.*
Feet		12		*Inches.*
Inches		3		*Barley Corns.*

Alſo,

Yards	*Mult. by*	4	*Produce*	*Quarters.*
Ells		5		*Quart. of Yards.*
Quarters		4		*Nails.*

7 *Of Superficial Meaſures of Land.*

Acres	*Mult. by*	4	*Produce*	*Roods.*
Roods		40		*Perches or* **Poles.**

8. *Of Time.*

Weeks	*Mult. by*	7	*Produce*	*Days.*
Days		24		*Hours.*
Hours		60		*Minutes.*

To reduce Integers of divers Denominations into the loweſt of thoſe Denominations

VI. Integers of divers denominations may be reduced into the laſt of thoſe denominations according to the fifth *Rule* aforegoing, by deſcending orderly to the next inferiour denomination,

nation; and adding to each Product ſuch Integers (if there be any) which are of the ſame name.

So 12 *Pounds* 13 *ſhillings*, and 10 *pence* may be reduced into 3046 *pence* in this manner, *viz.* 12*l.* multiplied by 20 (becauſe 20*s.* make one *l.*) produce 240 *Shillings*, unto which adding 13*s.* the ſum is 253 *Shillings.* Again, 253 *s.* multiplyed by 12 (becauſe 1 *ſhilling* is equal to 12 *Pence*) produce 3036 *Pence*, unto which if 10 *pence* be added the ſum is 3046 *pence*, as by the operation in the *Margin* is manifeſt.

l. *s.* *d.*
12--13--10.
20
240
add 13
253 *ſhillings.*
12
506
253
3036
add 10
3046 *pence.*

But after that general Method is well underſtood, the work of the laſt Example, and ſuch like may be contracted thus; *viz.* To convert 12 *pounds*, 13 *ſhillings*, 10 *pence*, all into *pence*, Firſt 12 multiplied by 0, (which ſtands in the Units place of 20) produceth 0, but inſtead of 0, I write down 3 under the line (to wit, the three that ſtands in the units place of the 13 *ſhillings in the ſum propounded*;) Then I proceed to multiply 12 by 2, ſaying twice 2 is 4, to which adding 1 for the ten in the ſaid 13 *ſhillings*) it makes 5, which I ſet on the left hand of 3 before written; Laſtly, twice 1 is 2, which I ſet on the left hand of 5; And ſo 12 *pounds*, 13 *ſhillings* and 10 *pence* are converted into 253 *ſhillings.*

l. *s.* *d.*
12--13--10.
20
253 *Shillings.*
12
516
253
3046 *pence.*

It remains to multiply the ſaid 253 by 12 (becauſe 12 *Pence* makes 1 *Shilling*) and to add 10 to the Product, which may be done thus; Firſt, twice 3 is 6, to which adding 10 (to wit, 10 *Pence* in the Sum firſt propounded) it makes 16, wherefore (according to the *Rule* of Multiplication) I ſet 6 under the line, and keep one in mind; Again, twice 5 with 1 in mind making 11, I write down 1, and keep 1 in mind; likewiſe twice 2 and 1 in mind making 5, I write down 5; Then 253 multiplied by 1 makes 253, which I ſet orderly under 516; Laſtly, thoſe two Products added together make 3046, which is the number of *Pence* contained in 12*l.*—13*s.*—10*d.* as before was found out by the general method.

So 35 *Ounces*, 16 *Penny Weights, and* 12 *Grains Troy* will be reduced into 17196 *Grains.*

VII. Reduction aſcending is performed by Diviſion, for if the number of Integers given be divided by ſuch a number of the ſame Integers, as are equal to one of the Integers required, the *Quotient* is the number of Integers ſought.

Reduction aſcending is performed by Diviſion.

So 220800 *Farthings* being divided by 4 (the number of *Farthings* in a *Penny*) give 55200 *Pence* in the *Quotient*; In like manner if 55200 *Pence* be divided by 12 (the number of *Pence* in a *Shilling*) the *Quotient* is 4600 *Shillings.* Laſtly, 4600 *Shillings* being divided by 20 (becauſe 20 *s.* make a *Pound ſterling*) the *quotient* is 230 *Pounds ſterling*) which are equal to 220800 *Farthings* firſt given. The operation is as followeth.

4)

```
                12)      20)
4) 220800(55200 (4600|0 (230l.
   .....   ....
            48
           ----
            72
            72
           ----
            00
```

In like manner, 34268 *Grains Troy* will be reduced to 5*l.* 11 *Ounces*, 7 *Penny Weight*, and 20 *Grains*. This kind of *Reduction* may be made the eaſier to the Learner by the following *Tables*.

1. *Of Engliſh Money.*

Farthings	*Divi. by*	4	*give*	*Pence.*
Pence		12		*Shillings.*
Shillings.		20		*Pounds.*

2. *Of Troy Weights.*

Grains	*Divi. by*	24	*give*	*Penny Weights.*
Penny Weight		20		*Ounces.*
Ounces		12		*Pounds Troy.*

Alſo in *Apothecaries Weights.*

Grains	*Divi. by*	20	*give*	*Scruples.*
Scruples		3		*Drams.*
Drams.		8		*Ounces Troy.*

3. *Of Averdupois Weight,*

Drams	*Divided by*	16	*give*	*Ounces.*
Ounces		16		*Pounds.*
Pounds		28		*Quarters of C.*
Quarters		4		*Hund. Weight.*

3. *Of Liquid Meaſures.*

	Divided by	*give*
Pints	2	*Quarts.*
Quarts	2	*Pottles.*
Pottles	2	*Gallons.*
Gallons	63	*Hogſheads.*

5. *Of Dry Meaſures.*

	Divided by	*give*
Pints	2	*Quarts.*
Quarts	2	*Pottles.*
Pottles	2	*Gallons.*
Gallons	2	*Pecks.*
Pecks	4	*Buſhels.*
Buſhels	8	*Quarters.*

6. *Of Long Meaſures.*

	Divided by	*give*
Barly Corns	3	*Inches.*
Inches	12	*Feet.*
Feet	3	*Yards.*
Yards	220	*Furlongs.*
Furlongs	8	*Engliſh Miles.*

Alſo,

	Divi. by	*give*
Nails	4	*Quarters of Yards.*
	4	*Yards.*
Quarters	5	*Ells Engliſh.*

Of Superficial Meaſures of Land.

	Divi. by	*give*
Perches or Poles	40	*Roods or Quarters of Acres.*
Roods	4	*Acres.*

8. *Of Time.*

	Divi. by	*give*
Minutes	60	*Hours.*
Hours	24	*Days.*
Days	7	*Weeks.*

Note, that if after Diviſion is finiſht in Reduction aſcending there be any remainder, it is of the ſame denomination with the Dividend.

Note alſo that Reduction deſcending and aſcending do mutually prove one another, by inverting the queſtion; for as in 56 *Pounds* ſterling, there will be found 53760 *Farthings*, by Reduction deſcending; So for Proof thereof, 53760 *Farthings* will be reduced to 56 *Pounds*, by Reduction aſcending.

Queſtions to exerciſe Reduction.

1. In 257*l.* how many ſhillings? *Anſwer*, 5140.

2. In 3076*l.* how many ſhillings? *Anſwer*, 61520.

3. In 902 ſhillings how many pence? *An.* 10824.

4. In 2179 ſhillings how many farthings? *Anſwer*, 104592.

5. In 49*l.*—13*s.*—7*d.* how many pence? *Anſwer*, 11923.

6. In 2053*l.*—14*s*—9*d.*—2*f.* how many farthings? *Anſw.* 1971590.

7. In 354*lb.* of Troy weight how many grains (of Gold-ſmiths weight?) *Anſwer*, 2039040.

8. In 300 Engliſh miles how many yards? *Anſwer*, 528000.

9. In 1 Engliſh mile, how many barley corns length? *Anſw.* 190080.

10. In 560 Acres how many Perches? *Anſwer*, 89600.

11 In 225 Acres, 3 Roods, and 30 Perches, how many Perches? *Anſw.* 36150.

12. In 11923 pence how many pounds? *Anſwer*, 49*l.*—13*s.*—7*d.*

13. In 5764684 farthings how many pounds? *Anſw.* 6004*l.*—17*s.*—7*d.*

14. In 234678 Perches, how many Acres? *Anſwer*, 1466 Acres, 2 Roods, and 38 Perches.

15. In 525960 minutes of an hour, how many days? *Anſw.* 365 days and 6 hours (or 1 year very near.)

16. In 10080 Pints, how many Hogſheads? *Anſw.* 20.

17. In 34678 grains of Apothecaries weight how many ounces Troy? *Anſw.* 72 Ounces, 1 Dram, 2 Scruples, and 18 Grains.

18. In 106735 Pints of wheat, how many Quarters? *Anſw.* 208 Quarters, 3 Buſhels, 2 Pecks, 1 Gallon, 1 Pottle, 1 Quart, 1 Pint.

19. In 3969301 Barley corns length, how many Miles? *Anſw.* 20 Miles, 7 Furlongs, 12 Yards, 2 Feet, 4 Inches, and 1 Barley corns length.

20. In 1900800 Barley corns length, how many Miles? *Anſ.* 10.

C H A P.

CHAP. VIII.

Of the Rule of Three Direct.

I. THE Rule of Three is ſo called, becauſe by three Numbers known or given, it teacheth to find a fourth unknown; it is alſo called the Golden Rule for the excellency thereof; Laſtly, it is called the Rule of proportion for the reaſon hereafter declared.

II. The Rule of Three is either ſingle or compound.

III. The ſingle Rule is, when three terms or numbers are propounded, and a fourth proportional unto them is demanded.

The Rule of Three.

IV. Four Numbers are ſaid to be proportionals, when the firſt containeth the ſecond, or is contained by the ſecond in the ſame manner as the third containeth the fourth, or is contained by the fourth: ſo theſe 4 Numbers are ſaid to be Proportionals, 8, 4, 12, 6, for as 8 containeth 4 twice, ſo doth 12 contain 6 twice, and therefore 8 is ſaid to have ſuch proportion to 4 as 12 hath to 6; likewiſe theſe are Proportionals, 4, 8, 6, 12. For as 4 is the half of 8, ſo is 6 the half of 12; and therefore 4 is ſaid to have ſuch proportion to 8 as 6 hath to 12.

V. The Terms or Numbers of the Rule of Three (to wit, the three Numbers given, and the fourth ſought) conſiſt of two different denominations, *viz.* two of the three given terms have one name, and the other given term with the term required

The divers denominations of the terms in the Rule of Three.

required have another: ſo this queſtion being demanded, if four Students ſpend 19 Pounds in certain Months, how much Money will ſerve 8 Students for the ſame time, and at the ſame rate of Expence? Here Students and Pounds are the two denominations of the terms in the queſtion, *viz.* 4 and 8 (being two of the terms propounded) have the denomination of Students, and 19 the other term given, together with the term required, have the denomination of Pounds.

VI. In the Rule of Three, two of the three given terms imply a ſuppoſition, and the third moves a queſtion: ſo in the aforementioned queſtion a ſuppoſition is made, that 4 Students ſpend 19 Pounds, and a queſtion is moved with the number 8, to wit, how many Pounds will 8 Students ſpend.

The right ordering of the terms given.

VII. In the Rule of Three, the numbers given muſt be ſo ranked, that the known number, or term upon which the queſtion is moved, muſt poſſeſs the third place in the Rule; alſo of the other two that which hath the ſame denomination with the third, muſt be in the firſt place: laſtly, the other known term, which is of the ſame denomination with the fourth term ſought (or anſwer of the queſtion) muſt poſſeſs the ſecond place: ſo in the queſtion before mentioned, the terms 4, 19, and 8, are to be thus placed, *viz.* 8 is the term upon which the queſtion is moved, and therefore to poſſeſs the third place in the Rule; 4 is of the ſame denomination with 8 *viz.* of Students, and therefore to be in the firſt place; Laſtly, 19 being of the ſame denomination with the term ſought, *viz.* of Money, is to be in the ſecond place: and ſo they will be placed in the Rule thus,

Students.	*Pounds.*	*Students.*
If 4 ——	—— 19 ——	—— 8

That is to ſay, if 4 Students ſpend 19 Pounds, what will 8 Students ſpend? And here for the better diſcerning of the term or number upon which the queſtion is moved, you may obſerve, that for the moſt part it is the known number in the queſtion which immediately followeth theſe or ſuch like words; *viz. How many? How much*? *What will? How long? How far?* &c.

VIII. The Rule of Three is either Direct or Inverſe.

IX. The Rule of Three Direct is, when the ſenſe or tenour of the queſtion requireth that the fourth number ſought muſt have ſuch proportion to the ſecond, as the third number hath to the firſt; ſo in the aforementioned queſtion, if 4 Students ſpend 19 Pounds, how many Pounds will 8 Students ſpend at the ſame rate of expence? It is evident that the thing required is to find a number which may have ſuch proportion to 19, as 8 hath to 4; that is, as 8 is the double of 4, ſo ought the fourth number to be the double of 19; for if 19 Pounds be required to maintain 4 Students a certain time, as much more muſt needs be required for the maintainance of 8 Students the ſame time, and therefore in this caſe we may ſay in a direct proportion, as 4 is to 8, ſo is 19 to a number which ought to be as much more as 19.

The Rule of Three Direct.

How to work the Rule of Three Direct, the three given terms being ſingle numbers.

X. In the direct Rule of Three, if you multiply the ſecond term by the third, or (which is all one) the third term by the ſecond, and then divide the Product by the firſt, the *Quotient* will give the fourth term or fourth proportional required: ſo in the queſtion before propounded, if you multiply 19 by 8, the product is 152, which if you divide by 4, the Quotient will give you 38 the fourth term demanded, and the work will ſtand thus.

```
   Stud. l, Stud.  l.
If 4---19-- 8--(38
        8
 4)152(38 pounds
   12
   ---
    32
    32
    --
     0
```

A ſecond Example may be this, if 8 Yards coſt 9 Pounds how much will 3 Yards coſt? *Anſwer*, 3 *l.*———7 *s.*——6 *d.*

This Queſtion being ſtated according to the ſeventh Rule of this Chapter, will ſtand as here you ſee; then multiplying (as before) the ſecond term 9 by the third term 3, the Product is 27, which being divided by the firſt term 8, the Quotient is 3 Pounds, and there is a remainder of three Pounds, which muſt be reduced into ſixty Shillings, and after thoſe Shillings are divided by 8, and the reſt of the work proſecuted according to the

```
y.  l.  y.  l. s. d.
8-- 9-- 3  (3:7:6.
        3
8)27(3 pounds
  24
  --
   3 the remaind.
  20
8)60(7 ſhillings.
  56
  --
   4 the remaind.
  12
8)48(6 pence.
```

Note at the latter end of the 11th Rule of the 6th Chapter, at length the entire quotient or anſwer of the queſtion is 3*l.*——7*s.*——6*d.*

A third Example, if 51 ounces of Silver plate be ſold for 13 Pound ſterling, what is the price of 1 Ounce of that Plate? *Anſ.* 5 *s.* — 1 *d.* and ſomewhat more. The operation is thus: After the three known terms of this queſtion are rightly ordered they will ſtand as here you ſee in the Example; then multiplying the ſecond term 13 by the third term 1, the Product will be alſo 13 (for multiplication by one makes no alteration;) which 13 being divided by 51, after the manner of operation delivered in the note upon the 5th Rule of the 7th Chapter, the entire Quotient or Anſwer of the Queſtion will at length be found to be 5 *s.* — 1 *d.* and ſomewhat more, but the ſurpluſage being leſs than a farthing is omitted as uſeleſs.

```
oz.   l.   oz.
51 —— 13 —— 1
            1
   ___________
           13
           20
   ___________
   51) 260 (5 ſhillings
       255
   ___________
         5
        12
   ___________
   51) 60 (1 penny.
       51
   ___________
        9
```

Example 4. What muſt be paid to a Labourer for his Wages for 27 Weeks at the rate of 4 *s.* for 1 Week? *Anſwer*, 5 *l.*——8 *s.*

After the three given terms are rightly placed in the Rule, they will ſtand as you ſee in the Example; then multiplying the third term 27 by the ſecond term 4, the product is 108, which I ſhould divide by the firſt term 1, but in regard

```
Week. Shil. Week.
1 —— 4 —— 27
           4
         ___
         108
```

diviſion

diviſion by 1 makes no alteration, the *Quotient* is alſo 108, ſo that the fourth term ſought is 108 ſhillings, which being reduced to Pounds, according to the ſeventh Rule of the ſeventh Chapter, give 5 *l.* 8 *s.* for the anſwer of the queſtion.

To prepare the terms of the Rule of Three, when they are compounded of divers denominations.

XI. In the Rule of Three, if after the queſtion is ſtated according to the ſeventh Rule of this Chapter, any of the three given terms be a compound term conſiſting of divers denominations, as Pounds, Shillings, and Pence; or Weeks Days, Hours, &c. ſuch compound term muſt firſt be reduced into the loweſt of thoſe denominations (by the 6th Rule of the ſeventh Chapter) to the end that the three given terms may be three ſingle numbers; alſo of theſe three ſingle numbers the firſt and third muſt always be of one and the ſame denomination: for if it happen that they expreſs things of different names, ſuch of the two which hath the greater name (or denomination) is to be reduced into the ſame name with the leſſer (by the 5th Rule of the ſeventh Chapter:) Theſe preparations being obſerved, the reſt of the work is to be proſecuted according to the tenth Rule of this Chapter. *Example,* What will 48 Ounces, 17 Peny-weight, and 20 Grains of Silver-Plate amount unto at the rate of 5 *s.*——6 *d.* the Ounce? *Anſwer,* 13 *l.*——8 *s.*——10 *d.*——3 *f.* very near.

This

oz.	*s.*	*d.*	*oz.*	*p. w.*	*gr.*
1	5	6	48	17	20
20	12		20		
20	66		977		
24			24		
480			3928		
			1954		
			23468 *Grains.*		

This queſtion being ſtated according to the ſeventh Rule of this Chapter, will ſtand in the Rule as you ſee in the Example, to wit, if 1 Ounce coſt 5 *s.*—6 *d.* what will 48 *oz.*—17 *p. w.*—20 *gr.* coſt? Here becauſe the third term is compounded of divers denominations, it muſt be reduced into the loweſt of thoſe denominations, to wit, grains; ſo by the ſixth Rule of the ſeventh Chapter, there will be found 23468 grains for the third term: likewiſe becauſe the ſecond term 5 *s.*—6 *d.* is a compound term, whoſe loweſt name is Pence, it muſt be reduced into Pence (by the aforeſaid Rule;) ſo there will be found 66 Pence for the ſecond term: Moreover becauſe the firſt term hath the name Ounce, and the third term the name Grain, the firſt term 1 Ounce muſt be converted into 480 Grains (which are equal to 1 Ounce;) then will the three terms or ſingle numbers ſtand in the Rule, as here you ſee *viz.* if 480 Grains coſt 66 Pence, how many Pence will 23468 Grains coſt?

gr.	*pence.*	*gr.*
480	66	23468

Now proceeding according to the tenth Rule of this Chapter, there will ariſe in the Quotient 3226 Pence, beſides a remainder of 408, which being reduced to 1632 Farthings, and thoſe divided by the firſt term 480 the quotient will be 3 Farthings, ſo that the entire quotient is 3226 Pence, 3 Farthings, and ſomewhat more (but the parts of a Farthing being of no moment, may be neglected.) Laſtly, the ſaid 3226 Pence being reduced according to the ſeventh Rule of the ſeventh Chapter, give 13*l.*——8*s.*——10*d.*——3*f.* ſo that 13*l.*——8*s.*——10*d.*——3*f.* and ſomewhat more, will be the Anſwer of the Queſtion.

The proof of the Rule of Three direct

XII. For the proof of the Direct Rule of Three, multiply the fourth term by the firſt, which done, if that Product be equal to the Product of the ſecond term multiplied by the third, the work is right, otherwiſe it is erroneous: ſo in the firſt Example, 38 the fourth term, being multiplied by the firſt term 4, the Product is 152, which is alſo the Product of 19 multiplied by 8. But if it happen that after the fourth term, or anſwer of the queſtion is found in the ſame denomination with the ſecond term, there is yet a remainder, ſuch remainder muſt be added to the Product of the firſt term, multiplied by ſuch fourth term, and then the ſum muſt be equal to the Product of the ſecond and third terms (the ſecond term conſiſting of the ſame denomination with the fourth:) ſo in the laſt Example the fourth term is 3226, and there happens to be a remainder of 408, which being added to the Product of the Multiplication of the ſaid 3226 by the firſt term 480, gives 1548888, which is the ſame with the Product of the third term 23468 multiplied by the ſecond term 66, as will appear when work'd.

XIII. When the firſt of the three given numbers in the Rule of three Direct, is 1 or unity, the queſtion may oftentimes be anſwered more ſpeedily than by the Rule of Three, even by thoſe who have but little skill in Arithmetick, as will partly appear by the following Examples, *viz.*

A compendious operation in the Rule of three direct, when the firſt term is 1 or unity

1. At 17*s.*—9*d.* the yard, what will 84 yards coſt? *Anſwer,* 74*l.*—11*s.* For Reaſon ſheweth that 84 yards muſt (at the ſaid rate) coſt 84 Angels, 84 Crowns, 84 half Crowns, and 84 Three pences, all which being computed and added together, will give the full value of 84 yards, *Viz.*

	l.	*s.*	*d.*
84 Angels make	42	00	00
84 Crowns	21	00	00
84 half Crowns	10	10	00
84 Three Pences	1	01	00
Sum	74	11	00

2. At the rate of 9*s.* the Buſhel of Wheat, what will 51 Quarters amount unto? *Anſwer,* 183 *l.* 12*s.*—0*d.*

It is evident that the price of 1 Quarter (which conſiſts of 8 Buſhels) will be 8 Angels wanting 8 Shillings: therefore,

	l.	*s.*	*d.*
from 8 Angels, to wit,	4	00	00
ſubſtract	0	08	00
remains the price of 1 Quarter	3	12	00

Then the value of 51 Quarters, at the rate of 3*l.*——12*s.*——0*d.* the Quarter, may be found in manner following, *Viz.*

	l.	*s.*	*d.*
51 times 3*l.* or 3 times 51*l.* is	51	00	00
	51	00	00
	51	00	00
51 Angels make	25	10	00
51 Shillings doubled make	5	02	00
the price of 51 Quarters	183	12	00

3. What is a Cheſt of Sugar worth, that weigheth neat weight (the Tare being ſubtracted) 7 *C.* 3 *q.* 7 *lb.* at the rate of 6*l.*——3*s.*——4*d.* for 1 *C*? *Anſwer*, 48*l.*——3*s.*——6*d.*——2*f.*

Tare is that wherein any thing is put, as a Bag for Pepper, a Cheſt for Sugar.

	l.	s.	d.	f.
7 times 6 pounds make	42	00	00	
7 times 3 Shillings	1	01	00	
7 Groats	0	02	04	
The half of 6*l.*--3*s.*--4*d.* for 2 *qu.* is	3	01	08	
The half of 3*l.*--1*s.*--8*d.* for 1 *qu.* is	1	10	10	
The fourth part of 1*l.*—10*s.*—10*d.* (becauſe 7 *l.* is a fourth part of 28*l.* or of 1 *qu.*) is	0	07	08	2
	48	03	06	2

Practical Rules of this nature cannot be compleatly underſtood without ſome skill in fractions, as will hereafter appear in the ſecond Chapter of the Appendix: And therefore I ſhall conclude this Chapter with the following Queſtions, whoſe Anſwers are annexed to them, and may be found out by the preceding Rules; but the operations are purpoſely omitted, and left as an exerciſe for the Learner.

Queſtions to exerciſe the Rule of Three direct.

1. If 17 yards of Cloth coſt 19*l.* 2*s.* 6*d.* what will 35 yards coſt at that rate; *Anſwer*, 39*l.* 7*s*, 6*d.*

2. If 35 yards coſt 39*l.* 7*s.* 6*d.* how many yards may be bought at that rate for 19*l.* 2*s.* 6*d*? *Anſwer*, 17 yards.

3. If 35 yards coſt 39*l.* 7*s.* 6*d.* what are 17 yards worth at that rate? *Anſwer*, 19*l.* 2*s.* 6*d.*

4. If 17 yards be ſold for 19*l.* 2*s.* 6*d.* how many yards will 39*l.* 7*s.* 6*d.* buy at that rate? *Anſw.* 35 yards.

5. What muſt I pay for the carriage of 17 hundred weight, 3 quarters and 11 pounds *Averdupois*, at the rate of 7 ſhillings the hundred weight? *Anſw. 6l.—4s.—11d.—1farth.*

6. If *6l.— 4s.—*11.—1 *f.* be pay'd for the carriage of 17 hundred weight, 3 quarters, and 11 pounds, what was paid for the carriage of 1 pound weight? *Anſw.* 3 Farthings.

7. What muſt I pay for 39 ounces, 7 penny weight, and 18 grains of white plate at the rate of 5*s.* and 5*d.* the ounce? *Anſ.* 10*l.*—13*s*,—4*d.* and three quarters of a farthing.

8. What muſt 1 *l.* (or 20 *s.*) pay towards a Tax, when 326*l.*—6*s.*—8*d.* is aſſeſſed at 41*l.*—16*s.*—2*d.*—3*f*? *Anſw.* 2 *s.*—6*d.*—2*f.*

9. What will the Intereſt of 876*l*—17*s.*—6*d.* amount unto for 1 year at the rate of 6*l.* for 100*l.* for the ſame time? *Anſw.* 52*l.*—12*s.*—3*d.*

10. If 3 yards in length of Engliſh meaſure be equal to 4 ells Flemiſh; how many Flemiſh ells are contained in 120 yards Engliſh? *Anſwer* 160 Flemiſh ells.

11. If 4 Flemiſh ells in length, be equal to 3 Engliſh yards; how many Engliſh yards in 300 Flemiſh ells? *Anſw.* 225 Engliſh yards.

12. If 3 ells in length of Engliſh meaſure, be equal to 5 Flemiſh ells, how many Flemiſh ells in 120 Engliſh ells? *Anſw.* 200 Flemiſh ells.

13. If 5 Flemiſh ells in length, be equal to 3 Engliſh ells; how many Engliſh ells in 145 Flemiſh ells? *Anſw.* 87 Engliſh ells.

14. If 3 ounces of Silk weight, be equal to 4 ounces of Venice weight; how many ounces Venice are equal to 60 ounces of Silk weight? *Anſwer* 80 ounces Venice.

15. A Merchant delivered at *London* 120*l.* ſterling, to receive 207*l* Flemiſh at *Amſterdam*; what was 1 *l.* ſterling valued at in Flemiſh money? *Anſw.* 1 *l.*—14 *s.*—6 *d.*

16. If a Bill of Exchange be accepted at *London*, for payment of 400*l.* ſterling, for the value deliver'd at *Amſterdam?* at 1 *l.*—13*s.*—6*d.* Flemiſh for 1*l.* ſterling how much Flemiſh money was deliver'd at *Amſterdam*? *Anſw.* 670 *l.* Flemiſh.

17. When the Exchange from *Antwerp* to *London* is at 1*l.*—4*s.*—7*d.* Flemiſh for 1*l.* ſterling; how much ſterling muſt I pay at *London* to receive 236*l.* Flemiſh at A*ntwerp*? *Anſw.* 192*l.* ſterling.

18. A Merchant deliver'd at *London* 370*l.* ſterling by Exchange for *Roan* at 74 *d.* ſterling for 50*s.* Tournois; how much Tournois ought he to receive at *Roan*? *Anſw.* 60000*s.* Tournois.

19. In 370 Ducats, at 4*s.*—2*d.* the Ducat; how many French Crowns at 6*s.*—2*d*? *Anſw.* 250 Crowns; For if 74*d.* give 1 Crown, 18500*d.* (or 370 Ducats) will give 250 Crowns.

20. In 516 Dollers, at 4*s.*—5*d.* the Doller; how many Guineas at 1 *l.*--1 *s.* 6 *d.* the peice? *Anſw.* 106 Guineas. For if 258 *d.* give 1 Guinea, 27348*d.* or 516 Dollers) will give 106 Guineas.

CHAP. IX.

Of the Inverſe Rule of Three.

1. THE Rule of Three Inverſe is, when the fourth term required ought to proceed from the ſecond term, according to the ſame rate or proportion that the firſt proceeds from the third: ſo this queſtion being propounded, if 8 Horſes will be maintained 12 days with a certain quantity of Provender, how many days will the ſame quantity maintain 16 Horſes? Here, as 8 is half 16, ſo ought the fourth term required to be half 12; for if certain buſhels of Provender ſerve 8 Horſes 12 days; 16 Horſes will eat up as much Provender in half that time: and therefore you cannot ſay here in a direct proportion (as before in the Rule of Three direct) as 8 to 16, ſo is 12 to another number which ought to be in that caſe as great again as 12; but contrariwiſe by an *inverted proportion*; beginning with the laſt term firſt, as 16 is to 8, ſo is 12 to another number, which ought to be in this caſe half 12. And by the due obſervation of this definition, together with that of the Rule of Three Direct (propounded in the ninth Rule of the eighth Chapter) when any queſtion belonging to the ſingle Rule of Three is propounded, you may readily diſcern by which of thoſe Rules it ought to be reſolved; for if the three terms given look for a fourth

horſes	*days*	*horſes*
8 ——	12 ——	16

fourth in a direct proportion as they ſtand ranked in the Rule, you muſt reſolve the queſtion by the direct Rule; contrariwiſe when the proportion is inverted or turned backwards, it ought to be reſolv'd by the Inverſe Rule of Three, which here followeth.

How to work the Inverſe Rule of Three.

II. In the inverſe Rule of Three, after the three given terms are rightly placed in the Rule, and reduced (if there be need) according to the eleventh Rule of the eighth Chapter, multiply the firſt term by the ſecond, or (which is the ſame) the ſecond term by the firſt, and then divide the Product by the third term, ſo the *quotient* will give you the fourth term required, or anſwer of the queſtion; thus in the queſtion premiſed in the laſt Rule, if you multiply 12 by 8, the Product is 96, which if you divide by 16 the *Quotient* gives you 6, the fourth term required, as by the ſubſequent operation is manifeſt.

horſes *days* *horſes* *days*
8——12——16——(6
8
————
16) 96 (6
96
————
0

III. For the more ready diſcovering, whether a queſtion propounded belongs to the Rule of Three Direct, or to the Rule Inverſe, obſerve the directions following, *viz.* 1, By the ſenſe and tenour of the queſtion conſider whether more be

How to diſcern whether a queſtion in the Rule of Three is to be reſolved by the Rule Direct, or by the Rule Inverſe.

required or leſs; that is, whether the number ſought muſt be greater or leſs than the ſecond term. Secondly, eſteeming the firſt and third terms as extreams in reſpect of the ſecond, this will be a general Rule; namely, When more is required, the leſſer extream is the *Diviſor*; but when leſs is required, the greater extream is the *Diviſor*. Laſtly, the *Diviſor* being found out, it will be apparent whether the Rule be Direct or Inverſe, for when the *Diviſor* is the firſt term, it is a Rule Direct; but when the *Diviſor* is the third term, the Rule is Inverſe. Another Example of the Rule inverſe may be this: If 12 Mowers do mow certain Acres in 4 days, in what time will 23 Mowers perform the ſame work? *Anſwer* 2 days, 2 hours, and ſomewhat more. Here, the 3 known terms being rightly placed in the Rule, will ſtand as you ſee in the Example; and ſince it is evident that 23 men will require leſs time than 12 men to finiſh the ſame work, therefore (by the Rule aforegoing) the greater of the two extream Numbers 23 and 12 muſt be the *Diviſor*; and becauſe the *Diviſor* 23 ſtands in the third place, this queſtion is to be wrought by the Rule Inverſe; wherefore multiplying the firſt term 12 by the ſecond term 4, the product is 48, which being divided by the firſt term 23, the *Quotient* gives 2 days, and there is a remainder

```
     M.    D.    M.

     12————4————23
           4
   ————————————————
   23) 48 (2 days
       46
   ————————————————
        2
       24
   ————————————————
   23) 48 (2 hours.
       46
   ————————————————
        2
```

mainder of 2 days, which being reduced to hours, and thoſe divided by 23, the *Quotient* will be 2 hours, and there is yet a remainder of 2 hours to be ſubdivided into 23 parts if you pleaſe; ſo that the fourth term ſought, or anſwer of the queſtion is 2 days, 2 hours and ſomewhat more.

Again, take this for a third Example, If I lend my Friend 356 pounds for one year and 35 days (the year being ſuppoſed to conſiſt of 365 days) how long time ought he to lend me 500 pounds to requite my courteſy? *Anſwer*, 284 days and ſomewhat more, there being a remainder, to wit. 400, after the Diviſion is finiſh'd, as by the ſubſequent operation is manifeſt.

l. *y.* *D.* *l.*
356——1 : 35——500

365
add 35

multiply { 400
356

5|00)1424|00 (284 *days.*

IV. The proof of the Inverſe Rule of Three is this, multiply the third term by the fourth, then if this Product be equal to the Product of the firſt term multiplied by the ſecond, the work s true, otherwiſe erroneous; ſo in the Example of the ſecond Rule, the Product of 16 and 6 equal to the Product of 8 and 12. But if it

The proof of the Rule of Three Inverſe.

happen that after the fourth term, for anſwer of the queſtion, is found in the ſame denomination with the ſecond term, there is yet a remainder, ſuch remainder muſt be added to the Product of the third term multiplied by the fourth, and then the ſum muſt be equal to the Product of the firſt and ſecond terms (ſuch ſecond term being of the ſame particular denomination with the fourth:) ſo in the laſt Example, the fourth term is 284 days, and there remains 400 after the diviſion is finiſh'd, this 400 being added to the Product of the Multiplication of the third term 500 by the fourth term 284, gives 142400, which is equal to the Product of the firſt term 356, multiplied by the ſecond term 400 days.

CHAP. X.

The double Golden Rule Direct, performed by two ſingle Rules.

I. THE Compound Golden Rule is, when more than 3 terms are propounded.

II. Under the Compound Golden Rule, is comprehended the double Golden Rule, and divers Rules of plural proportion.

The double Golden Rule.

III. The double Golden Rule is, when five terms being propounded, a ſixth proportional unto them is demanded: as in this queſtion, If 4 Students ſpend 19 pound in 3 months, how much will ſerve 8 Students 9 months? Or this, if 9 Buſhels of Provender ſerve 8 Horſes 12 days, how many days will 24 Buſhels laſt 16 Horſes?

The parts into which the terms of the ſame Rule are diſtributed.

IV. The five terms given in this Rule conſiſt of two parts, *Viz.* A ſuppoſition expreſſed in the three firſt terms; and a demand propounded in the two laſt: So in the firſt Example of the laſt Rule, this Clauſe (if four Students ſpend 19 pounds in 3 months) is the ſuppoſition, and this (how much will ſerve eight Students nine months) is the demand: likewiſe in the other Example of the ſame Rule, this clauſe (if nine Buſhels of Provender ſerve 8 Horſes 12 days) is the ſuppoſition, and this (How long, or how many days will 24 Buſhels laſt 16 Horſes) is the demand propounded.

The right ordering of the terms.

V Here for ranking the terms propounded in their due order, firſt obſerve amongſt the terms of ſuppoſition, which of them hath the ſame denomination with the term required; then reſerving that term for the ſecond place, write the other two terms of ſuppoſition one above another in the firſt place; And laſtly, the terms of demand likewiſe one above another in the third place of the Rule, in ſuch ſort that the uppermoſt may have the ſame denomination with the uppermoſt of thoſe in the firſt place. Example, If 4 Students ſpend 19 pounds in 3 months, how much will ſerve 8 Students 9 months? Here the three terms of ſuppoſition are 4, 19, and 3, and of theſe terms 19 hath the ſame denomination with the term required, *viz.* of Pounds, for you are to enquire how much Money is requiſite for the maintenance of 8 Students 9 months, wherefore reſerving 19 for the ſecond place I write 4 and 3 one above another thus; then drawing a line upon the right hand of 4, I write 19 in the ſecond place; this done the work will ſtand as in the Margin; Laſt of all the terms of demand being 8 and 9, and 8 having the denomination of Students, I place it in the ſame line with 4 and 19, and write 9 under

4——19
3

under it; all this performed, the terms in this queſtion rank themſelves as followeth:

Viz. *Thus,*

4——19——8
3 9

Or Thus,

3——19——9
4 8

In like manner, if the ſecond queſtion of the third Rule of this Chapter were propounded; the terms thereof ought to be diſpoſed

Thus,

8——12——16
9 24

Or Thus,

9——12——24
8 16

VI. Queſtions belonging to the double Golden Rule may be reſolved by two ſingle Rules of Three, or by the Golden Rule Compound of five Numbers.

The Proportions of the double Golden Rule, when it is performed by two ſingle Rules.

VII. When Queſtions of this nature are reſolved by two ſingle Rules, the proportions are as followeth:

I. As the uppermoſt term of the firſt place, is to the middle term; So is the uppermoſt term of the laſt place to a fourth Number:

II. As the lower term of the firſt place is to that fourth number; ſo is the lower term of the laſt place to the term required.

So in this Example before recited,

4—19—8 uſing tacitly the lower term of the
3   9 firſt place as a common number in the firſt proportion, *ſay*, *thus*,

I. If 4 Student ſpend 19 pounds (in three months) what will ſerve 8 Students the ſame time?

Or thus, If 4 Students ſpend 19 pounds, what will 8 ſpend?

Which Rule of Three will be diſcovered to be direct (by the third Rule of the ninth Chapter?) therefore the fourth proportional proceeding from the ſaid three given numbers 4, 19, and 8 is 38 (by the 10th Rule of the 8th Chap. aforegoing.) Again to find the term required, uſing tacitely the uppermoſt term of the third place as a common Number in this laſt proportion, *ſay as followeth.*

II. If in three months 38 pounds are ſpent (by 8 Students) how much will ſerve them for 9 months?

Or thus, If 3 give 38, what will 9 yield?

Which Rule of Three will likewiſe be diſcovered to be direct (by the third Rule of the 9th Chapter;) therefore the fourth proportional proceeding from the ſaid 3 numbers, 3, 38, and 9, you ſhall likewiſe find (by the 10th Rule of the 8th Chap- before-recited) to be 114, for 38 being multiplied by 9, the Product is 342. which divided by 3, yields you in the *Quotient* 114; So that I conclude, if four Students ſpend nineteen pounds in three months, 114 pounds will ſerve 8 Students 9 months; as you may further obſerve by the Work following:

4——19——8

3 9——(114

4——19——8——(38	3——38——9——(114
8	9
4) 152 (38	3) 342 (114
12	3
32	04
32	3
0	12
	12
	0

In like manner if two ſingle Rules of Three be formed (according to the preceeding 7th Rule) out of the five numbers given in the laſt mentioned queſtion, the ſame being ranked according to the latter manner of ordering the ſaid numbers in the fifth Rule, each of the ſaid two Rules of three will be a Rule direct, and the ſame anſwer of the queſtion, to wit, 114 pounds will be diſcovered, as you may ſee by the ſubſequent operation.

3——19——9

4 8——(114

3——19——9——(57	4——57——8——(114
9	8
3) 171 (57	4) 456 (114
15	4
21	05
21	4
0	16
	16
	0

VIII. The double Golden Rule is either Direct or Inverſe.

IX. The Double Golden Rule Direct is, when both the ſingle Rules do each of them look for a fourth term in a direct proportion: As in the Example of the ſeventh Rule, where each of the two ſingle Rules of Three is a Rule Direct.

The double Golden Rule Direct.

For another Example take this, If the Carriage of 8 C. weight 128 miles, coſt 48 Shillings, for how much may I have 4 C. weight carried 32 miles after the ſame rate? The terms of this queſtion according to the fifth Rule of this Chapter, rank themſelves in *this order:*

128 ———— ———— 48 ———— ———— 32
8 4

Now taking tacitly the lower term of the firſt place as a common number, I form the firſt Rule of Three according to the ſeventh Rule, *ſaying*,

I. If the carriage of a certain weight (to wit, 8 C.) 128 miles, will coſt 48 Shillings, what will the carriage of the ſame weight 32 miles coſt?

Here it is eaſie to diſcern, that the fewer miles any weight is carried, the leſs money will pay for the carriage of that weight; therefore the fourth number ſought by the ſaid Rule of three muſt be leſs than the ſecond number 48: And foraſmuch as by the third Rule of the ninth Chapter, when leſs is required, the greater extream (whether it be the firſt or third number) muſt be the Diviſor; therefore the firſt number 128 is the Diviſor, and conſequently the Rule of Three above propounded is a Rule direct, wherefore finding out the fourth number by the tenth Rule of the eighth Chapter, to be 12 Shillings, I proceed to the ſecond proportion, and ſay,

II. If the carriage of 8 C. (32 miles) coſt 12 Shillings, how much muſt I give to have 4 C. carried the ſame diſtance?

And here likewiſe finding a fourth number to be looked for in a direct proportion, I diſcover that fourth, by the ſaid tenth Rule of the eighth Chapter, to be *6s.* which is the term demanded, and the anſwer to the queſtion propounded: So that at laſt I conclude, if the carriage of 8 C. 128 miles coſt 48*s.* the carriage of 4 C. 32 miles will coſt *6s.* according to the ſame rate: See the whole Work.

```
        128——48——32
         8         4—(6
---------------------------------
        128——48——32—(12
              32
              96
             144
        128) 1536 (12
             128
              256
              256
                0
---------------------------------
           8—12—4—(6
               4
         8)   48 (6
              48
               0
```

CHAP.

CHAP. XI.

The Double Golden Rule Inverſe, performed by two ſingle Rules.

THE Double Golden Rule Inverſe is, when one of the ſingle Rules looks for a fourth term in an inverted proportion: As in the laſt Example propounded in the fifth Rule of the laſt Chapter. For if you rank the terms of that queſtion, according to the ſaid fifth Rule, *thus*,

The double Golden Rule Inverſe.

8——12——16
9   24

And then work by two ſingle Rules of Three, formed according to the ſeventh Rule of the laſt Chapter, you ſhall find by the third Rule of the ninth Chapter, that the firſt of the ſaid two Rules of Three will be inverſe, and the latter direct; for ſaying firſt, if 8 Horſes be maintained 12 days (by 9 buſhels of Provender) how many days will 16 Horſes be kept by ſo much Provender? Here the anſwer 6 days will be found out by the Rule of Three inverſe: Secondly, ſaying, if 9 buſhels of Provender be eaten up (by 16 Horſes) in 6 days, in how many days will 24 buſhels be ſpent? Here the anſwer 16 days will be found out by the Rule of Three direct.

But if you order the given terms of the ſame queſtion, *thus*,

9——

9——12——24
8 16

And then work by two ſingle Rules of Three, formed according to the ſeventh Rule of the laſt Chapter, you ſhall find by the third Rule of the ninth Chapter, that the firſt of the ſaid two Rnles of Three will be Direct, and the latter Inverſe; for ſaying, firſt, If 9 buſhels of Provender will laſt 12 days (to maintain 8 Horſes) how many days will 24 buſhels ſerve the ſame number of Horſes: The anſwer 32 days will be found out by the Rule of Three direct. Secondly, ſaying, If 8 Horſes will be maintained 32 days (by 24 buſhels of Provender) how long will 16 Horſes be kept by the ſame quantity of Provender? Here the anſwer 16 days will be found out by the Rule of Three Inverſe.

Wherefore whenſoever a queſtion belonging to the double Rule of Three is ſevered into two ſingle Rules of Three (according to the preceding Rules) if one of them happens to be a Rule Inverſe, that double Rule is called the double Rule Inverſe.

Now the Reſolution of the queſtion propounded being ranked after the firſt manner, is as followeth.

8——12——16
9 24—(16

8——12——16—(6
8

16) 96 (6
96

0

9——6——24—(16
6

9) 144 (16
9

54
54

0

Again, The Reſolution of the ſame queſtion, being ranked after the laſt manner, *is this*,

9——

9——12——24
8 16—(16

9——12——24—(32
12

48
24

9) 288 (32
27

18
18

0

8——32——16—(16
8

16) 256 (16
16

96
96

0

So that at laſt I ſay if 9 Buſhels of Provender ſerve 8 Horſes 12 days, 24 Buſhels will laſt 16 Horſes 16 days, which is the reſolution of the Queſtion propounded.

CHAP.

CHAP. XII.

The Golden Rule compounded of five Numbers.

I. THE Golden Rule compound of five numbers is, when the terms being ranked, as before, inſtead of the double terms we uſe their products, and then proceed to find the term required by one ſingle Rule of Three.

The Golden Rule Compound of five numbers performed by one ſingle Rule direct.

II. Here when the Queſtion propounded ought to be performed by the double Rule direct, multiplying the terms of the firſt place, the one by the other, take their product for the firſt term, the middle number for the ſecond, and the product of the two laſt terms for the third term; this done having found by the Rule of Three direct, a fourth proportional unto thoſe three, that fourth term ſo found is the number you look for; ſo this queſtion being again propounded, if 4 Students ſpend 19*l.* in 3 months, how much will ſerve 8 Students 9 months? and the terms thereof being ranked as before, *viz. thus,*

4——19——8
3 9

The product of 4 multiplied by 3 is 12, and the product of 8 multiplied by 9 is 72; wherefore I *ſay*, As 12 to 19, ſo 72 to the term required, which I find by the ſingle Rule of Three direct to be 114, So that if 4 Students ſpend 19*l.* in three months, 114*l.* will be requiſite for the maintenance of 8 Students 9 months; ſee the whole operation, *as followeth.*

```
4——19——8
3          9—(114
——         ———
12         72
           19
           ———
           648
           72
           ———
        12)1368(114
           12
           ———
            16
            12
           ———
            48
            48
           ———
             0
```

In like manner this being the Queſtion as before in the laſt Rule of the tenth Chapter) if the car-iage of 8 C. 128 miles, coſt 48*s.* What will the arriage of 4 C. 32 miles ſtand me in? the Anſwer hereunto will be 6*s.* as appears by the Work.

```
128——)
```

128——48——32
8 128 4

1024 384 128
96
48

1024)6144(6 *Shillings*
6144

0

The Golden Rule compound of five Numbers performed by one ſingle Rule Direct or Inverſe.

III. When the Queſtion propounded ought to be reſolved by the double Rule Inverſe, having multiplied the double terms acroſs, that is, the uppermoſt term of the firſt place by the lower of the laſt, and the uppermoſt of the laſt place by the lower of the firſt, write each product under the lower term by which it is produced: and then if the Inverſe proportion be found in the uppermoſt line, uſing thoſe products as ſingle terms, proceed to find the term required by the ſingle Rule of Three direct: But in caſe you find the Inverſe proportion in the lower line, perform the Work by the ſingle Rule of Three Inverſe.

So in the Example above mentioned, if 9 Buſhels of Provender ſerve 8 horſes 12 days, how long will 24 Buſhels laſt 16 Horſes? Here if you rank the terms *thus*, you ſhall find the Inverſe proportion in the firſt line, as is obſerved in the laſt Chapter: And therefore having ſubſcribed the products

8—12—16
9 24

products according to the direction given you in this Rule, I proceed to ſatisfie the demand of this queſtion by the ſingle Rule of Three direct, as appears by the Work following.

```
     8———12———16
     9          24—(16
   ———        ———
   144         192
                12
              ———
               384
              192
              ———
        144) 2304 (16
             144
             ———
              864
              864
              ———
                0
```

But the terms of this queſtion being ranked *thus*, the Inverſe proportion is found in the lower line, as you may obſerve likewiſe by the laſt Chapter: whereupon in this caſe to reſolve the Queſtion, I proceed by the ſingle Rule of Three Inverſe, as appears by the Work hereunto annexed: howſoever therefore you work the Queſtion, you ſhall find the term required to be 16; ſo that at laſt I conclude, as before in the laſt Chapter, If 9 buſhels of Provender ſerve 8 horſes 12 days, 24 buſhels will laſt 16 horſes 16 days.

```
9——12——24
8       16
```

9——12——24
8 16——(16

192 144
12

384
192

144) 2304 (16
144

864
864

0

CHAP. XIII.

The Rule of Fellowſhip.

Rules of plural Proportion.

I. THE Rules of plural Proportion are thoſe, by which we reſolve Queſtions, that are diſcoverable by more golden Rules than one, and yet cannot be performed by the double golden Rule mentioned before in the three laſt Chapters. Of theſe Rules there are divers kinds and varieties, according to the nature of the Queſtion propounded, for here the terms given are ſometimes four, ſometimes five, ſometimes more, and the terms required ſometimes more than one, *&c.*

II. Two particular Rules of plural proportion are theſe, the Rule of Fellowſhip and the Rule of Alligation.

The Rule of Fellowſhip.

III. The Rule of Fellowſhip is that, by which in accompts amongſt divers men (their ſeveral ſtocks together with the whole gain or loſs being propounded) the gain or loſs of each particular Man may be diſcovered: As in this Example, *A* and *B* were ſharers in a parcel of Merchandize, in the purchace of which, *A* laid out 7*l.* and *B* 11 *l.* and they having ſold this Commodity, find that their clear gains amount to 54*s.* Now here the Queſtion to be reſolved by this Rule is, what part of that 54*s.* accrews to *A*, and what to *B*, according to the rate of the ſeveral ſums or ſtocks which they adventured? Again, *A*, *B*, and *C*, freight a Ship from the *Canaries* for *England*, with 108 Tuns of Wine, of which *A* had 48, *B* 36, and *C* 24, the Mariners meeting with a ſtorm at Sea, were conſtrained for the ſafety of their lives, to caſt 45 Tun thereof over-board; here the Queſtion to be reſolved is, How many of the 45 Tun each particular Merchant hath loſt, according to the rate of his Adventure?

IV. The Rule of Fellowſhip is either ſingle or double.

V. The ſingle Rule is, when the ſtocks propounded do continue in the Adventure (or common Bank) equal times, to wit, one ſtock as long time as another.

How to work the ſingle Rule.

VI. In the ſingle Rule of Fellowſhip, take the total of all the ſtocks for the firſt term, the whole gain or loſs, for the ſecond, and the particular ſtocks for the third terms; this done, repeating the Rule of Three ſo often, as there are particular ſtocks in the Queſtion, the fourth terms produced upon thoſe ſeveral operations, are the reſpective gains or loſſes of thoſe particular ſtocks propounded: So in the firſt Example above-mentioned 7 *l.* and 11 *l.* are the ſtocks propounded, whoſe total is 18 *l.* which I take for the firſt term: Again, 54 *s.* the common gain, is the ſecond term, and 7 *l.* the firſt particular ſtock, is the third term of the firſt proportion; whereupon I ſay, as 18 *l.* to 54 *s.* ſo 7 *l.* to another number, which by the direct Rule of Three I find to be 21 *s.* *viz.* the part of the gain due to *A*, that expended the 7 *l.* ſtock. Then for the ſecond proportion, I ſay, as 18 *l.* to 54 *s.* ſo 11 *l.* to another number, which I likewiſe find by the Rule of Three direct to be 33 *s.* *viz.* the part of the gain due to *B*, for his 11 *l.* ſtock.

$$\left.\begin{matrix}7\\11\end{matrix}\right\}18\text{———}54\left\{\begin{matrix}7\text{———}21\\11\text{———}33\end{matrix}\right.$$

Again in the other premiſed Example, the particular loſs that happens to *A*, is 20 Tun. to *B* 15, and to *C* 10 Tun.

$$\left.\begin{matrix}48\\36\\24\end{matrix}\right\}108\text{———}45\text{———}\left\{\begin{matrix}48\text{———}20\\36\text{———}15\\24\text{———}10\end{matrix}\right.$$

VII. The double Rule of Fellowſhip

2. *Double.* is, when the ſtocks propounded are double numbers, *viz.* when each ſtock

hath relation to a particular time: Example, *A*, *B* and *C*, hold a Paſture in common, for which they pay 45 *l. per annum.* In this Paſture *A* had 24 Oxen went 32 days; *B* had 12 there 48 days, and *C* fed 16 Oxen there 24 days; now the Queſtion to be reſolved by this Rule is, what part each of theſe Tenants ought to pay of the 45 *l.* rent? and here you may *obſerve*, that the ſtocks propounded are *double* numbers, *viz.* each ſtock of Oxen hath reference to a *particular* time; for the reſpective ſtock of *A* is 24 Oxen, and its *particular* time is 32 days; again, the ſtock of *B* is 12 Oxen, and the *reſpective* time is 48 days: And laſtly, the ſtock of *C* is 16 Oxen, and its *peculiar* time is 24 days, which as you ſee are *double* numbers.

How to work the double Rule.

VIII. In the double Rule of Fellowſhip, multiply each particular ſtock by its reſpective time, and take the total of their Products for the firſt term, the whole gain or loſs for the ſecond, and the ſaid particular Products of the double numbers for the third term: This done, repeating, as before the Rule of Three, ſo often as there are Products of the double numbers; the fourth terms produced upon thoſe ſeveral operations, are the numbers you look for: So in the *Example* of the laſt Rule, the Product of 24 and 32 is 768, the Product of 12 and 48 is 576, and the Product of 16 and 24 is 384, the ſum of theſe Products is 1728, which is the firſt term in the Queſtion, then 45 *l.* the rent, is the ſecond term, and 768 the firſt Product, is the third term of the firſt proportion. Wherefore I ſay, as 1728 to 45 *l.* ſo 768 to another number, which I find by the direct Rule of *Three* to be 20*l.* *viz.* the part of the Rent that *A* ought to pay: Then for the ſecond proportion *I ſay*, as 1728 to 45 *l.* ſo 576 to 15 *l.* which is the part that *B* ought to *pay:* And laſtly, as 1728 to 45 *l.* ſo 384 to 10 *l.* *viz.* the part that *C* muſt pay.

$$\left.\begin{array}{l}768\\576\\384\end{array}\right\}1728\text{——}45\text{——}\left\{\begin{array}{l}768\text{—}20\\576\text{—}15\\384\text{—}10\end{array}\right.$$

A ſecond Example of the eighth Rule. Three Merchants, *A*, *B*, and *C* enter Partnerſhip, and agree to continue in a joint Adventure 16 months; *A* puts into the common ſtock at the beginning of the ſaid Term 100 pounds, at 8 months end he takes out 40 pounds, and 4 months after ſuch taking out he puts in 140 pounds. *B* puts in at firſt 200 pounds, at 6 months end he puts in 50 pounds more, and 4 months after the putting in of the 50 pounds, he takes out 100 pounds. *C* puts in at firſt 150 pounds, at 4 months end he takes out 50 pounds, and 8 months after ſuch taking out puts in 100 pounds. Now at the end of the ſaid 16 months, they had gained 357 pounds, the Queſtion is how much of the ſaid gain belongs to each Merchant for his ſhare.

In Queſtions of this Nature, two things are principally to be obſerved. 1. The whole time of Partnerſhip. 2. The reſpective time belonging to each Man's particular ſtock; ſo here, it is evident that the whole time is 16 months, and the particular ſtocks and times belonging to each Merchant will be as followeth, *viz.*

A Had

A had 100 *l.* in the common ſtock for 8 months, therefore 100 multiplied by 8 produceth	800
Alſo 60 *l.* for 4 months, therefore 60 multiplied by 4 produceth	240
Alſo 200 *l.* for 4 months, therefore 200 multiplied by 4 produceth	800
The Total of the Products of Money and Time for *A*, is	1840

B. had 200*l.* in the common ſtock for 6 months, therefore 200 multiplied by 6 produceth	1200
Alſo 250*l.* for 4 months, therefore 250 multiplied by 4, produceth	1000
Alſo 150*l.* for 6 months, therefore 150 multiplied by 6 produceth	900
The Total of the products of Money and Time for *B*, is	3100

C. had 150 *l.* in the common ſtock for 4 months, therefore 150 multiplied by 4 produceth	600
Alſo 100*l.* for 8 months, therefore 100 multiplied by 8 produceth	800
Alſo 200 *l.* for 4 months, therefore 200 multiplied by 4 produceth	800
The Total of the products of Money and Time for *C*, is	2200

Then adding the ſaid three totals together, to wit, 1840, 3100 and 2200, the ſum is 7140 wherefore proceeding as in the laſt Example, I ſay by the Rule of three direct, as 7140 is to the total gain 357 pounds; ſo is 1840 to 92 pounds the gain of *A*: again, As 7140 is to 357; ſo is 3100 to 155 the gain of *B*: Laſtly, as 7140 is to 357; ſo is 2200 to 110 the gain of *C*.

The Proof. *IX.* The Rule of Fellowſhip is proved by addition of the terms required, whoſe ſum ought to be equal to the ſecond term in the Queſtion, otherwiſe the whole work is erroneous: So in the firſt *Example* of the ſixth Rule afore-going, 21 *s.* and 33 *s.* being added together are equal to 54 *s.* the *ſecond term* in that Queſtion. Likewiſe in the laſt *Example* of the ſame Rule, as alſo in the firſt *Example* of the laſt Rule, the *ſum* of 20, 15, and 10, the terms required, are equal to 45, the *ſecond term* propounded.

CHAP. XIV.

The Rule of Alligation.

I. THE Rule of Alligation is that, by which we reſolve Queſtions, that concern the mixing of divers Simples together.

II. Alligation is either Medial or Alternate.

Alligation Medial. *III.* Alligation Medial is, when having the ſeveral quantities and rates of divers ſimples propounded, we diſcover the mean rate of a mixture compounded of thoſe ſimples. So 10 buſhels of Wheat at 4*s.* or (which is all one) 48*d.* the buſhel; 40 buſhels of Rye at 3*s.* or 36*d.* the buſhel; and 50 buſhels of Barley at 2*s.* or 24*d.* the buſhel; being mixed with 20 buſhels of Oats at 12*d.* the buſhel, the Rule of *Alligation medial* ſheweth you the mean price of that Miſtling.

The Operations and Proportions of the ſame Rule.

IV. In Alligation medial, firſt ſum the given quantities, then find the total value of all the ſimples: this done, the proportion will be as followeth.

As the ſum of the quantities is to the total value of the ſimples:
So is any part of the mixture propounded to the required mean rate or price of that part.

Repeating again the premiſed Example of the third Rule, I demand how much one buſhel of that Miſtling is worth? Now the ſum of 10, 40, 50, 20, (the given quantities) is 120 buſhels, and the value of the 10 Buſhels of Wheat at 48*d.* the buſhel amounts to 480*d.* for 48 being multiplied by 10, the product is 480: Again the value of the 40 buſhels of Rye at 36*d.* the buſhel, is 1440*d.* The value of the 50 buſhels of Barley at 24 *d.* the buſhel is 1200 *d.* And the value of 20 buſhels of Oats at 12*d.* the buſhel is 240*d.* All theſe values being added together, their total is 3360*d.* I ſay then by the Rule of *Three Direct*, if 120 buſhels give 3360 *d.* what will 1 buſhel yield? The Rule preſently anſwers me 28 *d.* whereupon I conclude, that a buſhel of that Miſtling may be afforded for 28*d.* that is, 2*s.* 4*d.* which is the reſolution of the *Queſtion propounded.*

120 —— 3360 —— 1 —— 28

In like manner if it be demanded what 8 Buſhels or a Quarter of that Miſtling is worth, the *Anſwer* will be 224*d.* which being divided by 12, and by that means reduced into *ſhillings* is 18*s.* 8*d.*

120——3360——8——224

The Proof. *V.* In *Alligation Medial*, the trial of the Work is by comparing the total value of the ſeveral Simples with the value of the whole mixture: For when thoſe ſums accord, the operation is perfect; So in the firſt Example of the laſt Rule.

		l.	*s.*	*d.*
The Value of	10 Buſhels of Wheat at 4*s.* the Buſhel is	2	0	0
	40 Buſhels of Rye at 3*s.* the Buſhel is	6	0	0
	50 Buſhels of Barley at 2*s.* the Buſhel is	5	0	0
	And 20 Buſhels of Oats at 12*d.* the Buſhel is	1	0	0
	All which amount to	14	0	0

which is likewiſe the value of 120 Buſhels at 28*d.* or 2*s.* 4*d.* the Buſhel, for that alſo amounts to 14*l.*

Alligation Alternate. *VI. Alligation Alternate* is, when having the ſeveral rates of divers Simples given, we diſcover ſuch quantities of them, as are neceſſary to make a mixture, which may bear a certain rate propounded.

Example: A Man being determined to mix 10 Buſhels of Wheat at 4*s.* or 48*d.* the Buſhel, with

Rye of 3*s.* or 36*d.* the Buſhel, with Barley of 2*s.* or 24*d.* the Buſhel, and with Oats of 1*s.* or 12 *d.* the Buſhel; the Rule of *Alligation Alternate* will diſcover unto you how much Rye, how much Barley, and how much Oats he ought to add unto the 10 Buſhels of Wheat; in ſuch ſort that the mixture of them altogether may bear a certain rate or price propounded.

The right ordering of the Terms.

VII. In Queſtions of *Alligation Alternate*, you muſt rank the terms in ſuch ſort, that the given rate of the mixture may repreſent the Root, and the ſeveral rates of the Simples may ſtand as Branches iſſuing from that Root: So the Example of the laſt Rule being propounded, I demand how much Rye, Barley and Oats, ought to be added to the 10 Buſhels of Wheat, that the mixture of all together may bear the rate or price of 28*d.* or 2*s.* 4*d.* the Buſhel: And therefore drawing a line of connexion, I place 28*d.* the given rate of the mixture, upon the left hand thereof, by it ſelf, repreſenting the *Root*, and likewiſe write the other rates propounded, *viz.* 48*d.* 36*d.* 24*d.* and 12*d.* one above another upon the right hand of that line of connexion, which rates are conceived to iſſue from 28*d.* as branches from the *Root*, the fabrick hereof appears plainly in the Margin.

$$28\left\{\begin{array}{l}48\\36\\24\\12\end{array}\right.$$

How to couple the Terms.

VIII. Having ranked the terms in their due order, link the branches together by certain Arches, in ſuch ſort, that one that is greater than the *Root* or rate of the mixture, may always be coupled with another that is leſs than the ſame: So in the premiſed Example, 48 may be linked with 12, and 36 with 24, or otherwiſe 48 may be coupled with 24, and 36 with 12, and then the Work will ſtand

Thus,



Or thus,



How to order the differences.

IX. Having alligated the branches, and found the differences betwixt them and the Root, write the differences of each branch juſt againſt his reſpective yoke-fellow. So the branches of the example aforegoing being linked after the firſt manner, and the difference between 28 and 48 (by the third or fourth Rule of the fourth Chapter of this Book) being 20, I place 20 juſt againſt 12, the reſpective yoke-fellow of 48. Again, 16, being the difference betwixt 28 and 12, I write it juſt againſt 48. In like manner 8 being the difference between 28 and 36, I place it right againſt 24. And laſtly, 4 the difference betwixt 28 and 24, I write juſt againſt 36: In the end the whole *Fabrick* of the Work (as the branches are thus linked) will ſtand as in the Example.



But the branches being linked after the other manner, the work will be thus diſpoſed:



For in this caſe 48 hath 24 for his yoke-fellow, and the reſpective *Comerado* of 36 is 12; and here the interchangeable placing of the *difference* (as in the premiſed Examples) is that which is more particularly termed *Alternation*.

X. When one branch is linked to divers other branches, and not to one alone, the *differences* ought to be as often tranſcribed, as it is ſo diverſly linked. So in the premiſed Example, you may (if you pleaſe) conceive 12 to be coupled both with 48 and 36; likewiſe 24 may be conceived to be linked with the ſame 48 and 36; wherefore the *difference* betwixt 28 and 12 being 16, I write it both juſt againſt 48 and 36; In like manner the *difference* between 28 and 24 being 4, I write it likewiſe over againſt the ſame numbers 48 and 36; Again, 20 being the *difference* betwixt 28 and 48, I place it juſt againſt 24 and 12; and 8 being the *difference* between 28 and 36 I write it likewiſe over againſt the ſame numbers 24 and 12; All this performed, the whole frame of the work will ſtand as in the *Margin*.



2. Take this for another Example: It is requi-

red to mix 10 Buſhels of Wheat at 48*d.* the buſhel with Rye of 36*d.* the buſhel, with Barley of 24*d.* the buſhel, and with Oats of 12*d.* the buſhel, and the queſtion now is, How much Rye, Barley, and Oats ought to be added to the 10 buſhels of Wheat, that the entire mixture may be afforded at 16*d.* the buſhel? Here the branches of this Queſtion (according to the eighth Rule of this Chapter) ought to be linked *thus*,



And as for the *Alternation* of the differences, it is evident (by the preſent Rule) that the difference between 16 and 12 being 4 ought to be thrice tranſcribed, *viz.* firſt juſt againſt 48, then againſt 36, and laſt of all againſt 24. Again, 32 the difference betwixt 16 and 48, as alſo 20 the difference between 16 and 36; and laſtly, 8 the difference betwixt 16 and 24, ought all to be placed juſt againſt 12.



3. I determining to mix 10 buſhels of Wheat at 48 *d.* the buſhel, with Rye of 36 *d.* the buſhel, with Barley of 24*d.* the buſhel, and with Oats

of 12 *d.* the Buſhel, deſire to know how much of each I ought to take, that I might afford the whole mixture at 40*d.* the buſhel: Here the whole Work being ordered according to the Rules aforegoing, it will ſtand as *followeth*,



4. A Man intending to mix 10 Buſhels of Wheat at 48 *d.* the Buſhel, with Rye of 36 *d.* the Buſhel with Barley of 24 *d.* the Buſhel, with Peaſe of 16*d.* the Buſhel, and with Oats of 12 *d.* the Buſhel, deſires to know how much Rye, Barley, Peaſe, and Oats he ought to add to the 10 Buſhels of Wheat, that the whole maſs of Corn ſo mixed might be afforded at 20 *d.* the Buſhel. This *Queſtion* being thus propounded, the terms thereof (by the Rules aforegoing) may be *Alligated* and the differences of the terms *Alternated*, as *followeth*,



5. Laſtly, A Goldſmith hath ſome Gold of 24 *Carects*, other of 21 *Carects*, and other ſome of 19 *Carects fine*, which he would ſo mix with *Alloy*, that 192 Ounces of the entire *mixture* might bear

17 *Careᴄts fine*, now the *Queſtion* is, how much of each ſort, as alſo how much *Alloy* he muſt take to accompliſh his deſire? Before you can well underſtand this *Queſtion*, it will be neceſſary to explain what a *Careᴄt fine*, and what *Alloy* is: The Mint-Maſters and Goldſmiths to diſtinguiſh the different *fineneſs* of Gold, eſteem an entire ounce to contain 24 *Careᴄts*, and one ounce of Gold that being tryed in the fire loſeth nothing of the weight, is ſaid to be 24 *Careᴄts fine*: Again, the ounce that being tryed loſeth one four and twentieth part of the weight, is ſaid to be 23 *Careᴄts fine*: In like manner that which loſeth two four and twentieth parts of the ounce, is eſteemed to be 22 *Careᴄts fine*, and ſo conſequently of the reſt: And as for the *Alloy*, it is Silver, Copper, or ſome other baſer Metal, with which the Goldſmiths uſe to mix their Gold, to the intent they may moderate, or abate the *fineneſs* thereof. Here you may alſo obſerve, that as the *fineneſs* of Gold is meaſured by *Careᴄts*, ſo is the *fineneſs* of *Silver* eſtimated by *ounces*: In ſuch ſort, that a *pound* of *Silver* which being tryed a certain time in the fire, loſeth nothing of the *weight*, is ſaid to be 12 *ounces fine*: But a *pound*, that being tryed loſeth ſomewhat of the *weight*, is ſaid to be the *remainder* of the *weight fine*. *Example*, a *pound* of *Silver*, that loſeth in the fire one *ounce* 8 *p*. is eſtimated to be 10 *ounces* 12 *p*. *fine*; and that which loſeth 2 *ounces*, 8 *p*. 10 *grains*, is ſaid to be 9 *ounces*, 11 *p*. 14 *grains fine*, &c. Now to rank the terms of the laſt mentioned *Queſtion*, as alſo the differences of the terms in their due order, becauſe the three given branches (*viz*. 24 *Careᴄts*,

What a Carect fine, and what Alloy is.

Carects, 21 *Carects*, and 19 *Carects*) are all greater than 17 *Carects* the *root* or rate of the *mixture*. I add o as another branch, which I conceive to be leſs than the *root*, and then proceed as in the former operations; the whole *frame* of the Work is expreſſed here, as followeth.



XI. When in one and the ſame line there are found more differences than one, add them together, and write the ſum juſt againſt the ſame differences before a ſtraight line drawn towards the right hand of the Work.

How to add the differences.

So the firſt Example of the laſt Rule being propounded, the ſum of 16 and 4 (the differences placed juſt againſt the firſt branch) being 20, I write it over againſt the ſame differences, before the new line drawn upon the right hand of the Work, and ſo conſequently the reſt in their due order, as appears by the Example hereunto annexed.



In like manner the laſt Example of the laſt Rule being offered, the whole Fabrick of the Work will ſtand as followeth:



XII. Alligation Alternate is, either Partial or Total.

Alternation Partial.

XIII. Alternation Partial is, when having the ſeveral rates of divers Simples, and the quantity of one of them given, we diſcover the ſeveral quantities of the reſt, in ſuch ſort that a mixture of thoſe Simples being made according to the quantity given, and the quantities ſo found, that mixture may bear a certain rate propounded: Of this kind is the Example of the ſixth Rule, as alſo all the Examples of the tenth Rule except the laſt.

The Proportions uſed in this Rule.

XIV. In Queſtions of Alternation Partial, the proportion is as followeth.

As the difference annexed to the firſt branch is to the ſeveral differences of the reſt:

So is the quantity propounded to the ſeveral quantities required.

So the Example of the ſixth and ſeventh Rules of this Chapter being again repeated, and the terms thereof, as alſo the differences of the terms being ordered after the firſt manner (ſhewed you in the ninth Rule aforegoing) it is evident that for every 16 Buſhels of Wheat that I take in the mixture, I ought to take 4 Buſhels of Rye, 8 Buſhels of Barley, and 20 Buſhels of Oats; and therefore *I ſay*,

The firſt Caſe.



I. As 16 the difference annexed to the firſt branch (being the rate of the Wheat) is to 4, the difference annexed to the next, being the rate of the Rye; ſo is 10 the given quantity of the Wheat to another number, which being found by the Rule of *Three direct*, to be two buſhels and an half (or two pecks) is the quantity of Rye neceſſary in the *mixture*.

II. As 16 to 8, ſo is 10 to another number; which being likewiſe found by the *Rule of Three* to be five buſhels, is the quantity of Barley neceſſary in the *mixture*.

III. As 16 to 20, ſo is 10 to another number, which being in like ſort found by the *Rule of Three* to be 12 buſhels, and half of a buſhel, is the quantity of Oats requiſite in the *mixture*.

So that at laſt I conclude, a heap of Corn being compoſed of 10 buſhels of Wheat, 2 buſhels and a half of Rye, 5 buſhels of Barley, and 12 buſhels and an half of Oats (when thoſe ſeveral Grains bear the *prices*, aforeſaid) may be afforded at 2*s.* 4*d.* the buſhel.

The ſame Example being ordered after the ſecond manner (expreſſed likewiſe in the 9th Rule of this preſent Chapter) *I ſay*, *2 Caſe.*

I. As 4 the *difference* annexed to the *rate* of the Wheat, is to 16 the *difference* annexed to the *rate* of the Rye; ſo is 10 the *given* quantity of the Wheat, to 40 buſhels the *required* quantity of the Rye.

II. As 4 to 20, ſo is 10 to 50 buſhels the requiſite quantity of the Barley.

III. As 4 to 8, ſo is 10 to 20 buſhels the quantity of the Oats *neceſſary* in the mixture.



So that I conclude *again*, a maſs of Corn being compounded of 10 buſhels of Wheat, 40 buſhels of Rye, 50 buſhels of Barley, and 20 buſhels of Oats, (when thoſe Grains bear the prices propounded in this Example) may be afforded at 2 *s.* 4 *d.* the buſhel as before.

3 *Caſe.* 3. That Example being diſpoſed after the third manner (expreſſed in the tenth and eleventh Rules of this Chapter) *I ſay*,

I. As 20 the *ſum* of the differences annexed to the *rate* of the Wheat, is to 20 the *ſum* of the differences annexed to the *rate* of the Rye; ſo is 10 the *given* quantity of the Wheat, to 10 buſhels the *required* quantity of the Rye.

II. As 20 to 28, ſo is 10 to 14 buſhels the *requiſite* quantity of the Barley.

III. As 20 to 28, ſo is 10 to 14 buſhels, the quantity of Oats *demanded* in the mixture.



16	4	20
16	4	20
20	8	28
20	8	28

Whereupon this *third time* likewiſe I conclude, that (thoſe Grains ſtill retaining the given rates) 10 buſhels of Wheat, 10 buſhels of Rye, 14 buſhels of Barley, and 14 buſhels of Oats being all mixed together will conſtitute a *maſs* of Corn, that may be afforded at 28 *d.* or 2 *s.* 4 *d.* the buſhel.

By this Example thus *diverſified* it plainly appears, that the quantities required may be altered as often as the Queſtion given will admit divers *Alligations*, and yet the mixture produced will ſtill hold the *rate* propounded; but when the *Queſtion* propounded will admit but one only way of *Alligation*, the quantities required to make the *mixture*, cannot be varied; ſo the ſecond *Example* of the tenth Rule of this Chapter, being again produced, and ordered according to the direction of the eleventh Rule aforegoing, *I ſay*,

I. As 4 to 4, ſo 10 to 10 buſhels of Rye.
II. As 4 to 4, ſo 10 to 10 buſhels of Barley.
III. As 4 to 60, ſo 10 to 150 buſhels of Oats.



4	4
4	4
4	4
32, 20, 8.	60.

So that for this queſtion *I conclude*, to 10 buſhels of Wheat you ought to add 10 buſhels of Rye, 10 buſhels of Barley, and 150 of Oats, to the end that a *mixture* of Corn might be made, which may be ſold at 16 *d.* the buſhel: And here the quantities found (*viz.* 10, 10, and 150) cannot be *altered*, becauſe the terms of this Queſtion will not admit any other variety of *Alligation.*

The Proof.

XV. In Alternation Partial, the proof is likewiſe by comparing the total value of the ſeveral ſimples, with the value of the whole mixture: So in the ſecond example of the laſt Rule, the total *value* of the 10 buſhels of Wheat, 40 buſhels of Rye, 50 buſhels of Barley, and 20 buſhels of Oats, amounts to 14 *l.* which is alſo the value of the whole mixture at 2 *s.* 4 *d.* the buſhel, as appears by the example of the fifth Rule of this preſent Chapter.

Alternation total.

XVI. Alternation total is, when having the total quantity of all the ſimples, together with their ſeveral rates, we produce their ſeveral quantities, in ſuch ſort, that a mixture of them being made according to the quantities ſo found, that mixture may bear a certain rate propounded: Of this ſort is the laſt example of the tenth Rule aforegoing; as alſo *this:* A Goldſmith having divers ſorts of Gold, *viz.* ſome of 24 Careɛts, other of 22 Careɛts, ſome of 18 Careɛts, and other ſome of 16 Careɛts *fine*, is deſirous to melt of all theſe ſorts ſo much together, as may make a *maſs* containing 60 ounces of 21 Careɛts *fine*: Now this Rule of *Alternation total* ſheweth you how much you are to take of each ſort, to the end the whole maſs *may* contain juſt 60 ounces of 21 Carects, the *fineneſs* propounded.

The Proportions.

XVII. In Queſtions of Alternation total the proportion is, as followeth.

As the ſum of all the differences is to the total quantity of all the ſimples: So is the correſpondent difference of each rate to the reſpective quantity of the ſame rate.

So the laſt example of the laſt Rule being propounded, *I ſay*,

I. As 12 the ſum of the differences is to 60 ounces the *total* quantity of all the ſimples: ſo is 5 the correſpondent *difference* of 24 Carects the firſt rate, to 25 ounces, *viz.* the required quantity of the Gold of the ſame rate, which may be taken to make the mixture propounded.

II. As 12 to 60, ſo is 3 the correſpondent *difference* of 22 Carects the ſecond rate, to 15 ounces, *viz.* the quantity of the Gold of 22 Carects, that ought to be uſed in the mixture.

III. As 12 to 60; ſo is 1 to 5 ounces of the Gold of 18 Carects fine.

IV. As 12 to 60, ſo is 3 to 15 ounces of the Gold of 16 Carects *fine*, which are requiſite to be taken for the mixture propounded.



Whereupon *I conclude*, that 25 ounces of 24 Carects *fine*, 15 ounces of 22 Carects, 5 ounces of 18 Carects, and 15 ounces of 16 Carects *fine*, being all melted together will produce a *maſs* of Gold containing 60 ounces of 21 Carects *fine*, which is the reſolution of the *Queſtion* propounded.

Again, The laſt Example of the tenth Rule being here repeated, and ordered according to the direction of the eleventh Rule, *I ſay*,

I. As 64 to 192, ſo 17 to 51 ounces of 24 Carects *fine*.

II. As 64 to 192, ſo is 17 to 51 ounces of 21 Carects *fine*.

III. As 64 to 192, ſo is 17 to 51 ounces of 19 Carects *fine*.

IV. As 64 to 192, ſo is 13 to 39 ounces of *Alloy*.



And therefore for *concluſion* I ſay, that 51 ounces of Gold, 24 Carects *fine*, 51 ounces of 21 Carects *fine*, 51 ounces of 19 Carects *fine*, and 39 ounces of *Alloy* being all mixed together, will produce a *maſs* containing 192 ounces of Gold, 17 Carects *fine*, which is the *ſatisfaction* of the *queſtion* premiſed.

And here *obſerve* (as before in the Expoſition of the fourteenth Rule of this Chapter) that the operations of the firſt of theſe Examples may be *varied* according to the diverſity of the *Alligations* which it will admit, whereas the laſt Example is not ſubject to any *variety*, the *Alligations* thereof remaining always the ſame.

XVIII. Here the operation is perfect, when the ſum of the quantities found agrees with the total quantity propounded; So in the firſt Example of the laſt Rule, 25, 15, 5, and 15 (the quantities found) being all added together amount to 60, which is the total quantity propounded. *The Proof.*

CHAP. XV.

The Rule of Falſe.

I. THE Rule of Falſe is always performed by falſe and ſuppoſitial Numbers taken at pleaſure after the Propoſition is made, and the queſtion propounded; for things are ſaid to be found out by the *Rule of Falſe*, when by falſe terms *ſuppoſed*, we *diſcover* the true terms *required*.

II. The *Rule of Falſe*, is either of ſingle or double Poſition.

III. The Rule of ſingle Poſition is when at once, *viz.* by one falſe Poſition we have means to diſcover the true reſolution of the Queſtion propounded. *The Rule of ſingle Poſition.*

For Example: *A*, *B*, and *C*, determining to buy together a certain quantity of Timber, that ſhould coſt them 36 *l.* agree among themſelves that *B* ſhall pay of that ſum a *third* part more than *A*, and that *C* ſhall pay a *fourth* more than *B*. Now the Queſtion is, What particular ſum each of theſe parties ought to pay of the 36 *l.* To reſolve this Queſtion; firſt, put the caſe that *A* ought to pay 6 *l.* of the 36 *l.* and then *B* muſt pay 8 *l.* becauſe he pays one third part more than *A.* And laſtly, *C* ought to pay 10 *l.* becauſe he is to lay out one fourth part more than *B.* This done although by addition of theſe three ſums, *viz.* 6, 8, and 10, I find that I have made a wrong *Poſition* (their total amounting only to 24 *l.* which ought to have been 36 *l.*) nevertheleſs by thoſe *ſuppoſitional* Numbers, I have means to diſcover the true ſums which the ſeveral parties ought to pay: For I ſay by the Rule of *Three Direct.*

I. As 24 to 36, ſo is 6 to 9 *l.* the part that *A* muſt pay.

II. As 24 to 36, ſo is 8 to 12 *l.* the part that *B* ought to pay.

III. As 24 to 36, ſo is 10 to 15 *l.* the part of the 36 *l.* that *C* muſt pay.

The Proof.

IV. Here for trial of this Rule the total of the ſums found ought to accord with the ſum given: So in the Example of the laſt Rule, 9, 12, and 15 being all added together amount to 36, the ſum propounded.

The Rule of double Poſition.

V. The Rule of double *Poſition* is, when two falſe *Poſitions* are ſuppoſed for the reſolution of the queſtion propounded. As in *this*, A Workman having threſht out 40 quarters of Grain (part thereof being Wheat, and the reſt Barley) received for his labour 28 *s.* being paid after the rate of 12 *d.* for every quarter of Wheat, and 6 *d.* for each quarter of Barley: Now here the queſtion is how many of thoſe 40 quarters were Wheat, and how many Barley? Here therefore I firſt ſuppoſe at random, that there were 26 quarters of Wheat, and 14 of Barley, and then to diſcover whether I have gueſſed right or wrong, I find how much Money is due to the Workman at the rate of 12 *d.* the Quarter of Wheat, and 6 *d.* the Quarter of Barley, which I find to be 33 *s.* (*viz.* 26*s.* for the 26 Quarters of Wheat, and 7 *s.* for the 14 quarters of Barley) which he ought to have received, if my *ſuppoſition* had been right; but becauſe it differs from 28 *s.* the true ſum that he received, I perceive I have miſt the mark, and therefore diſcovering how much I have err'd by finding the *difference* betwixt 28*s.* and 33*s.* I keep in mind 5 their *difference*, which is called the *firſt errour*, or the *errour of the firſt Poſition*: Again, I propound for the *ſecond Poſition*, that there was 30 quarters of Wheat, and 10 quarters of Barley; and then the *ſecond errour* I find to be 7; for there is then due to the Workman for the 30 quarters of Wheat 30 *s.* and for the 10 quarters of Barley 5 *s.* in all 35 *s.* which differs from 28 *s.* the true ſum that he received, by 7 *s.* and here by theſe two *falſe Poſitions*, together with their *errours*, you may diſcover how many quarters of Wheat, and how many of Barley the Workman threſht, as ſhall be further explained by the *Rules following.*

The Operation.

VI. In the Rule of double *Poſition* having drawn two lines acroſs, and placed the terms of the falſe *Poſition* (*viz.* thoſe that have the ſame Denomination) at the uppermoſt end of that Croſs, as alſo each *errour* under his reſpective *Poſition* at the lower end of the ſame Croſs, multiply each *errour* by the contrary

Poſition; that is, the ſecond *errour* by the firſt *Poſition*, and the firſt *error* by the ſecond *Poſition*; this done, when both the *errours* are of one and the ſame kind (*viz.* both exceſſes or both defects) ſubtract the leſs product out of the greater, and then the remainder is your Dividend; but if the *errours* be of different kinds, (*viz.* one of them an exceſs, and the other a defect) add thoſe Products together, and then the ſum will be your Dividend, which if you divide by the difference of the *errours*, (when they are of one and the ſame kind) or by their ſum (when they are of different kinds) the *Quotient* will give you a number you look for, having the ſame Denomination with the falſe *Poſitions* placed at the upper end of the Croſs.

Example 1. The Queſtion of the laſt Rule being again propounded, I place theſe terms, *viz.* 26 (having the Denomination of the Quarters of Wheat in the *firſt Poſition*) and 30 (having the ſame Denomination in the *ſecond Poſition* at the upper end of the Croſs:) As alſo 5 and 7 the two *errours* reſpectively under them at the lower end of the ſame Croſs, as you may ſee it exemplified by the Pattern *following*.

Note, *That this Character,* —— *ſignifies that the leſſer of the two Numbers, betwixt which it is found, ought to be ſubtracted from the greater.*



This done, having multiplied 26 by 7, the *Product* is 182, and likewiſe 30 by 5, the *Product* is 150, which being deducted out of 182 (becauſe the *errours* here are both of the ſame kind, that is, are each of them an exceſs above 28 *s.* the ſum that the Workman received) the *remainder* is 32, which being divided by 2 (the difference betwixt 5 and 7 the two errours) leaves in the *Quotient* 16, for the quarters of Wheat that the Workman threſht, whoſe complement to 40 *viz.* 24, are the quarters of Barley, that he likewiſe threſht; ſo at laſt I conclude, the Workman received 28 *s.* for his wages in threſhing out 40 quarters of Grain (being part Wheat, part Barley) at 12 *d.* the quarter of Wheat, and 6 *d.* the quarter of Barley, threſhed in all 16 quarters of Wheat, and 24 quarters of Barley.

Example 2. The ſame queſtion being again propounded, I ſuppoſe for my *firſt Poſition* that there are 8 quarters of Wheat, and 32 quarters of Barley, and then the firſt errour will be 4 *s.* for 8 *s.* being accounted for the 8 quarters of Wheat, and 16 *s.* for the 32 quarters of Barley, make in all 24*s.* which wants 4 *s.* of 28 *s.* the ſum received: Again, *ſuppoſing* that there are 12 quarters of Wheat, and 28 quarters of Barley, the ſecond errour will be 2 *s.* for 12 *s.* being allowed for the 12 quarters of Wheat, and 14 *s.* for the 28 quarters of Barley, the ſum is 26 *s.* which comes 2 *s.* ſhort of 28 *s.* the right ſum: now then 8 being multiplied by 2, the Product is 16; likewiſe 12 by 4 produceth 48, out of which if you deduct 16 (becauſe the errours in this caſe happen to be both defects under 28 *s.* the ſum received) the remainder is 32, which being divided by 2 (the difference of the errours) gives you in the quotient 16, *viz.* the quarters of Wheat, *as before.*



Example 3. The ſame demand being the third time produced, I take for my *firſt Poſition* 10 quarters of Wheat, and 30 quarters of Barley, and then proceeding as before, the firſt errour will prove 3 *s.* which upon that *Poſition* I want of 28 *s.* the right ſum: Again, here for the *ſecond Poſition* I take 26 quarters of Wheat, and 14 quarters of Barley, and then the ſecond errour will be 5 *s.* which upon that *Poſition* I have exceeded 28 *s.* the true ſum: Now then multiplying 10 by 5, the Product is 50, and 26 by 3, the Product is 78: And here (becauſe the errours are of different kinds, one of them being a *defect*, and the other an *exceſs* of 28 *s.* the true ſum) you are to add 50 and 78 the two Products together, whoſe ſum is 128, which being divided by 8, the ſum of 3 and 5 the two errours, gives you in the quotient 16 for the quarters of Wheat, *as before* in the former reſolutions, So that what *Poſitions* ſoever you take in this *Queſtion*, you ſhall always find, that the Workman threſhed 16 quarters of Wheat, and 24 quarters of Barley, which is the Reſolution of the Queſtion propounded.



Note, *That this Character* + *intimates that the Numbers, betwixt which it is found, ought to be added together.*

VII. Here the trial is the ſame with that which is uſed in finding out the errours: So in the Example premiſed 16 and 24 being the numbers found, and 16 *s.* being allowed for the 16 quarters of Wheat, likewiſe 12 *s.* for the 24 quarters of Barley, their ſum is 28*s.* which was the ſum received by the *Workman.*

Example 4. A certain Man being demanded what was the Age of each of his 4 Sons? Anſwered, that his eldeſt Son was 4 Years elder than the ſecond; his ſecond was 4 Years elder than the third; his third Son was 4 Years elder than the fourth or youngeſt; and his fourth or youngeſt was half the Age of the eldeſt, the Queſtion is, what was the Age of each Son? Here I gueſs the Age of the eldeſt Son to be 16, then it may be infer'd from the Queſtion, that the Age of the ſecond Son was 12, the Age of the third 8, and the Age of the fourth or youngeſt 4, this 4 ſhould be half 16 (for the Queſtion ſaith, that the Age of the youngeſt was half the Age of the eldeſt) but it wants 4 of what it ought to be; wherefore I make a ſecond Poſition, and take 20 for the Age of the eldeſt, then the Age of the ſecond muſt neceſſarily be 16, the Age of the third 12, and the Age of the fourth 8, which ſhould be half 20, but it wants 2: now (according to the Rule) multiplying 16 (the firſt Poſition) by 2 (the ſecond errour) the Product is 32: alſo multiplying 20 (the ſecond Poſition) by 4 (the firſt errour) the Product is 80, and becauſe the errours are both of one kind, to wit, both defective; I ſubtract the leſſer Product from the greater, ſo the remainder is 48 for a Dividend, alſo ſubtracting the leſſer errour from the greater, the remainder is 2 for a Diviſor: Laſtly, dividing 48 by 2, the quotient is 24, and ſuch was the Age of the eldeſt Son, therefore the Age of the ſecond was 20; the Age of the third 16; and the Age of the fourth 12; which is half the Age of the eldeſt, as was declared by the queſtion.



The Doctrine of Vulgar Fractions.

CHAP. XVI.

Notation of Vulgar Fractions.

I. THus far of *Arithmetick* in *whole numbers*, only the Doctrine of *Fractions* enſueth, which depends upon this ſuppoſition, that Unity, or at leaſt one whole thing, whatſoever it be, may in mind be conceived diviſible into any number of equal parts: ſome will not allow 1 or unity to be a number, when it is conſidered in the abſtract, and ſeparated from matter, but foraſmuch as that Prince of Arithmeticians, *Diophantus* of *Alexandria*, in divers of his ſubtil Problems doth mention unity as a number, and propounds it to be divided into numbers, I ſhall take the like liberty to eſteem 1 or unity as a number, and likewiſe ſuppoſe it diviſible into any number of equal parts.

A Fraction.

II. A broken number, otherwiſe called a *Fraction*, is only part of an integer or whole thing, as if you would expreſs in figures the length of a piece of cloth, that contains *three fourths*, or (which is all one) *three quarters* of a yard, you are to write it thus, $\frac{3}{4}$, that is, an entire yard being ſuppoſed to be divided into four equal parts, the length of the piece propounded is three of thoſe four parts: In like manner (a Foot being divided into 12 Inches) you muſt write ſix Inches thus $\frac{6}{12}$, that is, *ſix twelve parts* of a foot; or if the foot be divided into one hundred equal parts, to expreſs five and twenty of thoſe parts, ſet them down thus $\frac{25}{100}$ that is five and twenty hundredth parts of a foot.

III. A Fraction conſiſts of two parts, the *Numerator* and the *Denominator*, which are placed one above the other, and ſeparated by a little line.

IV. The *Numerator* is the number placed above the line, and the *Denominator* is the number placed underneath: ſo in the aforementioned *Fraction* $\frac{3}{4}$, the number 3 placed above the line is the *Numerator*, and the number 4 placed underneath is the *Denominator.* Alſo in this Fraction $\frac{6}{12}$, the *Numerator* is 6, and the *Denominator* is 12. The *Denominator* is ſo called, becauſe it denominates or declares into how many equal parts the Integer or whole thing is ſuppoſed to be divided, and the *Numerator* is ſo called, becauſe it numbreth or expreſſeth how many of thoſe equal parts of the Integer are ſignified by the Fraction.

3 *Numerator.*
4 *Denominator.*

V. A Fraction is either proper or improper.

A proper Fraction. *VI.* A proper Fraction is that that whoſe Numerator is leſs than the Denominator, ſuch are the *Fractions* before-mentioned $\frac{3}{4}$, $\frac{6}{12}$, $\frac{25}{100}$, and the like.

VII. A proper Fraction is either ſingle or compound.

A ſingle Fraction *VIII.* A ſingle Fraction is that which conſiſts of one Numerator, and one Denominator; ſuch are $\frac{3}{4}$, $\frac{6}{12}$, $\frac{25}{100}$, and the like.

IX. A ſingle Fraction doth often ariſe in Diviſion of whole Numbers, for when Diviſion is finiſht, if any number remain, it is to be eſteemed as the Numerator of a Fraction, which hath the Diviſor for a Denominator, and is to be annexed to the Integer or Integers in the quotient as part of the quotient; which Fraction doth always expreſs certain parts (or at leaſt a part) of an integer or entire unit, which hath the ſame Denomination with one of the Integers in the quotient; ſo if 17 pounds be given to be divided equally amongſt 5 perſons, there will ariſe 3 entire pounds in the quotient, and there will be a remainder or ſurpluſage of 2 pounds, which 2 is to be placed, as the Numerator of a *Fraction*, over the Diviſor 5 as a *Denominator*; ſo will the *Fraction* be $\frac{2}{5}$, and the compleat quotient will be $3\frac{2}{5}$, that is, 3 pounds, and 2 fifth parts of a pound for each perſons ſhare.

5) 17 ($3\frac{2}{5}$

A ſingle Fraction doth likewiſe ariſe, when a leſſer whole number is given to be divided by a greater, for in ſuch caſe the *Dividend* is to be made the *Numerator* of a Fraction, and the *Diviſor* the *Denominator*; which Fraction is the true quotient, and doth always expreſs a certain part (or at leaſt parts) of an Integer, which hath the ſame name with the *Dividend:* ſo if 3 pounds ſterling be given to be divided equally among 4 Perſons, the ſhare of each, that is, the quotient will be $\frac{3}{4}$, to wit, 3 fourth parts of a pound.

In like manner, if 5 be given to be divided by 8, the quotient is $\frac{5}{8}$, ſo that the *Numerator* of a Fraction is always a *Dividend*, the *Denominator* is a *Diviſor*, and the *Fraction* it ſelf is the *quotient.*

A Compound Fraction.

X. A Compound Fraction (otherwiſe called a Fraction of a Fraction) is that which hath more *Numerators* and *Denominators* than one, and may be diſcovered by the word (*of*) which is interpos'd between the parts of ſuch compound Fraction: ſo $\frac{2}{3}$ of $\frac{3}{4}$ is a Fraction of a Fraction, or *compound Fraction*, and expreſſeth two thirds of three fourths of an *Integer*, viz. a pound ſterling being ſuppoſed the *Integer*, and firſt divided into four parts, three of thoſe four parts are equal to 15 *s.* Again, if the ſaid 15 *s.* be divided into three parts, two of thoſe three parts are equal to 10 *s.* therefore the *compound Fraction* $\frac{2}{3}$ of $\frac{3}{4}$ of a pound ſterling doth expreſs 10 *s.* In like manner the compound Fraction $\frac{1}{4}$ of $\frac{3}{4}$ of $\frac{4}{5}$ of a pound ſterling, that is, one fourth of three fourths of four fifths of a pound ſterling doth expreſs 3 *s.* as will be farther manifeſted by the ſixteenth and ninth Rules of the ſeventeenth Chapter.

An improper Fraction.

XI. An improper Fraction is that, whoſe Numerator is either greater, or at leaſt equal unto the Denominator: ſo this Fraction $\frac{16}{4}$, that is 16 fourths, is called an *Improper Fraction*, and ſo is this $\frac{4}{4}$; for indeed a Fraction of this kind may well be ſurnamed *Improper*, becauſe it will not admit the definition of a true Fraction, ſince it is always greater than an entire unity, or at leaſt eqaul unto it; ſo ſixteen Farthings, or $\frac{16}{4}$ of a penny are equal to 4 entire pence; and 4 Farthings, or $\frac{4}{4}$ of a penny are equal to 1 penny; therefore when the Numerator is greater than the Denominator, ſuch *improper* Fraction ſignifieth more than 1 or an Integer, but when the Numerator is equal to the Denominator (be it what number ſoever) ſuch *improper* Fraction is always equal to unity, or 1 Integer.

A mixt number.

XII. A mixt number conſiſts of entire unities (or Integers) or at leaſt of unity (or 1 Integer) and a Fraction annexed: So 5$\frac{11}{12}$, 1$\frac{3}{4}$, and ſuch like; are called mixt numbers; So that if a piece of Timber be five feet and eleven inches in length, you are to write that length thus, 5$\frac{11}{12}$; In like manner, one Mile and three quarters or fourths of a mile are to be written thus, 1$\frac{3}{4}$.

CHAP. XVII.

Reduction of Vulgar Fractions.

I. THe ſame parts of *Numeration*, as have been wrought in *whole Numbers* in the preceeding Chapter, are likewiſe to be performed in *fractions*, but firſt of all *Reduction* of *Fractions* in divers kinds muſt be known, which being the principal skill in the doctrine of Fractions, muſt be diligently obſerved by the Learner.

II. A number is ſaid to be a common Meaſure or Diviſor unto two or more numbers given, when it will meaſure or divide every one of the numbers given, and leave no remainder; ſo 4 is a common meaſure unto the numbers 12 and 20; for if 12 be divided by 4, the *Quotient* will be exactly 3, without any remainder or ſurpluſage; alſo if 20 be divided by the ſame Diviſor 4, the quotient will be

Preciſely 5 without any remainder; in like manner 5 is a *common Diviſor* unto theſe three numbers 10, 25 and 40.

To find the greateſt common meaſure unto any two Numbers.

III. Two numbers being given their greateſt common Diviſor; that is, the greateſt number which will meaſure or divide each of the numbers given without leaving any remainder, may be found out in this manner; *viz.* Divide the greater number by the leſs, then divide the Diviſor by the remainder (if there be any) and ſo continue dividing the laſt Diviſors by the remainders, until there be no remainder (neglecting the quotients;) ſo is the laſt Diviſor the greateſt common Diviſor unto the numbers given.

Thus, If the greateſt *common Diviſor* unto the numbers 91 and 117 be ſought, divide the greater number 117 by 91, the remainder is 26, by which dividing 91, the remainder is 13; by which dividing 26 the remainder is 0; ſo is 13 the greateſt *common Diviſor* unto the numbers 117 and 91, as is manifeſt in dividing each of them by 13; for 13 is found in 91 preciſely 7 times, and in 117 preciſely 9 times. In like manner, 29 will be found a common Diviſor unto 116 and 145; and 51 a common Diviſor unto 561 and 612.

```
91)117(1
   91
  ----
   26)91(3
      78
     ----
      13)26(2
         26
         --
          0
```

*To reduce a Fraction into the leaſt terms,*viz. *1. By a general Rule.*

IV. A ſingle Fraction may be reduced into the leaſt terms by dividing the Numerator & Deno-

minator by their greateſt common meaſure(or divi-
or;) for the quotients will be the Numerator and
Denominator of a fraction equal to the former, and
n the leaſt terms.

So if the fraction $\frac{91}{117}$ be given to be reduced in-
o the *leaſt terms*, ſearch out the greateſt common
Diviſor unto 91 and 117 by the laſt *Rule*, which
vill be found 13, and then dividing 91 by 13, the
quotient will be 7 for a new Numerator; alſo divi-
ing 117 by 13, the quotient will be 9 for a new
Denominator: ſo the fraction $\frac{91}{117}$ is reduced into the
eaſt terms, *viz.* into the fraction $\frac{7}{9}$. In like manner $\frac{216}{145}$
vill be reduced into $\frac{4}{5}$; And $\frac{561}{612}$ unto $\frac{11}{12}$: But here
ou are to obſerve, that if the greateſt *common Divi-
or* unto the Numerator and Denominator be 1,
uch Fraction is in its *leaſt terms* already: ſo the fra-
tion $\frac{189}{151}$ cannot be reduced into lower terms, be-
auſe the greateſt *common Diviſor* will be found 1,
(by the third *Rule* of this Chapter;) the like may
appen of infinite others: And although the laſt be
general *Rule* for the Reduction of Fractions into
heir *leaſt terms*, yet there are other practical *Rules*;
which in ſome caſes will be more ready (eſpecially
nto beginners) *viz.*

2. *By particular Rules.*

V. When the Numerator and De-
ominator are even numbers, they
nay be meaſured or divided by 2.
Therefore in ſuch caſe you may (as is taught in
he Rules of the 6th Chapter) take the half of the
Numerator for a new Numerator, alſo the half of
he Denominator for a new Denominator. So
f $\frac{16}{64}$ be given, draw at length the
ine which ſeparates the Numera-
or from the Denominator, and

16	8	4	2	1
64	32	16	8	4

croſs'd the ſame with a downright ſtroke near the Fraction, as you may ſee in the *Margin*, then take the half of 16, which is 8, for a new Numerator, and the half of 64, which is 32, for a new Denominator; Again, the half of 8 is 4, for a new Numerator, alſo the half of 32 is 16, for a new Denominator, and proceeding in the like manner, there will be found $\frac{1}{4}$ equivalent unto $\frac{16}{64}$.

VI. When the Numerator and Denominator do each of them end with 5, or one of them ending with 5, and the other with a Cypher, they may be both meaſured or divided by 5. So $\frac{225}{475}$ will be reduced into $\frac{9}{19}$ and $\frac{50}{425}$ into $\frac{2}{17}$, as by the operation in the *Margin* is manifeſt.

$$\frac{225}{475}\Big|\frac{45}{95}\Big|\frac{9}{19}$$

$$\frac{50}{425}\Big|\frac{10}{85}\Big|\frac{2}{17}$$

VII. Whenſoever you can eſpy any other number, which will exactly divide the Numerator and Denominator (although it be not the greateſt common Diviſor) you may divide the Numerator and Denominator by ſuch number as before: So $\frac{28}{84}$ may be firſt reduced into $\frac{7}{21}$ by 4, and $\frac{7}{21}$ may be reduced into $\frac{1}{3}$ by 7, as by the operation is manifeſt.

$$\frac{28}{84}\Big|\frac{7}{21}\Big|\frac{1}{3}$$

VIII. When the Numerator and Denominator do each of them end with a Cypher or Cyphers, cut off equal Cyphers in both, and the fraction will be reduced into leſſer terms: So $\frac{400}{500}$ is reduced into $\frac{4}{5}$, and $\frac{700}{9000}$ into $\frac{7}{90}$.

$$\frac{4}{5}\Big|\frac{00}{00}$$

$$\frac{7}{90}\Big|\frac{00}{00}$$

To find the value of a ſingle Fraction in the known parts of the Integer.

IX. The value of a ſingle fraction in the known parts of the Integer, may be found out in this manner, *viz.* multiply the Numerator of the fraction propounded by the number of known parts of the next inferiour denomination which are equal to the Integer, and divide that product by the Denominator, ſo is the quotient the value of the fraction in that inferiour denomination, and if there happen to be any fraction in the quotient, you may find the value thereof in the next inferiour denomination, by the ſame Rule, and ſo proceed till you come to the leaſt known parts.

So the value of $\frac{9}{16}$ of a pound *ſterling* will be found 11 *s.* 3*d.* *viz.* multiply the Numerator 9 by 20 (the number of ſhillings which are equal to 1 pound *ſterling*) the Product is 180, which being divided by the Denominator 16, the quotient is $11\frac{4}{16}$ ſhillings. In like manner, the value of $\frac{4}{16}$ of a ſhilling will be found 3 pence, for multiplying the Numerator 4 by 12 (the number of pence in a ſhilling) the product is 48, which being divided by the Denominator 16, the quotient is 3 pence. Alſo the value of $\frac{7}{13}$ of a pound *ſterling*, will be found 10 *s.* $9\frac{3}{13}$*d.* And $\frac{31}{96}$ of a pound *Troy* will be found equivalent unto 3 ounces 17 penny weight and 12 grains.

$$\begin{array}{r}
9 \\
20 \\
\hline
16)\ 180\ (11\frac{4}{16} \\
16 \\
\hline
20 \\
16 \\
\hline
4 \\
12 \\
\hline
16)\ 48\ (3 \\
48 \\
\hline
0
\end{array}$$

To reduce a mixt number into an improper fraction.

X. A mixt number may be reduced into an improper fraction equivalent unto the mixt number, In this manner, *viz.* Multiply the Integer or Integers in the mixt number by the Denominator of the fraction annexed to the Integer or Integers, and unto the Product add the Numerator of the ſaid fraction; ſo is the ſum the Numerator of an improper fraction, whoſe Denominator is the ſame with that of the ſaid fraction annexed.

So $4\frac{11}{12}$ will be reduced into the improper fraction $\frac{59}{12}$, for 4 being multiplied by 12, the Product is 48, unto which adding the Numerator 11, the ſum is 59 for a new Numerator, which being placed over the Denominator 12, gives the improper fraction $\frac{59}{12}$; which is equivalent unto $4\frac{11}{12}$ (as will appear by the 13*th Rule* of this Chapter.) In like manner $7\frac{1}{2}$ will be reduced into $\frac{15}{2}$.

To reduce a whole number into an improper fraction.

XI. A whole number is reduced into an improper fraction, by placing the whole number given as a Numerator, and 1 as a Denominator.

So 14 Integers will be reduced into the *improper fraction* $\frac{14}{1}$, and one Integer into the *improper fraction* $\frac{1}{1}$.

XII. A whole number is reduced into an improper fraction which ſhall have any Denominator aſſigned, in multiplying the whole number given by the Denominator aſſigned, and placing the Product as a Numerator over the ſaid Denominator.

As if 13 be given to be reduced into an *improper fraction*, whoſe Denominator ſhall be 4 multiply 13 by 4, the Product is 52, which being placed over 4, gives the *improper fraction* $\frac{52}{4}$ equivalent unto 13 (as will appear by the next *Rule.*) In like manner 13 may be reduced into $\frac{91}{7}$.

XIII. An improper fraction may be reduced into its equivalent whole number or mixt number in this manner, *viz.* divide the Numerator by the Denominator, and the quotient will give the whole number or mixt number ſought; So the improper fraction $\frac{59}{12}$ will be reduced into this *mixt number* $4\frac{11}{12}$, for if 59 be divided by 12, the quotient is $4\frac{11}{12}$. Alſo this improper fraction $\frac{52}{4}$ will be reduced into the whole number 13.

To reduce an improper Fraction into its equivalent whole or mixt number.

XIV. Fractions having unequal Denominators may be reduced into fractions of the ſame value, which ſhall have equal Denominators, by this Rule and the next following, *viz.* when two fractions having unequal Denominators are propounded to be reduced into two other fractions of the ſame value, which ſhall have a common Denominator, multiply the Numerator of the firſt fraction (that is either of them) by the Denominator of the ſecond, and the Product ſhall be a new Numerator (correſpondent unto the Numerator of that firſt fraction;) alſo multiplying the Numerator of the ſecond fraction by the Denominator of the firſt, the Product is a new Numerator (correſpondent unto the Numerator of the ſecond fraction;) laſtly, multiply the Denominators one by the other, and the Product

To reduce fractions to a common denominator, viz. 1. *When two fractions are propounded.*

Product is a common Denominator to both the new Numerators.

Thus, If the Fractions $\frac{2}{3}$ and $\frac{4}{5}$ be propounded, multitiply 2 by 5, the product 10 is a new Numerator correſpondent unto 2: alſo multiply 4 by 3, the product 12 is a new Numerator correſpondent unto 4: laſtly, multiply 3 by 5, and the product 15 ſhall be a common *Denominator* unto the new Numerators, ſo the fractions $\frac{10}{15}$ and $\frac{12}{15}$ are found out, which have *equal Denominators*, and each of theſe new fractions is equal unto its correſpondent fraction firſt given, *viz.* $\frac{10}{15}$ is equal unto $\frac{2}{3}$ and $\frac{12}{15}$ is equal unto $\frac{4}{5}$, as will be manifeſt by the 4th *Rule* of this Chapter.

$$\begin{array}{ccc} 2 & \times & 4 \\ \text{--} & & \text{--} \\ 3 & & 5 \\ 10 & & 12 \\ \hline 15 & & 15 \end{array}$$

2 *When three or more Fractions are to be reduced into others that ſhall have a Common Denominator.*

XV. When three or more Fractions having unequal Denominators, are given to be reduced into other Fractions of the ſame value with thoſe given, but ſuch as ſhall have one common Denominator; multiply *continually* (according to the thirteenth Rule of the fifth Chapter) the Numerator of the firſt Fraction into all the Denominators, except its own, *i. e.* the Denominator of that firſt Fraction; and reſerve the laſt Product for a new Numerator inſtead of that firſt Numerator: In like manner, multiply *continually* the Numerator of the ſecond Fraction into all the Denominators, except the Denominator of the ſecond Fraction, and reſerve the laſt Product for a new Numerator, inſtead of the ſecond Numerator; Proceed in like manner to find out new Numerators for the reſt of the given Fractions: Laſtly, multiply *continually* all the Denominators one into another, and the laſt Product ſhall be a common Denominator to all the new Numerators.

As for Example, if theſe three Fractions, $\frac{3}{8}$, $\frac{2}{5}$, $\frac{5}{7}$, having unequal (or different) Denominators, be given to be reduced into three other Fractions of the ſame value, which ſhall have equal Denominators (or one common Denominator.) Firſt, I multiply *continually* the firſt Numerator 3 into the ſecond and third Denominators 5 and 7, ſaying 3 times 5 makes 15, which multiplied by 7 produceth 105, for a new Numerator inſtead of the firſt Numerator 3; Secondly, I multiply *continually* the ſecond Numerator 2 into the firſt and third Denominators 8 and 7, ſaying, twice 8 is 16, which multiplied by 7 produceth 112, for a new Numerator inſtead of the ſecond Numerator 2; Thirdly, I multiply *continually* the third Numerator 5 into the firſt and ſecond Denominators 8 and 5, ſaying 8 times 5 makes 40, which multiplied by 5 produceth 200, for a new Numerator inſtead of the third Numerator 5; Fourthly and laſtly, I multiply *continually* all the Denominators 8, 5 and 7 one into another, ſaying, 8 times 5 makes 40, which multiplied by 7 produced 280 for a Denominator to each of the three new Numerators 105, 112 and 200 before found out; And ſo theſe three Fractions $\frac{105}{280}$, $\frac{112}{280}$, and $\frac{200}{280}$, are diſcovered, which have one common Denominator 280, and each of them is equal in value unto its correſpondent Fraction firſt given, *viz.* $\frac{105}{280}$, is equal unto $\frac{3}{8}$; Alſo $\frac{112}{280}$ is equal unto $\frac{2}{5}$; and $\frac{200}{280}$ is equal unto $\frac{5}{7}$; as may eaſily be prov-

$$\begin{array}{ccc} \frac{3}{8} & \frac{2}{5} & \frac{5}{7} \\ \frac{105}{280} & \frac{112}{280} & \frac{200}{280} \end{array}$$

ed

ved by the fourth Rule of this Chapter.

After the ſame manner theſe four Fractions $\frac{2}{3}$, $\frac{3}{4}$, $\frac{4}{5}$ and $\frac{5}{6}$ are reducible into theſe, $\frac{240}{360}$, $\frac{270}{360}$, $\frac{288}{360}$ and $\frac{300}{360}$, which have 360 for a common Denominator, and are equal in value reſpectively to the four Fractions given to be reduced.

Note, Although by the foregoing fourteenth and fifteenth Rules, any multitude of Fractions may be reduced to a common Denominator; yet becauſe Fractions in their leaſt Terms are fitteſt for uſe, I ſhall ſhew how leſſer Denominators than thoſe that will be diſcovered by the ſaid Rules, may often times be found out, *viz.*

I. When the unequal Denominators of two Fractions have a common Diviſor greater than 1, divide the Denominators ſeverally by their greateſt common Diviſor (found out by the fore-going third Rule of this Chapter;) and then multiply croſs-wiſe in this manner, *viz.* the Numerator of the firſt Fraction by the latter Quotient, and the Numerator of the latter Fraction by the firſt Quotient, and reſerve the Products for new Numerators; laſtly, multiply the Denominator of the firſt Fraction by the latter Quotient (or the Denominator of the latter Fraction by the firſt Quotient,) ſo ſhall the Product be a common denominator to the ſaid new Numerators: As for Example, if $\frac{5}{12}$ and $\frac{7}{18}$ be propoſed to be reduced to a common Denominator, I divide each of the Denominators, 12 and 18, by their greateſt common Diviſor 6, and the



the Quotients are 2 and 3; then I multiply 5 the Numerator of the firſt Fraction by 3 the latter Quotient, alſo 7 the Numerator of the latter Fraction by 2 the firſt Quotient, and the Products 15 and 14 I reſerve for new Numerators inſtead of 5 and 7; Laſtly, I multiply 12 the Denominator of the firſt Fraction by 3 the latter Quotient (or 18 the Denominator of the latter Fraction by 2 the firſt Quotient,) and the Product 36 is a Denomiuator to each of the new Numerators 15 and 14: So $\frac{15}{36}$ and $\frac{14}{36}$ are found out, which have the leaſt common Denominator unto which the given Fractions $\frac{5}{12}$ and $\frac{7}{18}$ can be reduced; Alſo $\frac{15}{36}$ is equal to $\frac{5}{12}$, and $\frac{14}{36}$ to $\frac{7}{18}$.

II. Whenſoever the Denominator of a Fraction can be divided by the Denominator of a ſecond Fraction, without any Remainder; then if by the Quotient you multiply ſeverally the Numerator and Denominator of ſuch ſecond Fraction, a third will ariſe, having the ſame value with the ſecond, and the ſame Denominator with the firſt Fraction: By this Rule Three or more Fractions may often times be reduced to a leſſer common Denominator, than that which will be diſcovered by the foregoing Rule XV. As for Example, let theſe ſix following Fractions be given to be reduced to a common Denominator, *viz.*

$$\frac{13}{36},\ \frac{11}{18},\ \frac{7}{12},\ \frac{4}{9},\ \frac{5}{6},\ \frac{2}{3}.$$

Becauſe 36 the Denominator of the firſt Fraction, being divided by the five other Denominators ſeverally,

rally will give theſe Quotients 2, 3, 4, 6, and 12 without any Remainder, I multiply the Numerator and Denominator of each of the five latter Fractions, by its correſpondent Quotient; *viz.* 11 and 18 by 2 the firſt Quotient; Alſo 7 and 12 by 3 the ſecond Quotient, and in like manner the reſt; ſo inſtead of thoſe five latter Fractions, five others (hereunder placed after the firſt of thoſe ſix) are produced, *viz.*

$$\frac{13}{36},\ \frac{22}{36},\ \frac{21}{36},\ \frac{16}{36},\ \frac{30}{36},\ \frac{24}{36}.$$

All which Fractions laſt expreſs'd have a common Denominator 36, and are equal in value reſpectively to thoſe given to be reduced.

To reduce a compound fraction to a ſingle fraction. See continual multiplication in the laſt Rule of the 5th Chapter.

XVI. A compound Fraction (otherwiſe called a *Fraction of a Fraction*) may be reduced into a ſingle Fraction in this manner, *viz.* Multiply all the Numerators *continually*, and take the Product for a new Numerator, alſo multiply all the Denominators *continually*, and the Product ſhall be a new Denominator.

Thus, if the *compound Fraction* $\frac{2}{3}$ of $\frac{3}{4}$ be given to be reduced into a ſingle Fraction, multiply the Numerators 2 and 3, one by the other, ſo is the Product 6 a new Numerator. Alſo multiplying the Denominators 3 and 4 one by the other, the product 12 is a new Denominator, ſo $\frac{6}{12}$ (or $\frac{1}{2}$ is the ſingle Fraction ſought, being equivalent unto $\frac{2}{3}$ of $\frac{3}{4}$, the compound Fraction given to be reduced.

$\frac{2}{3}$ of $\frac{3}{4}$

$\frac{6}{12}$ or $\frac{1}{2}$

In like manner, this compound Fraction $\frac{2}{3}$ of $\frac{3}{4}$ of $\frac{4}{5}$ will be reduced unto $\frac{24}{60}$, or $\frac{2}{5}$; for the Numerator 2, 3, 4, being multiplied *continually* produce the new Numerator 24, and the Denominators 3, 4, 5 multiplied *continually* produce the new Denominator 60: Laſtly, the new Fraction $\frac{24}{60}$ (by the fourth Rule of this Chapter) will be reduced unto $\frac{2}{5}$, which is equal to $\frac{2}{3}$ of $\frac{3}{4}$ of $\frac{4}{5}$: But to make the meaning hereof more evident, ſuppoſe the Integer to be one Pound of Engliſh Money; then

$\frac{4}{5}$ of 1 *l.* (*viz.* of 20 *s.*) —— 16 *s.*
$\frac{3}{4}$ of thoſe $\frac{4}{5}$ (*viz.* of 16 *s.*) is —— 12 *s.*
$\frac{2}{3}$ of thoſe $\frac{3}{4}$ (*viz.* of 12 *s.*) is —— 8 *s.* or $\frac{2}{5}$ *l.*

whereby 'tis manifeſt that $\frac{2}{3}$ of $\frac{3}{4}$ of $\frac{4}{5}$ *l.* is equal to $\frac{2}{5}$ *l.*

By this Rule a fraction or mixt number of a leſſer name may be reduced to a Fraction of a greater name. As if 3 $\frac{1}{2}$ pence be propounded to be reduced into an improper Fraction of a Pound ſterling, the operation will be in this manner, *viz.* 3 $\frac{1}{2}$ or $\frac{7}{2}$ of a penny is $\frac{7}{2}$ of $\frac{1}{12}$ of $\frac{1}{20}$ of a Pound ſterling, which compound Fraction will (by the aforeſaid Rule) be reduced, to $\frac{7}{480}$ *l.* In like manner 42 $\frac{3}{16}$ minutes of an hour are equal to $\frac{45}{64}$ of an hour, for $\frac{675}{16}$ (that is 42 $\frac{3}{16}$) of $\frac{1}{60}$ are equal to $\frac{675}{960}$ (or in its leaſt terms) $\frac{45}{64}$.

Here you may alſo obſerve, that when a compound fraction is one of the given terms in any queſtion, it is firſt of all to be reduced to a ſingle fraction by the aforeſaid ſixteenth Rule.

To find whole Numbers, which ſhall have the ſame reaſon as any Fractions or mixt numbers given.

XVII. Two or more fractions being given, there may be whole numbers found, which ſhall have the ſame reaſon or proportion as the fractions given, *viz.* When the fractions given have unequal denominators, reduce them into equivalent fractions which ſhall have a common denominator (by the 14th or 15th Rule of this Chapter;) then rejecting the common Denominator, the Numerators ſhall have the ſame reaſon or proportion as the fractions firſt given given.

So $\frac{3}{5}$ and $\frac{5}{8}$ being given, will firſt of all be reduced into their equivalent fractions $\frac{24}{40}$ and $\frac{25}{40}$; then rejecting the common Denominator 40, the Numerators 24 and 25 have the ſame reaſon with $\frac{3}{5}$ and $\frac{5}{8}$ *viz.* As $\frac{3}{5}$ is to $\frac{5}{8}$, ſo is 24 to 25: Alſo it the fractions $\frac{1}{8}$, $\frac{1}{4}$ and $\frac{1}{2}$ were given, there will be found 8, 16 and 32, which are in the ſame proportion one to the other as the fractions given: In like manner if mixt Numbers be given, there may be whole numbers found which ſhall have the ſame *reaſon* or *proportion*, as the mixt numbers; ſo $5\frac{2}{3}$ and $3\frac{5}{8}$ being given, will be firſt reduced into the improper fractions $\frac{17}{3}$ and $\frac{29}{8}$ (by the tenth Rule of this Chapter:) alſo the ſaid $\frac{17}{3}$ and $\frac{29}{8}$ will be reduced into $\frac{136}{24}$ and $\frac{87}{24}$; then rejecting the common Denominator 24, the Numerators 136 and 87 will have the ſame *reaſon* as $5\frac{2}{3}$ and $3\frac{5}{8}$, *viz.* As 136 is to 87, ſo is $5\frac{2}{3}$ to $3\frac{5}{8}$: Alſo $16\frac{1}{2}$ and 18 being given, there will be found 33 and 36, which being divided by their common Diviſor 3 (found out by the third Rule of this Chapter) will give 11 and 12 which have the ſame *reaſon* as $16\frac{1}{2}$ and 18.

CHAP.

CHAP. XVIII.

Addition of Vulgar Fractions and mixt Numbers.

I. WHen the numbers given to be added are ſingle fractions, and have equal denominators, add all the Numerators together, ſo is the ſum the Nnmerator of a fraction, whoſe Denominator is the ſame with the common Denominator, which new fraction is the ſum of the fractions given to be added.

To add ſingle fractions, viz. *1. When they have equal denominators.*

So $\frac{3}{9}$ and $\frac{2}{9}$ being given to be added, their ſum will be found $\frac{5}{9}$, *viz.* the ſnm of the Numerators, 3 and 2, is 5, which being placed over the common Denominator 9, gives $\frac{5}{9}$: In like manner the ſum of theſe fractions $\frac{7}{8}$, $\frac{5}{8}$, $\frac{1}{8}$, and $\frac{2}{8}$, will be found $\frac{17}{8}$, which (by the 13 *Rule* of the ſeventeenth Chapter) will be found equivalent unto 2 $\frac{1}{8}$; ſo that 2 $\frac{1}{8}$ is the ſum of the fractions given to be added.

II. When the Fractions given to be added have unequal Denominators, they are firſt to be reduced into fractions of the ſame value, which ſhall have a common Denominator (by the fourteenth or fifteenth Rule of the ſeventeenth Chapter;) and then they may be added by the firſt Rule of this Chapter.

2. When they have unequal denominators.

So if $\frac{2}{3}$ and $\frac{4}{5}$ were given to be added, their ſum will be found 1 $\frac{7}{15}$ for (by the fourteenth Rule of the ſeventeenth Chapter) $\frac{2}{3}$ and $\frac{3}{5}$ will be reduced into their equivalent fractions $\frac{10}{15}$ and $\frac{9}{15}$ which having equal Denominators may be added according to the firſt rule of this Chapter, and ſo the ſum will be found $1\frac{4}{15}$: In like manner the ſum of theſe fractions $\frac{21}{24}$ and $\frac{3}{4}$ will be found $1\frac{5}{8}$. Alſo the ſum of theſe ſix Fractions $\frac{13}{36}$, $\frac{11}{18}$, $\frac{7}{12}$, $\frac{4}{9}$, $\frac{5}{6}$, $\frac{2}{3}$, after they are reduced to a common Denominator (according to the latter Example in the Note at the end of the fifteenth Rule of the ſeventeenth Chapter) will be found $\frac{126}{36}$, that is, $3\frac{1}{2}$.

2 3
X
3 5
10
9
$\frac{19}{15}$ *that is* $1\frac{4}{15}$

The Addition of compound Fractions.

III. When any of the fractions given to be added is a compound Fraction, ſuch compound fraction is firſt of all to be reduced into a ſingle fraction (by the ſixteenth Rule of the ſeventeenth Chapter) and then you may proceed as before.

So $\frac{3}{5}$ and $\frac{2}{3}$ of $\frac{1}{4}$ being given to be added, their ſum will be found $\frac{23}{30}$ for the compound fraction $\frac{2}{3}$ of $\frac{1}{4}$ will (by the 16th Rule of the 17th Chapter) be reduced to $\frac{2}{12}$ (or in its leaſt terms) $\frac{1}{6}$, which added to the ſingle fraction $\frac{3}{5}$ (according to the ſecond Rule of this Chapter) gives $\frac{23}{30}$. Here you may obſerve, that the fractions given to be added in all the former caſes, are ſuppoſed to be fractions of Integers, which have one and the ſame *particular Denomination*, *viz.* If one of the fractions, given to be added, be a fraction of a pound *ſterling*, all the reſt ought to be fractions of a pound

By Denomination is meant the name of any Integer or thing.

pound *ſterling*, and the like is to be underſtood of other denominations.

IV. When fractions of Integers, of different denominations are given to be added, they are firſt of all to be reduced into fractions of Integers which ſhall have one and the ſame particular denomination (by the ſixteenth Rule of the ſeventeenth Chapter;) and then they may be added by the firſt or ſecond Rule of this Chapter.

To add fractions of Integers which have different denominations.

So if $\frac{7}{9}$ of a pound *ſterling*, $\frac{3}{5}$ of a ſhilling, and $\frac{5}{8}$ of a penny were given to be added, reduce the two latter into fractions of a pound *ſterling* (by the ſixteenth Rule of the ſeventeenth Chapter) *viz.* $\frac{3}{5}$ of a ſhilling is $\frac{3}{5}$ of $\frac{1}{20}$ of a pound *ſterling*, which compound fraction being reduced into a ſingle fraction gives $\frac{3}{100}$ *li.* Likewiſe $\frac{5}{8}$ of a penny, is $\frac{5}{8}$ of $\frac{1}{12}$ of $\frac{1}{20}$ of a pound *ſterling*, which compound fraction being reduced, gives $\frac{1}{384}$ *li.* Laſtly, $\frac{7}{9}$ *li.* $\frac{3}{100}$ *li.* and $\frac{1}{384}$ *li.* being added according to the ſecond Rule of this Chapter, their ſum will be found $\frac{280068}{345600}$, or in its leaſt terms $\frac{23339}{28800}$ *li.*

V. When mixt numbers are given to be added, find firſt of all the ſum of the fractions (by the firſt and the ſecond Rule of this Chapter;) then add the Integer or Integers (if there be any found) in the ſum of the fractions, unto the whole numbers, and collect the ſum of them as you were taught by the Rules of the third Chapter.

To add mixt Numbers.

So if $3\frac{1}{2}$, $4\frac{1}{3}$ and $16\frac{5}{8}$ were given to be added, their ſum will be found $24\frac{11}{24}$ *viz.* the ſum of the fractions $\frac{1}{2}$, $\frac{1}{3}$ and $\frac{5}{8}$, will be found (by the ſecond Rule of this Chapter) to be $1\frac{11}{24}$, and the ſum of the whole numbers, 3, 4, and 16, is 23, unto which adding 1 (the Integer found in the ſum of the fractions) the ſum is 24; ſo that 24 $\frac{11}{24}$ is the ſum of the mixt numbers given to be added.

CHAP. XIX.

Subtraction of Vulgar Fractions and mixt Numbers.

The ſubtraction of ſingle fractions, viz.
1. When they have a common Denominator.

I. WHen the Numbers given are both ſingle Fractions and have equal Denominators, ſubtract the leſſer Numerator from the greater, and place the remainder over the common Denominator, ſo is ſuch new fraction the difference between the fractions given.

Thus the difference between the fractions $\frac{9}{11}$ and $\frac{7}{11}$ is $\frac{2}{11}$, which is found by ſubtracting the leſſer Numerator 7 from the greater Numerator 9, and placing the remainder 2 over the common Denominator 11: alſo the difference between the fractions $\frac{11}{21}$ and $\frac{17}{21}$ is $\frac{6}{21}$, that is, the fraction $\frac{17}{21}$ ex- $\frac{11}{21}$ by $\frac{6}{21}$.

2. When they have unequal Denominators.

II. When the numbers given are both ſingle Fractions and have not a common Denominator, reduce them into fractions of the ſame value which ſhall have a common Denominator (by the fourteenth or fifteenth Rule of the ſeventeenth Chapter) and then find their differences by the laſt Rule.

So the difference between the fractions $\frac{6}{7}$ and $\frac{7}{8}$ will be found $\frac{1}{56}$, *viz.* reducing the fractions given into their equivalent fractions $\frac{48}{56}$ and $\frac{49}{56}$, which have a common Denominator, the difference ſought will be found $\frac{1}{56}$, by the firſt Rule of this Chapter. Likewiſe $\frac{7}{12}$ being ſubtracted from $\frac{11}{13}$, there will remain $\frac{41}{156}$.

The ſubtraction of mixt numbers, viz. 1. *By a general Rule.*

III. When one of the numbers given is a whole number or a mixt number, alſo when both of them are mixt numbers, reduce ſuch whole, or mixt numbers into an improper Fraction or Fractions, by the 10th or 11th Rule of the ſeventeenth Chapter, and then the operation will be according to the firſt or ſecond Rule of this Chapter.

So 7 $\frac{3}{5}$ being given to be ſubtracted from 12, the remainder will be found 4 $\frac{2}{5}$; *viz.* Firſt 7 $\frac{3}{5}$ will be reduced into the improper Fraction $\frac{38}{5}$, alſo 12 will be reduced to $\frac{12}{1}$, then theſe two improper fractions $\frac{38}{5}$ and $\frac{12}{1}$ will be reduced into their equivalent fractions $\frac{38}{5}$ and $\frac{60}{5}$ (which have a common Denominator.) Laſtly, the difference between $\frac{38}{5}$ and $\frac{60}{5}$ is $\frac{22}{5}$, or 4 $\frac{2}{5}$. In like manner 9 $\frac{1}{2}$ being given to be ſubtracted from 12 $\frac{1}{5}$, the remainder will be found 2 $\frac{7}{10}$; as by the ſubſequent operation is manifeſt.

12	7 $\frac{3}{5}$	12 $\frac{1}{5}$	9 $\frac{1}{2}$
$\frac{12}{1}$	$\frac{38}{5}$	$\frac{61}{5}$	$\frac{19}{2}$
60		122	
38		95	
$\frac{22}{5}$ that is 4 $\frac{2}{5}$		$\frac{27}{10}$ that is 2 $\frac{7}{10}$.	

Although the three laſt Rules be ſufficient for all caſes in *Subtraction* of *Fractions*, mixt numbers, or whole and mixt; nevertheleſs the following *Rules* will be more expeditious in the ſubtraction of mixt numbers, or whole and mixt, eſpecially when the Integers conſiſt of many places, as will be manifeſt by the operation, *viz.*

2. By particular Rules, viz. 1 A whole number from a mixt number.

IV. When a whole number is given to be ſubtracted from a mixt number, ſubtract the ſaid whole number from the Integer or Integers of the mixt number (as is taught by the Rules of the fourth Chapter) and unto the remainder annex the fractional part of the mixt number given, ſo is the mixt number thus found, the remainder or difference ſought.

$$\begin{array}{r} 24\,\frac{5}{8} \\ \underline{7\phantom{\,\frac{5}{8}}} \\ 17\,\frac{5}{8} \end{array}$$

As if 7 be given to be ſubtracted from 24 $\frac{5}{8}$, the remainder will be 17 $\frac{5}{8}$ as by the operation is manifeſt.

2. A Fraction from an Integer.

V. When a fraction is given to be ſubtracted from an Integer, ſubtract the Numerator from the Denominator, and place that which remains over the Denominator, which new fraction thus found, is the remainder or difference ſought.

So $\frac{3}{5}$ being ſubtracted from an Integer, or 1, the remainder is $\frac{2}{5}$: Alſo $\frac{13}{19}$ being ſubtracted from 1, the remainder is $\frac{6}{19}$.

3. A Fraction from a whole number greater than 1.

VI. When a fraction is given to be ſubtracted from a whole number greater than 1, ſubtract the ſaid fraction from one of the Integers given by the laſt Rule;) ſo the remaining fraction being annexed to the number of Integers leſſened by unity or 1, gives the remainder or difference ſought.

Thus $\frac{5}{7}$ being ſubtracted from 17, the remainder is 16 $\frac{2}{7}$; alſo $\frac{7}{12}$ being ſubtracted from 39, the remainder is 38 $\frac{5}{12}$.

4. A mixt number from a whole number.

VII. When a mixt number is given to be ſubtracted from a whole number, ſubtract firſt of all (by the fifth Rule of this Chapter) the fractional part of the mixt number from an Integer borrowed from the whole number given, and ſet down the remaining fraction, then adding the Integer borrowed unto the Integer or Integers of the mixt number, ſubtract the ſaid ſum from the whole number given (as is taught in ſubtraction of whole numbers;) ſo that which remains, together with the remaining fraction before found, is the remainder or difference ſought.

So if 9 $\frac{7}{12}$ be ſubtracted from 50, the remainder is 40 $\frac{5}{12}$, as by the operation is manifeſt.

$$\begin{array}{r} 50 \\ 9\,\frac{7}{12} \\ \hline 40\,\frac{5}{12} \end{array}$$

5. A Fraction from a mixt Number by this and the next Rule.

VIII. When a fraction is given to be ſubtracted from a mixt number, and the ſaid fraction is leſs than the fractional part of the mixt number, ſubtract the leſſer fraction from the greater by the firſt or ſecond Rule of this Chapter, then the remaining fraction being annexed to the Integer or Integers of the mixt number, gives the remainder or difference ſought.

So $\frac{5}{9}$ being ſubtracted from 12 $\frac{7}{8}$ the remainder is 12 $\frac{23}{72}$, as by the operation is manifeſt.

$$\begin{array}{r} 12\,\frac{7}{8} \\ 0\,\frac{5}{9} \\ \hline 12\,\frac{23}{72} \end{array}$$

IX. When a Fraction is given to be ſubtracted from a mixt Number, and the ſaid Fraction is greater than the fractional part of the mixt number, ſubtract the ſaid greater Fraction from an Integer borrowed from the mixt Number (by the fifth Rule of this Chapter) and add the remaining Fraction unto the fractional part of the mixt Number (by the firſt or ſecond Rule of the eighteenth Chapter) ſo the Fraction found by that addition, being annexed to the Integers of the mixt Number leſſened by an Integer, or 1, gives the remainder or difference ſought.

Thus $\frac{5}{9}$ being ſubtracted from 13 $\frac{3}{8}$, the remainder 12 $\frac{59}{72}$, *viz.* ſubtracting $\frac{5}{9}$ from 1, the remainder is $\frac{4}{9}$, which added to $\frac{3}{8}$ gives $\frac{59}{72}$, which being annexed to 12 (the number of Integers in the mixt number leſſened by 1 or unity) gives 12 $\frac{59}{72}$ the remainder ſought.

$$\begin{array}{r} 13\,\frac{3}{8} \\ 0\,\frac{5}{9} \\ \hline 12\,\frac{59}{72} \end{array}$$

6. A mixt number from a mixt number by this and the next Rule.

X. When a mixt number is given to be ſubtracted from a mixt Number, and the fractional part of the mixt number to be ſubtracted, is leſs than the fractional part of the mixt Number from which you are to ſubtract, ſubtract the ſaid leſſer Fraction from the greater (by the firſt or ſecond Rule of this Chapter) and ſet down the remaining Fraction: Alſo ſubtract the Integers of the leſſer mixt Number from the Integers of the greater (as in ſubtraction of whole Numbers,) ſo is the mixt Numbhr thus found, the remainder or difference ſought.

So if $17\frac{3}{8}$ be given to be ſubtracted from $20\frac{5}{7}$, the remainder will be found $3\frac{19}{56}$, *viz.* ſubtracting $\frac{3}{8}$ from $\frac{5}{7}$, the remainder is $\frac{19}{56}$, alſo ſubtracting 17 from 20, the remainder is 3.

$$\begin{array}{r} 20\frac{5}{7} \\ 17\frac{3}{8} \\ \hline 3\frac{19}{56} \end{array}$$

XI. When a mixt Number is given to be ſubtracted from a mixt Number, and the fractional part of the mixt number to be ſubtracted is greater than the fractional part of the mixt Number from which you are to ſubtract, ſubtract the ſaid greater Fraction from an Integer borrowed from the greater mixt Number (by the fifth Rule of this Chapter) and add the remaining Fraction unto the fractional part of the greater mixt Number (by the firſt or ſecond Rule of the 18th Chapter;) ſo is the ſum to be reſerved as the fractional part of the remainder ſought; then add the Integer borrowed unto the Integer or Integers of the leſſer mixt Number, and ſubtract the ſum from the Integers of the greater mixt Number (as in ſubtraction of whole Numbers;) ſo that which remains, together with the Fraction before reſerved, is the remainder or difference ſought.

Thus if $20\frac{7}{8}$ be given to ſubtracted from $35\frac{3}{5}$ the remainder will be found $14\frac{29}{40}$, *viz.* ſubtracting $\frac{7}{8}$ from an Integer or 1, the remainder is $\frac{1}{8}$, which added to $\frac{3}{5}$ gives $\frac{29}{40}$, then adding theInteger borrowed unto 20, it will be 21, which ſubtracted from 35, the remainder is 14, ſo the remainder or difference ſought is $14\frac{29}{40}$.

$$\begin{array}{r} 35\frac{3}{5} \\ 20\frac{7}{8} \\ \hline 14\frac{29}{50} \end{array}$$

When you cannot clearly diſcern which is the greater of two Fractions, having unequal Denominators, reduce them into Fractions of the ſame value which ſhall have a common Denominator (by the fourteenth Rule of the ſeventeenth Chapter) and then it will be apparent which of the two Fractions is the greater. As if it be deſired to know which of theſe two Fractions $\frac{6}{7}$ and $\frac{11}{13}$ is the greater, after they are reduced to $\frac{78}{91}$ and $\frac{77}{91}$, it is evident that the former exceeds the latter by $\frac{1}{91}$.

To diſcern the greater of two fractions.

CHAP. XX.

Multiplication of Vulgar Fractions, and mixt Numbers.

I. When the Numbers given to be multiplied are both ſingle Fractions, multiply the Numerators one by the other, and take the Product for a new Numerator; alſo multiply the Denominators one by the other, and the Product is a new Denominator, which new Fraction is the Product ſought.

To multiply ſingle fractions.

So $\frac{7}{12}$ and $\frac{5}{8}$ being given to be multiplied, the Product will be found $\frac{35}{96}$, for 7 multiplied by 5 produceth 35 for a new Numerator, and 12 multiplied by 8 produceth 96 for a new Denominator: alſo $\frac{5}{7}$ and $\frac{3}{7}$ being multiplied one by the other, the Product will be found $\frac{15}{49}$. Here you may obſerve that in the multiplication of proper *Fractions*, the Product is always leſs than either of the terms given; for in multiplication ſuch proportion as unity or 1 hath to either of the terms given, the ſame proportion hath the other term to the Product.

To multiply mixt Numbers,

II. When one of the numbers given is a whole number or a mixt number; alſo, when both of them are mixt numbers, reduce ſuch whole number, mixt number or numbers into an improper Fraction or Fractions by the tenth or eleventh Rules of the ſeventeenth Chapter, and then the operation will be the ſame as in the laſt Rule.

So $8\frac{2}{3}$ being given to be multiplied by 5, the Product will be found $43\frac{1}{3}$; *viz.* $8\frac{2}{3}$ being reduced into the improper Fraction $\frac{26}{3}$: Alſo 5 unto $\frac{5}{1}$ multiply 26 by 5, the Product is 130 for a new Numerator: Alſo multiplying 3 by 1, the Product is 3 for a new Denominator, which new Fraction $\frac{130}{3}$ being reduced (according to the thirteenth Rule of the ſeventeenth Chapter) will be $43\frac{1}{5}$ the Product ſought. In like manner $7\frac{1}{2}$ being multiplied by $5\frac{3}{5}$, the Product will be found 42. Here obſerve, that when either of the terms given is a compound Fraction, it is firſt of all to be reduced into a ſingle Fraction, and then the operation is as before.

Note, 1. Sometimes the work of Multiplication in Fractions may be very uſefully contracted by this following Rule, *viz.*

When two Fractions propos'd to be multiplied (whether they be proper or improper) are ſuch, that the Numerator of the one, and the Denominator of the other, may be ſeverally divided by ſome common Diviſor without a remainder; you may take the Quotients inſtead of the ſaid Numerator and Denominator, and then multiply as before in the firſt Rule of this Chapter: As for example, if, $\frac{6}{7}$ be to be multiplied by $\frac{5}{12}$; becauſe 6 the Numerator of the firſt, and 12 the Denominator of the latter Fraction, being ſeverally divided by their common Diviſor 6, give the Quotients 1 and 2, I ſet theſe (or imagin them to be ſet) in the places of 6 and 12; by which exchange there ariſe $\frac{1}{7}$ and $\frac{5}{2}$, theſe multiplied one by the other (according to the firſt Rule of this Chapter) produce $\frac{5}{14}$ the deſired Product of $\frac{6}{7}$ into $\frac{5}{12}$, in the ſmalleſt terms.

Again to multiply $\frac{16}{48}$ by $\frac{3}{16}$; becauſe the Numerator of the firſt Fraction and the Denominator of the latter, being each divided by 16 give the Quotients 1 and 1, I ſet 1 and 1 in the places of 16 and 16; likewiſe becauſe 48 the Denominator of the firſt, and 3 the Numerator of the latter Fraction, being each divided by their common Diviſor 3, give 16 and 1, I take 16 and 1 inſtead of 48 and 3, ſo by thoſe exchanges there ariſe $\frac{1}{16}$, and $\frac{1}{1}$, which multiplied one by the other produce $\frac{1}{16}$, which is the Product in the ſmalleſt terms made by the multiplication of $\frac{16}{48}$ into (or by) $\frac{3}{16}$.

2. To take any part or parts of a Number propounded, is nothing elſe but to multiply the ſaid Number by the Fraction which declareth what part is to be taken: So if you deſire to know what is $\frac{5}{8}$ of 320, multiply $\frac{320}{1}$ by $\frac{5}{8}$, or $\frac{40}{1}$ by $\frac{5}{1}$, and the Product will be 200. In like manner $\frac{2}{3}$, of 45 $\frac{3}{8}$ is 30 $\frac{1}{4}$ Alſo $\frac{1}{4}$ of 120 is 30.

3. Sometimes the work of multiplication in mixt

numbers may be compendiouſly performed after the manner of theſe following examples, *viz.* if it be required to multiply 120 $\frac{1}{4}$ by 48 $\frac{1}{2}$, firſt multiply the whole numbers mutually, to wit, 120 by 48, and place the particular products orderly one under the other as in Multiplication of whole numbers; then multiply the ſaid whole numbers firſt given by the Fractions alternately, *viz.* take $\frac{1}{4}$ of 48 which is 12, alſo take $\frac{1}{2}$ of 120 which is 60, and place the ſaid 12 and 60 orderly to be added to the former particular Products: Laſtly, add altogether, and to the ſum annex the product of the two Fractions, to wit in this example, the product of the Multiplication of $\frac{1}{4}$ by $\frac{1}{2}$, which is $\frac{1}{8}$, ſo the total Product required will be 5832 $\frac{1}{8}$, as you ſee by the example in the Margin. In like manner, if 18 $\frac{1}{2}$ be multiplied by 40 $\frac{1}{3}$, the Product will be 746 $\frac{1}{6}$; and if 29 $\frac{1}{2}$ be multiplied by 50, the Product will be 1475, as you ſee by the examples following.

120 $\frac{1}{4}$
48 $\frac{1}{2}$
960
480
12
60
5832 $\frac{1}{8}$

18 $\frac{1}{2}$	29 $\frac{1}{2}$
40 $\frac{1}{3}$	50
720	1450
20	25
6	1475
746 $\frac{1}{6}$	

4. When a Fraction is to be multiplied by a number which happens to be the ſame with the Denominator, take the Numerator for the Product; ſo if this Fraction $\frac{3}{4}$ be propounded to be multiplied by the Denominator 4, the Product will

be $\frac{12}{4}$ that is 3, which is the ſame with the Numerator 3. In like manner if $\frac{5}{8}$ be multiplied by the Denominator 8, the Product is equal to 5 the Numerator of the ſaid $\frac{5}{8}$.

CHAP. XXI.

Diviſion of Vulgar Fractions and mixt Numbers.

I. WHen the numbers given are both ſingle Fractions, multiply the Denominator of the Diviſor by the Numerator of the Dividend; and take the Product for a new Numerator: alſo multiply the Numerator of the Diviſor by the Denominator of the Dividend, and the Product is a new Denominator; which new Fraction is the quotient ſought.

The Diviſion of ſingle fractions.

So if $\frac{4}{9}$ be given to be divided by $\frac{3}{5}$, the quotient will be found $\frac{20}{27}$; *viz.* multiplying 5 by 4 the Product is 20 for a new Numerator, alſo multiplying 3 by 9, the Product is 27 for a new Denominator, ſo is $\frac{20}{27}$ the quotient ſought; in like manner if $\frac{5}{8}$ be given to be divided by $\frac{2}{7}$, the quotient will be found $\frac{35}{16}$, that is, 2 $\frac{3}{16}$, as you ſee in the Example: here you may obſerve, that in Diviſion by proper fractions, the quotient is always greater than either of the Fractions given; for in Diviſion, as the Diviſor is in proportion to 1 or unity, ſo is the Dividend to the Quotient.

$\frac{3}{5}$) $\frac{4}{9}$ ($\frac{20}{27}$

$\frac{2}{7}$) $\frac{5}{8}$ ($\frac{35}{16}$

II. When one of the numbers given is a whole number or a mixt number; alſo when both are mixt numbers, reduce ſuch whole number or mixt number or numbers into an improper fraction or fractions, by the tenth or eleventh Rule of the ſeventeenth Chapter, and then the operation will be the ſame as in the laſt Rule.

So if 42 be divided by $7\frac{1}{2}$. the quotient will be found $5\frac{3}{5}$, for $7\frac{1}{2}$ and 42 will be reduced into theſe improper fractions $\frac{15}{2}$ and $\frac{42}{1}$, then multiplying 42 by 2, the Product is 84 for a new Numerator, alſo multiplying 15 by 1, the product is 15 for a new Denominator, ſo is $\frac{84}{15}$. the quotient ſought, which is equal to $5\frac{3}{5}$ (as is evident by the thirteenth Rule of the ſeventeenth Chapter.) In like manner, if $6\frac{1}{2}$ be divided by $3\frac{2}{5}$, the quotient will be $1\frac{31}{34}$. Alſo if $5\frac{1}{3}$ be divided by $12\frac{1}{2}$ the quotient will be $\frac{32}{75}$.

$7\frac{1}{2})\ 42\ ($

$\frac{15}{2})\ \frac{42}{1}\ (\frac{84}{15}$

$15)84(5\frac{9}{15}$

Note, Sometimes the Work of Diviſion in Fractions may be very uſefully contracted by this following Rule, *viz.* When either the two Numerators, or the two Denominators of the Fractions propoſed can be divided ſeverally by ſome common Diviſor without a remainder, you may take the Quotients inſtead of the ſaid Numerators or Denominators, and then divide by the firſt Rule of this Chapter: As for Example if $\frac{12}{17}$ be to be divided by $\frac{8}{15}$, becauſe the Numerators 12 and 8 being each divided by their common Diviſor 4 will give the Quotients 3 and 2; I take theſe inſtead of 12 and 8, by which exchange there ariſe $\frac{3}{17}$ and $\frac{2}{15}$, the former of which being divided by the latter, (according

ing to the firſt Rule of this Chapter) given $\frac{15}{34}$, which is the Quotient in the leaſt terms that ariſeth by dividing $\frac{12}{17}$ by $\frac{8}{5}$.

Again, to divide $\frac{25}{8}$ by $\frac{15}{8}$; becauſe the Numerators 25 and 15 being ſeverally divided by their common Diviſor 5 give the Quotients 5 and 3, likewiſe becauſe the Denominators 8 and 8 being each divided by 8 give the Quotients 1 and 1, I ſet 5 and 3 in the places of the Numerators 25 and 15, alſo 1 and 1 in the places of the Denominators 8 and 8, whence ariſe $\frac{5}{1}$ and $\frac{3}{1}$. Laſtly, dividing $\frac{5}{1}$ by $\frac{3}{1}$, that is 5 by 3, there ariſeth $\frac{5}{3}$, that is 1 $\frac{2}{3}$, which is the deſired Quotient of $\frac{25}{8}$ divided by $\frac{15}{8}$.

Queſtions to exerciſe the Rules of Vulgar Fractions before delivered.

Queſt. 1. The difference of two numbers is 1 $\frac{13}{24}$, the leſſer number is 2 $\frac{1}{8}$, what is the greater? *Anſw.* 3 $\frac{2}{3}$, (found by *Addition.*)

Q. 2. What number is that, which if added to 3 $\frac{5}{8}$ gives the ſum 8 $\frac{23}{88}$? *Anſ.* 4 $\frac{7}{11}$ (found by *Subtraction*)

Queſt. 3. There is in three Bags the ſum of 121 $\frac{9}{20}$ *l.* *viz.* in the firſt Bag 50 $\frac{5}{8}$ *l.* in the ſecond 40 $\frac{4}{15}$ *l.* what is in the third Bag? *Anſwer* 30 $\frac{1}{2}$ *l.* (found by *Addition* and *Subtraction.*)

Queſt. 4. Two Merchants *A* and *B*, having certain ſhares in a Ship, the ſhare of *A* is $\frac{7}{10}$ of the Ship, that of *B* $\frac{9}{13}$, what is the difference between their parts? *Anſw.* The ſhare of *A* exceeds the ſhare of *B* by $\frac{1}{130}$. (found by *Subtraction.*)

Queſt. 5. What is $\frac{5}{8}$ of 130 $\frac{2}{3}$? *Anſw.* 81 $\frac{2}{3}$ (found by *Multiplication.*)

Queſt. 6. What number is that, which being multiplied by $\frac{3}{5}$ produceth 25 $\frac{2}{5}$? *Anſw.* 42 $\frac{1}{3}$ (found by *Diviſion.*)

Now followeth the Doctrine of *Decimal Fractions.*

The Doctrine of Decimal Fractions.

CHAP. XXII.

Notation of Decimal Fractions.

1. IT is hard to determine, who was the firſt that brought *Decimal Arithmetick* to light, though it be a late Invention; but without doubt it hath received much improvement within the compaſs of a few years, by the induſtry of *Artiſts*, and now ſeems to be arrived at perfection. The excellency thereof is beſt known to ſuch as can apply it to the practical part of the *Mathematicks*, and to the Conſtruction of *Tables*, which depend upon ſtanding or conſtant proportions, ſuch are *Trigonometrical Canons*, *Tables* for computing of *compound Intereſt*, &c. in which caſes decimal operations do afford ſo great help, (that in my opinion) many Ages have not produced a more uſeful invention. But it may be objected, that *Decimal Arithmetick* for the moſt part gives an imperfect ſolution to a queſti-

The proper uſe of Decimal Arithmetick.

a queſtion. This I grant, yet the anſwer ſo given may be as uſeful as that which is exactly true; for in common affairs, the loſs of $\frac{1}{1000}$ part of a *Grain*, or of an *Inch*, &c. to wit, any quantity which cannot be ſeen, is inconſiderable: but I would not be miſtaken, for in extolling *Decimals* I do not cry down *Vulgar Fractions*, ſince experience ſheweth that *Decimal Fractions* are commonly abuſed, by being applyed to all manner of queſtions about *Money*, *Weight*, &c. when indeed many queſtions may be reſolved with much more facility by *Vulgar Arithmetick*, as may partly appear by this Example, *viz.* at 9*l.*——6*s.*——8*d.* the hundred weight of Tobacco, what will 987 hundred weight coſt? *Anſw.* 9212*l.* which by the common *Rule* of *Practice* by *Aliquot parts* is found out in a quarter of the time, that will neceſſarily be required to work it by *Decimals*, which at laſt will give an imperfect Anſwer; I might inſtance the like inconvenience divers ways, were it not for loſs of time; ſo that the right uſe of *Decimals* depends upon the diſcretion of the *Artiſt*.

Decimal Fractions ſometimes abuſed.

II. When a ſingle Fraction hath for its denominator a number conſiſting of 1 or unity in the extream place towards the left hand, and nothing but a Cypher or Cyphers towards the right, it is more particularly called a *Decimal*: of this kind are theſe that follow, $\frac{5}{10}$, that is, five tenths; $\frac{5}{100}$, five hundredth parts; likewiſe theſe are decimal fractions, $\frac{34}{1000}$, $\frac{205}{10000}$, $\frac{1023}{10000}$, &c.

The definition of a Decimal Fraction.

III. A Decimal fraction may be expreſs'd without

out the Denominator, by prefixing a point or comma before (to wit, on the left hand of) the Numerator, ſo $\frac{5}{10}$ may be written thus .5 or thus ,5, $\frac{25}{100}$ thus, .25, or thus, ,25.

IV. In Decimals when the Numerator conſiſts not of ſo many places as the Denominator hath Cyphers, fill up the void places in the Numerator with Cyphers prefixed on the left hand: ſo $\frac{5}{100}$ is written thus, .05; likewiſe $\frac{50}{1000}$ thus .050; and $\frac{205}{10000}$, thus, .0205, likewiſe $\frac{6}{1000}$, thus .006.

V. In Decimals thus expreſs'd, the Denominator is diſcoverable by the places of the Numerator: for if the Numerator conſiſts of one place, the Denominator conſiſts of 1 or unity with one Cypher; if of two places, the Denominator conſiſts of 1 with two Cyphers annexed; if of three, the Denominator conſiſts of 1 or unity with three Cyphers annexed: ſo the Denominator of .25 is 100; the Denominator of .050 is 1000, and the Denominator of .096 is 1000.

VI. Cyphers at the end of a Decimal do neither augment nor diminiſh the value thereof: ſo .2, .20, .200, .2000 are *decimals*, which have one and the ſame value, for $\frac{20}{100}$ being abbreviated by the eighth *Rule* of the ſeventeenth *Chapter*, will be made $\frac{2}{10}$ and ſo will $\frac{200}{1000}$ or $\frac{2000}{10000}$.

VII. Wherefore Decimal fractions are eaſily reduced to a common Denominator (which is a troubleſome work in *Vulgar Fractions*;) for if all the Numerators of as many decimal fractions as are given, be made to conſiſt of the ſame number of places, by annexing a Cypher or Cyphers at the end (that is on the right hand) of ſuch Numerators as are defective, they will all be reduced to a common Denominator, ſo theſe *Decimals*, .2, .03, .027, (which ſignifie $\frac{2}{10}$, $\frac{3}{100}$, $\frac{27}{1000}$) may be reduced into theſe, .200, .030, .027, which have a 1000 for a common Denominator.

VIII. The order of places in any Decimal proceedeth from the left hand to the right, contrary to the order of places in Integers, which is from the right hand to the left: ſo in this *Decimal*, .247 the figure 2 ſtandeth in the firſt place (being the outermoſt towards the left hand, and next to the point,) the figure 4 ſtandeth in the ſecond *place*, and 7 in the *third*. Alſo in this *Decimal* .0245, a Cypher ſtands in the *firſt place*, 2 in the *ſecond*, 4 in the *third*, and 5 in the *fourth*.

IX. Every place in the Numerator of a Decimal Fraction hath a peculiar Denominator, or proper value, *viz.* the Denominator of the firſt place is 10, of the ſecond, 100; of the third, 1000, *&c.* ſo that the firſt place of a Decimal ſignifies tenth parts of an Unite or Integer; the ſecond place, hundreth parts of an Integer; the third place, thouſandth parts of an Integer, *&c.* Hence it is manifeſt, that this *Decimal* .3254 (every place thereof being conſidered apart by it ſelf) conſiſts of .3, .02, .005, .0004, (*viz.* $\frac{3}{10}$, $\frac{2}{100}$, $\frac{5}{1000}$, $\frac{4}{10000}$, which being reduced to a common Denominator (by the ſeventh *Rule* of this *Chapter*) will give theſe, .3000, .0200, .0050, .0004, (to wit, $\frac{3000}{10000}$, $\frac{200}{10000}$, $\frac{50}{10000}$, $\frac{4}{10000}$) all which collectively make .3254 (or $\frac{3254}{10000}$)

X. In whole numbers, the firſt place above (that is on the left hand of) the place of Units, ſignifies Tens of Units; but the firſt place beneath, (that is on the right hand of) the place of Units, ignifies tenth parts of 1 or Unity, and is called the irſt place of Decimal parts, or place of Primes; ikewiſe the ſecond place above the place of Units ignifies hundreds of Units, but the ſecond place beneath the place of Units, ſignifieth hundredth parts of 1 or Unity, and is called the *ſecond place of Decimals*, or place of ſeconds; ſo that as the values of the places in Integers, do aſcend in a decuple proportion from the place of Units towards the left hand, ſo the values of the places of Decimals do deſcend in a ſubdecuple proportion beneath the place of Units towards the right hand: *viz.* Among the places of Integers, every following place towards the left hand, is ten times the value of the next preceding place: But among the places of decimal Parts, every following place towards the right hand is one tenth part of the value of the next preceding place: All which will be evident by the following *Table.*

A TABLE for the Notation of Integers and Decimals.

Integers. of Units.						*Decimal parts.* of 1 or Unity.				
&c.	*Ten Thouſands*	*Thouſands*	*Hundreds*	*Tens*	*Units (under 10)*	*Ten parts*	*Hundredth parts*	*Thouſandth parts*	*Ten Thouſandth parts*	&c.
	7	3	2	8	5	8	2	3	7	
&c.	*Fifth place*	*Fourth place*	*Third place*	*Second place*	*Firſt place*	*Firſt place*	*Second place*	*Third place*	*Fourth place*	&c.

In the foregoing Table you may obſerve, that the places of Integers, or whole numbers are ſeparated from the places of *Decimal parts* of 1 (or unity) by a point; ſo the number on the left hand of the point expreſſeth 73285 Integers or Units, but the number on the right hand of the point expreſſeth only 8237 parts of 1 (or an Integer) ſuppoſed to be divided into 10000 equal parts. In like manner this number 5 .8 ſignifies 5 Integers and eighth tenth parts of an Integer, and this number 285 .82 ſignifies 285 Integers (or Units) and $\frac{82}{100}$ parts of an Integer.

CHAP. XXIII.

Concerning the Reduction of Vulgar Fractions to Decimal Fractions.

I. IF the greateſt Integer of *money*, as alſo of *weight*, *meaſure*, &c. were ſubdivided decimally, to wit, a *pound* of Engliſh money into ten equal pieces of Coin, and every one of theſe into ten other equal pieces, &c. and *weights*, *meaſures*, &c. after the ſame manner: the doctrine of Arithmetick would be taught with much more eaſe and expedition than now it is; but it being improbable that ſuch a reformation will ever be brought to paſs, I ſhall proceed in directing a courſe to the ſtudious for obtaining the frugal uſe of ſuch Decimal fractions as are in his power.

II. Foraſmuch as in *Arithmetical* queſtions, ſome of the given numbers do for the moſt part happen to be fractions, a way muſt be ſhew'd how to reduce a *Vulgar Fraction* to a *Decimal Fraction*; yet in ſome caſes there is no need of this *Reduction*; for example, a *foot* in length is vulgarly ſubdivided into 12 inches, an inch into 4 quarters, and each quarter into two half quarters; but a *foot* may as eaſily, and a great deal more commodiouſly be divided, firſt into ten equal parts, and then each of thoſe into ten other equal parts, and each of theſe into ten other equal parts; (or at leaſt ſuch diviſion may be ſuppoſed or imagined when it cannot actually be made.(This *foot* in length ſo divided being applied to the ſides of *ſuperficial figures*, or of *ſolids* will at firſt ſight give the quantities of lines in *feet* and *decimal parts* of a *foot* (as readily as a *foot* vulgarly divided will ſhew you how many *feet*, *inches*, *quarters*, and *half quarters* are contained in any line) from whence the *ſuperficial* or *ſolid content* may be found in *feet* by *multiplication* only; and how much this excels the *vulgar way*, I ſhall partly manifeſt in the fifth Rule of the 26th Chapter. The like ſubdiviſion I would have to be made of a *Yard*, *Perch*, &c.

How to reduce a vulgar fraction to a decimal fraction.

III. A ſingle fraction, which is no decimal fraction, may be reduced into a decimal of the ſame value, or infinitely near (for all vulgar fractions cannot be exactly reduced to decimals) by the Rule of Three Direct; for as the Denominator of any ſingle fraction whatſoever, is to the Numerator thereof, ſo is any other Denominator to his correſpondent Numerator: *Example*, let it be required to reduce $\frac{5}{8}$ into a Decimal, whoſe Denominator is aſſigned to be 1000, ſay by the *Rule of three*, if the Denominator 8 hath 5 for a Numerator, what will the Denominator 1000 require for

Numerator? Multiply and divide as the Rule of
ree direct doth require, ſo will the fourth *propor-*
nal he found to be 625, which is the Numerator
ught; therefore $\frac{625}{1000}$ or .625, is a *decimal fraction*
ual in value to $\frac{5}{8}$. Another *Example.* Let it be re-
ired to reduce $\frac{7}{240}$ into a *decimal fraction*, whoſe
enominator ſhall be 100000, ſay by the Rule of
ree, if 240 the Denominator give 7 for a Nume-
tor, what will the *Denominator* 100000 require
r a *Numerator?* *Anſw.* 2916 and ſomewhat more;
it that which the ſaid 2916 wants of being a true
umerator is leſs than $\frac{1}{1000}$ part of an Integer,
erefore the *decimal fraction* $\frac{2916}{100000}$ or .02916 is al-
oſt equal to $\frac{7}{240}$, which $\frac{7}{240}$ cannot be exactly re-
uced into a *decimal fraction.* The like will happen
the reduction of moſt *vulgar fractions* to *decimals*,
which caſe, the *Denominator* of the *decimal* muſt
e aſſigned to be ſo great, that what is wanting in
he *Numerator* may be an inconſiderable value.

IV. Upon the aforeſaid ground, the known or
ccuſtomary parts of *Money*, *Weight*, *Meaſure*, *Time*,
c. may be reduced to *decimals*: for if you deſire to
now what *decimal fraction* of a pound *ſterling* is e-
ual in value to one *ſhilling*, conſider firſt that a
ound is the *Integer*, and that 20 *ſhillings* are equal
o that *Integer*, therefore 1 ſhilling is $\frac{1}{20}$ of a pound,
ow if we conceive one pound to be divided into
00000 parts, *viz.* if we aſſign 100000 for the *Deno-*
ninator of a *decimal fraction*, the *Numerator* will be
ound by the laſt Rule to be 5000, ſo that $\frac{5000}{100000}$ or
05000 or .05 (for Cyphers at the end of a *decimal*
re of no uſe, as hath been ſhewn in the 6th *Rule* of
he 22 *Chapter*) is a *decimal fraction* of a *pound*, and is
exactly

actly equal to 1*s.* or $\frac{1}{20}$ part of a pound *ſterling.*

In like manner foraſmuch as 240 *pence* are equal to a pound of Engliſh money, 7 *pence* are $\frac{7}{240}$ parts of a pound, which *fraction* will be reduced into this *decimal* .02916*l.* which is very near equal to $\frac{7}{240}$ for it wants not $\frac{1}{100000}$ part of a pound. Moreover ſince 960 *farthings* are equal to a *pound* Engliſh, one *farthing* is $\frac{1}{960}$ part of a pound, which will be reduced into this *decimal* .00104*l.* very near; but if you pleaſe to proceed near to the truth, you will find this *decimal* .00104166, &c. to anſwer a *farthing*, and ſo by augmenting the Denominator with Cyphers, you may proceed infinitely near, when you cannot attain unto the truth it ſelf. After the ſame method may the vulgar *Sexagenary Fractions* uſed in *Aſtronomy* be reduced to *decimals*; for ſince a *degree* is uſually ſubdivided into ſixty parts called *minutes* or *primes*; a *prime* or *minute* into ſixty parts called *ſeconds*; a *ſecond* into ſixty *thirds*; a third into ſixty *fourths*, &c. and conſequently a *degree* is equal unto 60 *minutes* (or *Primes*) or unto 3600 *ſeconds*, or 216000 *thirds*, or 12960000 *fourths*, &c. It is evident that 7 *minutes* (or *Primes*) are $\frac{7}{60}$ parts of a *degree*, which by the third *Rule* of this *CHAPTER* may be reduced into the Decimal .1166, &c. Alſo 29 *thirds* are $\frac{29}{216000}$ parts of a *degree* which may be reduced into the Decimal .000134, &c. ′ ″ ‴ ⁗

Moreover, 58 : 33 : 14 : 12, that is, 58 *primes*, 33 *ſeconds*, 14 *thirds*, and 12 *fourths* may be reduced to a *decimal* in this manner, *viz.* reduce them all into *fourths* (according to the ſixth *Rule* of the ſeventh *Chapter*) ſo will you find 12647652 *fourths*, which are $\frac{13647652}{13984690}$ parts of a *degree*, which *vulgar fraction* may be reduced into this *decimal* of a *degree*, to wit .975899, &c. (by the third *Rule* of this *Chapter.*)

This to the ingenious will be a ſufficient light for the finding of the *Decimals* congruent to the *ſhillings*, *pence*, and *farthings*, which are under a *pound ſterling*; alſo the *Decimals* of the known parts of *Weight*, *Meaſure*, *Time*, &c. as they are expreſs'd in the following *Table*, wherein you may obſerve, that moſt of the *Decimals* conſiſt of 7 or 8 figures, yet in ordinary practice, you ſhall have occaſion to uſe only the firſt five, and ſometimes fewer.

The TABLE
OF
REDUCTION.

TABLET I.
Of English Money, the Integer being a Pound.

Shillings.	*Decimals of a Pound.*
19	.95
18	.9
17	.85
16	.8
15	.75
14	.7
13	.65
12	.6
11	.55
10	.5
9	.45
8	.4
7	.35
6	.3
5	.25
4	.2
3	.15
2	.1
1	.05

Pence with Farthings.	*Decimals of a Pound.*
	.0489583
	.0479166
	.046875
11	.0458333
	.0447916
	.04375
	.0427082
10	.0416666
	.040625
	.0395833
	.0385416
9	.0375
	.0364583
	.0354166
	.034375
8	.0333333
	.0322916
	.03125
	.0302083
7	.0291666
	.028125
	.0270833
	.0260416
6	.025

	.0239583
	.0229166
	.021875
5	.0208333
	.0197916
	.01875
	.0177708
4	.0166666
	.015625
	.0145833
	.0135416
3	.0125
	.0114583
	.0104166
	.009375
2	.0083333
	0072916
	00625
1 Pen. & 1 Far.	.0052083
Penny	.0041666
3. Farth.	.003125
2. Farth.	.0020833
1. Farth.	.0010416

TABLET II.

Of Troy weight, the Integer being an Ounce.

Penny weights.	*Decimals of an Ounce.*
19	.95
18	.9
17	.85
16	.8
15	.75
14	.7
13	.65
12	.6
11	.55
10	.5
9	.45
8	.4
7	.35
6	.3
5	.25
4	.2
3	.15
2	.1
1	.05

Grains	*Decimals of an Ounce*
23	.0479166
22	.0458333
21	.04375
20	.0416666
19	.0395833
18	.0375
17	.0354166
16	.0333333
15	.03125
14	.0291666
13	.0270833
12	.025
11	.0229166
10	.0208333
9	.01875
8	.0166666
7	.0145833
6	.0125

5	.0104166
4	.0083333
3	.00625
2	.0041666
1	.0020833

TABLET III.

Of Averdupois great weight, the Integer being an hundred weight, to wit, 112 pounds.

quarters of 1 hundred.	*decimals of 1 hundred.*
3	.75
2	.5
1	.25

Pounds	*decimals of 1 hundred.*
27	.2410714
26	.2321428
25	.2232142
24	.2142857
23	.2053571
22	.1964285
21	.1875
20	.1785714
19	.1696428
18	.1607142
17	.1517857
16	.1428571
15	.1339285
14	.125
13	.1160714
12	.1071428
11	.0982142
10	.0892857
9	.0803571
8	.0714285
7	.0625
6	.0535714
5	0446428
4	.0357142
3	.0267857
2	.0178571
1	.0089285

Ounces.	*decimals of 1 hundred.*
15	.0083705
14	.0078125
13	.0072544
12	.0066964
11	.0061383
10	.0055803
9	.0050223
8	.0044642
7	.0039062
6	.0033482
5	.0027901
4	.0022321
3	.0016741
2	.0011160
1	.0005580

quarters of 1 Ounce.	*decimals of 1 hundred.*
3	.0004185
2	.0002790
1	.0001395

TABLET

TABLET IV.

Of Avoirdupois little weight, the Integer being a Pound.

Ounces.	*Decimals of a Pound.*
15	.9375
14	.875
13	.8125
12	.75
11	.6875
10	.625
9	.5625
8	.5
7	.4375
6	.375
5	.3125
4	.25
3	.1875
2	.125
1	.0625

Drams.	*Decimals of a Pound.*
15	.05859375
14	.0546875
13	05078115
12	.046875
11	.04296875
10	.0390625
9	.03515625
8	.03125
7	.02734375
6	.0234375
5	01953125
4	.015625
3	.01171875
2	.0078125
1	.00390625

Quarters of a Dram	*Decimals of a Pound.*
3	.0029296
2	.0019531
1	.0009765

TABLET V.

Of liquid Meaſures, the Integer being a Gallon.

Pints.	*Decimals of a Gallon.*
7	.875
6	.75
5	.625
4	.5
3	.375
2	25
1	.125

Quarters of a Pint.	*Decimals of a Gallon.*
3	.09375
2	.0625
1	.03125

TABLET VI.

Of dry meaſures, the Integer being a Quarter.

Buſhels.	*decimals of a quarter.*
7	.875
6	.75
5	.625
4	.5
3	.375
2	.25
1	.125

Pecks.	*decimals of a quarter.*
3	.09375
2	.0625
1	.03125

Quarters of a peck.	*decimals of a quarter.*
3	.0234375
2	.015625
1	.0078125

Pints.	*decimals of a quarter*
3	.005859
2	.003906
1	.001953

TABLET VII.

Of long meaſures, one Yard or one Ell being the Integer.

quarters of 1 yard or 1 ell.	*decimals of 1 yard or 1 ell.*
3	.75
2	.5
1	.25

Nails.	*decimals of 1 ya.*
3	.1875
2	.125
1	.0625

quarters of nails.	*decimals of 1 ya.*
3	.046875
2	.03125
1	.015625

TABLET VIII.

Of the Reduction of Inches, &c. to decimals, the Integer being a foot in length.

Inches.	*decimals of a foot.*
11	.9166666
10	.8333333
9	.75

8

8	.6666666
7	.5833333
6	.5
5	.4166666
4	.3333333
3	.25
2	.1666666
1	.0833333

quarters of an Inch.	*decimals of a foot.*
3	.0625
2	.0416666
1	.0208333
half a quart. of an Inch.	.0104166

TABLET IX.

Of dozens, the Integer being a groſs.

dozens.	*decimals of a groſs.*
11	.9166666
10	.8333333
9	.75
8	.6666666
7	.5833333
6	.5
5	.4166666
4	.3333333
3	.25
2	.1666666
1	.0833333

parts of a dozen.	*decimals of a groſs.*
11	.076388
10	.069944
9	.0625
8	.055555
7	.048611
6	.041666
5	.034722
4	.027777
3	.020833
2	013888
1	.006944

TABLET X.

Of Time, a day being the Integer.

Hours.	*decimals of a day.*
23	.9583333
22	.9166666
21	.875
20	.8333333
19	.7916666
18	.75
17	.7083333
16	.6666666
15	.625
14	.5833333
13	.5416666
12	.5
11	.4583333
10	.4166666

9	.375
8	.3333333
7	.2916666
6	.25
5	.2083333
4	.1666666
3	.125
2	.0833333
1	.0416666

Minutes.	*decimals of a day.*
59	.0409722
58	.0402777
57	.0395833
56	.0388888
55	.0381944
54	0375
53	.0368055
52	.0361111
51	.0354166
50	.0347222
49	.0340277
48	.0333333
47	.0326388
46	.0319444
45	.0312500
44	.0305555
43	.0298611
42	.0291666
41	.0284722
40	.0277777
39	.0270833
38	.0263888
37	.0256944
36	.0249999
35	.0243055
34	.0236111
33	.0229166
32	.0222222
31	.0215277
30	0208333
29	.0201388
28	.0194444
27	.01875
26	.0180555
25	.0173611
24	.0166666
23	.0159722
22	.0152777
21	.0145833
20	.0138888
19	.0131944
18	.0125
17	.0118055
16	.0111111
15	.0104166
14	.0097222
13	.0090277
12	.0083333
11	.0076388
10	.0069444
9	.00625
8	.0055555
7	.0048611
6	.0041666
5	.0034722

4	.0027777
3	.0020833
2	.0013888
1	.0006944

V. This Table aforegoing conſiſts of ten ſeveral Tablets, of which the firſt (intituled *Engliſh money)* contains in the firſt column thereof the particular Fractions *(viz.* the *ſhillings, pence,* and *farthings)* of a pound *ſterling* ; and in the other column the *decimals,* unto which they may be reſpectively reduced: So in the ſame Tablet .65 is the *decimal,* anſwerable to 13*s.* .0208333 to 5*d.* and .003125 to 3*f.* Likewiſe, .0489583 is the decimal of 11*d.* together with 3 farthings: Alſo .03125 is the decimal of 7 pence half-penny.

Tablet 1. Of Engliſh money.

VI. The next Tablet (intituled *Troy weight(* contains in the firſt column thereof the particular fractions (*viz.* the *Penny weights,* and *Grains)* of an ounce *Troy* and in the other their reſpective decimals: ſo .6 is the correſpondent *decimal* of 12 penny weight, and .0020833 of 1 grain. Likewiſe .025 is the decimal of 12 grains.

2. Of Troy weight.

VII. The third Tablet (intituled *Averdupois great weight)* contains in the firſt colum thereof the Fractions (*viz.* the *Quarters, Pounds, Ounces,* and the *Quarters of an Ounce* of an Hundred according to *Averdupois weight,* and in the other their proper decimals: ſo .5 is the *decimal* of two quarters or half a hundred,.1517857 of 17 pounds

3. Of Averdupois great weights.

.0033482 of 6 ounces, and .0004185 the *decimal* of 3 quarters of an ounce.

4. *Of Averdupois little weight.*

VIII. The fourth (intituled *Averdupois little weight*) ſheweth you the fractions (*viz.* the *Ounces*, *Drams*, and *quarters* of a *dram*) of a pound *Averdupois*, together with their reſpective decimals: So the *decimal* of 3 Ounces is .1875, the *decimal* of 9 Drams is .03515625, and the *decimal* of one quarter of a Dram is .0009765.

5. *Of Liquid meaſures.*

IX. The fifth (intituled *Liquid meaſures*) hath the fractions (*viz.* the *Pints* and *quarters* of a *pint*) of a Gallon, and likewiſe their ſeveral decimals: So the *decimal* of 5 Pints is .625, and the *decimal* of two quarts or half a pint is .0625.

6. *of Dry meaſures.*

X. The ſixth (intituled *Dry meaſures*) gives you the fractions (*viz.* the *Buſhels*, *Pecks*, *quarters* of *Pecks* and *Pints*) of a quarter, together with their peculiar decimals: ſo .375 is the *decimal* of three Buſhels, .03125, of one Peck, .0234375 of $\frac{3}{4}$ of a Peck, and .003906 of two Pints.

7. *Of Long meaſures.*

XI. The ſeventh (intituled *Yards* and *Ells*) offers you the fractions (*viz.* the *Quarters* of *Yards* and *Ells*, *Nails*, and *quarters* of *Nails of Yards*) and their reſpective decimals: ſo .25 is the *decimal* of one quarter of a Yard or Ell, .125 of two Nails, and .046875 of three quarters of a Nail.

XII. The eighth (intituled *Reduction of Inches, &c. to decimals of a foot*) preſents unto you the fractions (to wit, the Inches, quarters and half quarters of an Inch) of a *foot*, together with their correſpondent decimals: So .4166666 is the *decimal* of 5 Inches, .0625 of $\frac{3}{4}$ of an Inch, and .0104166 of $\frac{1}{8}$ or half a quarter of an Inch.

8. Of things accompted by the dozen.

XIII. The ninth Tablet (intituled *Dozens*) yields you the Fractions (*viz.* the Dozens and particulars) of a Groſs, as alſo their reſpective *decimals*: ſo .25 is the *decimal* of 3 Dozen and .048611 of 7 particulars.

9. Of Time.

XIV. The tenth and laſt Tablet (intituled *Time*) gives you the Fractions (*viz.* the Hours and Minutes) of a Day: So .625 is the *Decimal* of 15 hours .0375 of 54 minutes, and .0006944 of one minute.

The uſe of the ſame Table for the Reduction.

1. Of ſingle fractions to decimals.

XV. When a ſingle Fraction of any of the premiſed Tablets is propounded to be reduced to a decimal, find it in the firſt Column of the Tablet, unto which it belongs; this done, juſt againſt that Fraction ſo found, you ſhall have the decimal required: So 13 *s.* being propounded, taking the firſt premiſed *Tablet*, I find 13*s.* in the firſt Column of the Tablet of *money*, and juſt againſt the ſame thirteen ſhillings, I *obſerve* .65, before which having prefixed a point, and by that means ſigned it for a *decimal* (according to the third Rule of the 22 Chapter of this Book) I conclude the ſame .65 ſo ordered, to be the correſpondent *decimal* of thirteen ſhillings the fraction propounded. In like manner .0229166 is the *decimal* of 11 grains in the Tablet of *Troy* weight; and ,0357142 the *decimal* of 4 *lb.* in the Tablet of *Averdupois great weight*, &c.

XVI. When two or more Fractions are propounded, and it is required to find a decimal equivalent unto the ſum of them, find the decimal of each of the Fractions given according to the laſt Rule; then adding together the decimals ſo found that intire ſum is the decimal ſought: *So* 13*s* 5*d.* being produced to a decimal, is 670833; for the decimal of 13*s* is .65, and the decimal of 5*d* .020833, which being added together (by the ſecond Rule of the 24th Chapter of this Book) amount to .670833, *viz.* the decimal which repreſents 13*s* 5*d.* the Fraction propounded: In like manner the decimal of 9 peny weight, and 13 Grains is .4770833, and the decimal of ½ *C.* 19 *lb.* 7 Ounces, is .67354, &c.

13*s.*	.65
5*d.*	.020833
	.670833
9 *p.w.*	.45
13 *gr.*	.027083
	.477083
½ *C.*	.5
19*lb.*	.16964
7 *ounc.*	.00390
	.67354

And here as you ſee meer Fractions reduced, ſo likewiſe may the Fractions of mixt numbers be reduced to Decimals; for example, theſe numbers 97

lb.

b. 7 ounces, 13 ¼ drams. *Item* of 67 Gallons 5 ¾ pints. *Item* 28 Quarters, 0 Buſhels and 2 ⅓ Pecks, after reduction are 97 .4891, 67 .7187, and 28 .0781.

97.4375
.0507
.0009
———
97.4891

67.625
.0937
———
67.7187

28.0625
.0156
———
28.0781

Again 22 ½ yards, 3 ¼ Nails; *Item* 36 Groſs, 3 Dozen and 5 particulars, being reduced, are 22 .7031, 36 .2847.

22.5.
.1875
.0156
———
22.7031

36 ,25
.0347
———
36 .2847

XVII. When a Decimal is propounded to know what Fraction it repreſents, ſearch the ſame Decimal in the ſecond Column of the Tablet, unto which it belongs, where if you find it expreſly, the number juſt againſt it in the firſt Column is the fraction you look for: So .65 (repreſenting the fraction of a Pound ſterling) being given, I find it in the ſecond Column of the Tablet of Money, and over againſt it in the firſt Column I find 13*s.* which is the fraction repreſented by .65, the decimal propounded. In like manner 3 .025 (repreſenting 3 ounces and .025 of an ounce *Troy*) being propounded, the number repreſented by it, is 3 Ounces, 0 *p. w.* 12 grains.

3. *Of Decimals to ſingle Fractions.*

XVIII. When in the ſecond Column of the

Tablet, unto which you are directed, you cannot preciſely find the decimal propounded, ſearch that which being leſs, comes neareſt unto it, and take the number that anſwers unto it in the firſt Column for the greateſt fraction of the number required: Then deducting the decimal ſo found out of the decimal given, find likewiſe the remainder as another decimal, and take his correſpondent number for the next fraction of the number required; and ſo proceed in that order, till you have diſcovered the intire number repreſented by the decimal propounded.

Example. .6739 being propounded, I demand the fraction of a pound ſterling repreſented by it; the decimal in the Tablet of money, which being leſs comes neareſt to .6739 is .65, whoſe correſpondent number in that Tablet is 13, which are the ſhillings of the number required; then ſubtracting (by the 1 Rule of the 25th Chapter of this Book) .65 out of .6739, the remainder is .0239, and the neareſt decimal in the ſame Tablet to .0239, is .0208, whoſe correſpondent number is 5, which are the pence of the number required. Laſt of all deducting .0208 out of 0239, the remainder is .0031, which gives you in the firſt Column 3, being the farthings of the number required: So that I conclude the intire fraction repreſented by the decimal .6739, is 13 *s*. 5 *d*. 3 *f*.

.6739*l*. *ſterling*.
Subtract 13 *s*. ——— .65
.0239
Subtract 5 *d*. ——— .0208
3 *f*. ——— .0031

In like manner 7.359*C*, being reduced by the Tablet of *Averdupois great weight* is $7\frac{1}{4}$*C*. 12*lb*. 4 ounc. And 94.58*lb*. reduced by the Tablet of *Averdupois little weight* is 94 *lb*. 9 ounces and 6 drams.

7.359 C.
Subtract 1 *quarter*——— .25
109
Subtract 12 *lb*——— .107
4 *oz*.——— .002
94.58 *lb*.
Subtract 9 *oz*. ——— .56
6 *Drams*.——— .02

CHAP. XXIV.

Addition of Decimal Fractions.

I. TO ſuch as well underſtand the *Notation* of *Decimal Fractions*, all the varieties of their *Numeration*, to wit, *Addition*, *Subtraction*, &c. will be as eaſie as the operations by whole numbers; therefore he that would be a good Proficient in *Decimal Arithmetick*, muſt throughly underſtand the 22 and 23 *Chapters* aforegoing.

II. When divers decimal fractions are given to be added together, they muſt firſt of all be orderly placed one under another according to the Doctrine of their Notation. So if theſe *Decimal Fractions*, to wit; .125, .39 and .7 were given to be added, they muſt be written down thus;

.125
.39
.7

or if you will have the ſame number of places to be in all the *decimals* given, without altering their values, they may be written thus,

.125
.390
.700

Not thus,

.125
.39
.7

For the Figures or Cyphers, which are of like degrees or places muſt be ſubſcribed directly one under another, *viz. tenth parts of* or *primes* muſt be written down directly underneath *tenths*; alſo *hundredth parts* or *ſeconds* muſt be placed under *hundredth parts*. as you ſee in the firſt Example, where .3 or three tenth parts in the ſecond decimal ſtands directly under .1 or one tenth part in the firſt decimal; likewiſe .7 or ſeven tenths in the third decimal ſtands directly under the tenths in the former, and ſo of the reſt.

In like manner, when mixt numbers, which conſiſt of Integers, and decimal parts are given to be added, due reſpect muſt be had of their ſubſcription one under another: ſo if theſe mixt numbers, to wit, 32 .056, 7 .07, and 1 .9 were given to be added, they muſt be written down thus,

32 .056
7 .07
1 .9

III. Having placed the decimals and drawn a line underneath in manner aforeſaid, add them together,

ether, beginning with the outermoſt rank towards he right hand (as hath been taught in Addition of vhole numbers of one denomination in the third hapter:) ſo if the decimals in the firſt Example f the ſecond *Rule* of this *Chapter* were given to be dded, I firſt ſubſcribe 5, which is all that ſtands in he firſt rank towards the right hand; then proeeding to the ſecond rank, I ſay 9 and makes 11, wherefore I write down 1, vhich is the exceſs of 11 above 10, and or the 10 I carry 1 in mind to the next ank, ſaying 1 in mind added to 7 makes 8 vhich added to 3 and 1 make 12, wherefor I write 2, which is the exceſs of 12 above 10, nder the line, reſerving 1 in mind for the 10, then prefix a point before 2, which ſtands in the firſt lace of decimals; and on the left hand of the point, o wit in the place of Units or firſt place of Integers, I write down 1 (being the 1 in mind) which lone, I find that the ſum of the Decimals given s 1 .215, that is, one Integer (whether it be a Perch, Yard, Foot, &c.) and $\frac{215}{1000}$ parts of an Integer, as you ee in the Example. In like manner theſe mixt numbers 32.056; 7.07 and 1 .9 being given to be added, their ſum will be found to be 41 .026, that is, 41 Integers and $\frac{26}{1000}$ parts of an Integer, as you ſee in the Margin; more *Examples* for the learners exerciſe are theſe.

.125
.39
.7
———
1.215

32.056
7.07
1 .9
———
41 .026

.65	24.7	503 . 75
.025	0.35	0 . 32
.03	5.27	0 . 12
.705	30 . 31	504 . 19

CHAP.

CHAP. XXV.

Subtraction of Decimal Fractions.

.837
.784
.053

I. HAving firſt written down the greater of the two numbers given (whether it be a whole number, mixt number, or decimal) and the leſſer underneath the greater, according to the directions in the ſecond Rule of the 24 Chapter, proceed as you are taught in Subtraction of whole numbers (by the Rules of the 4th Chapter:) So if this decimal fraction .784 were given to be ſubtracted from this decimal .837, the remainder will be .053, that is $\frac{53}{1000}$ parts of an Integer; in like manner if this mixt number 78 .919 were given to be ſubtracted from 295 .094, the remainder will be 216$\frac{175}{1000}$. In each of which examples you may obſerve that 10 is borrowed as often as need requires, according to the Rules of Subtraction of whole numbers of one denomination: Note alſo, when the decimals in both the numbers given conſiſt not of the ſame number of places, that decimal which is defective in places towards the right hand, muſt have the void places filled up with cyphers, or at leaſt cyphers muſt be ſuppoſed to be annexed: So if this decimal .04338 be given to be ſubtracted from this .65, the remainder will be found to be .60662, and the Work will ſtand as in the Margin, where you ſee the three void places are ſupplyed with cyphers, and then the operation is as in whole numbers, by borrowing 10 as often as the lower fi-

295 .094
78 .919
216 .175

.65000
.04338
.60662

gure cannot be ſubtracted from the upper. More Examples of ſubtraction of Decimals are theſe following.

24.04338	37	.394
.65	0.104	.35
23.39338	36.896	.044

CHAP. XXVI.

Multiplication of Decimal Fractions.

I. WHen two numbers are given to be multiplied, and are both mixt numbers, or both Decimal fractions, or one of them a whole number, and the other a decimal or mixt number (which are all the caſes that can happen) there is no neceſſity of writing them down preciſely one under the other as in Addition and Subtraction, for the product or number ſought in Multiplication depends not upon any regular placing of the two numbers given: So if this mixt number 56.3 were given to be added to this mixt number 1.30526, they ought to be written one under the other, as you ſee (according to the ſecond Rule of the 24th Chapter;) but if they are to be multiplied one by the other, they may be written thus,

1.30526
56.3

1 . 30526
56.3

II. In any of the Caſes which may happen in Multiplication of Decimals, multiply the numbers given as if they were whole numbers, then cut off always from the product by a point, comma, or line, ſo many places towards the right hand, as there are places of decimal parts in both the numbers given to be multiplied; that done, the figure or figures (if any happen to be) on the left hand of the ſaid point or line of ſeparation doth declare the Integer or Integers in the product, and thoſe on the right hand of the point are decimal parts of an Integer: So if this mixt number 56 .3 (that is, 56 Integers and $\frac{3}{10}$ of an Integer) be given to be multiplied by this next number 1 .30526, the product will be found 73 .486138, that is, 73 Integers and $\frac{486138}{1000000}$ parts of an Integer; for having choſen that to be the Multiplicator, which will cauſe leaſt work, and ſubſcribed it under the Multiplicand (to wit, 56 .3 underneath 1 .30526) I proceed according to the Rules of Multiplication of whole numbers, *viz.* having drawn a line underneath the numbers given, I multiply all the Multiplicand, to wit, 1. 30526, as if it were a whole number, by 3 the firſt multiplying figure, and ſubſcribe the product thereof, which is 391578 underneath the line, and proceeding in like manner with the other multiplying figures 6 and 5, at laſt I find the total of the particular products to be 73486138; and becauſe there are ſix places of decimal parts in both the numbers given (to wit, 5 places of parts in the multiplicand, and 1 place in the multiplicator) I cut off 6 places to the right hand from the total before produced, ſo will it ſtand thus 73|486138: Wherefore I conclude that the true product is 73 $\frac{487138}{1000000}$ or 73 .486138, that is, 73 Integers and almoſt $\frac{1}{2}$ of an Integer.

$$\begin{array}{r}
1\cdot30526 \\
56\cdot3 \\ \hline
391578 \\
783156 \\
652630 \\ \hline
73|486138
\end{array}$$

In like manner, if this mixt number 246.25 (that is 246 $\frac{25}{100}$) were given to be multiplied by 35 Integers, the true product will be found 8618. 75, that is 8618 Integers and $\frac{75}{100}$ parts of an Integer, as you ſee by the operation in the Margin, where you may obſerve that two places are cut off from the total number produced of the multiplication, towards the right hand, becauſe there are two places of decimals in the multiplicand (the multiplicator conſiſting of Integers only;) but if there had been decimal parts alſo in the multiplicator, ſo many more places ſhould have been cut off, as was ſhewed in the firſt *Example.*

```
 246.25
     35
 ------
 123125
 73875
 ------
8618.75
```

Again, If theſe two decimals .87 and .9 (to wit $\frac{87}{100}$ and $\frac{9}{10}$) were given to be multiplied one by the other, the true product will be found to be .783, that is $\frac{783}{1000}$ parts of an Integer as you ſee in the *Example*, where you may obſerve that the product is a fraction only; for after 3 places (being the number of places of decimals in both the numbers given to be multiplied) are cut off to the right hand, there remains no Integer on the left hand.

```
 .87
  .9
----
.783
```

III. When the multiplication is finiſht, if there ariſe not ſo many places in all as ought to be cut off by the ſecond Rule of this Chapter (which may often happen when the product is a fraction;) in ſuch caſe, as many places as are wanting, ſo many cyphers muſt be prefixed to the product on the left hand thereof, and then a point muſt be prefixt to ſign the product ſo increaſed for a decimal: So theſe decimals .0375 and .05 being given to be multiplied one by the other, I multiply 375 by 5, and there ariſeth 1875: Now according to the ſecond Rule of this Chapter, I ſhould cut off 6 places to the right hand, and here are but 4 in all; wherefore I prefix two Cyphers, to wit, as many as there are places wanting, and then prefixng a point, the true product will be .001875 or $\frac{1875}{1000000}$. In like manner if this mixt number 5.525 be multiplied by this decimal .0026, the true product will be found to be .0143650 (or $\frac{143650}{10000000}$) as you may ſee by the operation in the Margin, where one Cypher is prefixed to the Numbers ariſing from the total Multiplication to diſcover the true product.

.0375
.05
———
.001875
———
5.525
.0026
———
.33150
.11050
———
.0143650

To reduce Decimals to the known parts of the Integer.

IV. Decimal parts of an Integer may be reduced to the known or accuſtomed parts of ſuch Integer by Multiplication only, for it the decimal Fraction given be multiplied by that number which declareth how many known parts are equal to the Integer, the Product gives the number of known parts required: So this decimal Fraction of a Pound ſterling, to wit, .8687*l.* being propounded, I multiply it firſt by 20 (the number of Shillings contained in a Pound) and the product gives 17 Shillings and .3740 parts of a Shilling;

Shillings; which decimal .3740 being multiplied by 12 (the number of Pence in a Shilling) produceth 4 pence, and .488 parts of a Penny; Laſtly, multiplying .488 by 4 (the number of Farthings, which make a Penny) the product gives one Farthing, and .9520 parts of a Farthing, which are very near in value to another Farthing, ſo it appears that .8687 parts of a Pound ſterling are 17 *s.* 4 *d.* 2 *f.* very near. After the ſame manner, a *decimal fraction* of any Integer whatſoever may be reduced into the known or accuſtomed parts of ſuch Integer.

```
            .8687 l.
               20
        ----------
Shil.     17|3740
               12
        ----------
             7480
            3740
        ----------
Pence      4|4880
                4
        ----------
Farth.     1|9520
```

A briefer way to value any decimal part of a Pound of *Engliſh Money*, without loſs of a Farthing, may be this, *viz.* the figure (if any happen) in the firſt place of the decimal being doubled, gives Shillings; alſo if there be 5, or a figure greater than 5 in the ſecond place, one Shilling more is to be added to the former; laſtly, when 5 is taken from the figure in the ſecond place, if every Unit in the remainder be accounted as Ten, and the figure in the third place as Units, theſe Tens and Units taken as one number and leſſened by 1, give the number of Farthings, which with the Shillings before found declare the value of the decimal propounded; likewiſe if the figure in the ſecond place

A brief way to find the value of any decimal Fraction of a Pound of Engliſh Money.

(when any happens) be leſs than 5, every unit in ſuch figure is to be accounted ten as before: ſo in the decimal before-mentioned, to wit, .8687*l.* the figure 8 in the firſt place being doubled gives 16 ſhillings, alſo becauſe 5 is contained in 6 which ſtands in the ſecond place, one ſhilling more is to be added to the aforeſaid 16 ſhillings, which will now be made 17*s.* that done, the remainder of the ſaid 6 after 5 is ſubtracted, to wit, 1 being eſteem'd as 10, and added to 8 (which ſtands in the third place, and to be eſteemed as units) gives 18, from which abating 1, the remainder is 17 farthings or 4 pence and a farthing; ſo that the value of the ſaid decimal .8687 *l.* is found as before to be 17 ſhillings 4 pence 1 farthing. After the ſame manner this decimal of a pound of *Engliſh money*, to wit, .319 will be reduced to 6 ſhillings and 18 farthings, or 6 ſhillings 4 pence 2 farthings, which wants leſs than a farthing of the exact value of the decimal .319*l.*

V. Having explained all the caſes in *Multiplication* of *Decimals*, I ſhall here give the learner a taſte of their excellent uſe by ſome familiar queſtions whereby it will be evident, that what is oftentimes performed by many tedious *Multiplications* and *Diviſions* in the vulgar way, is effected for the moſt part by one or two *Multiplications* in *Decimals*.

See the queſtions from 49 *to* 73 *in the* 10*th Chapter of the Appendix.*

The firſt *Example* may be this: Suppoſe there is a certain piece of *Wainſcot*, in form of a *Rectangled Parallelogram*, commonly called a *long ſquare*, whoſe breadth is 3 *yards*, ¾ of a *yard*, 1 *nail* and ¼ of a *nail*; and the length of 6 *yards*, and ½ of a *yard*, the queſtion is to know how many *ſquare yards* are contained in that piece of *Wainſcot*; here becauſe it is deſired that the *ſuperficial content* may be given in *yards*, the parts of a *yard* as well in the breadth as in the length of the *Wainſcot* which are before expreſs'd by the accuſtomed parts of *quarters*, *nails*, &c. muſt be reduced into decimal parts of a *yard*, which are as eaſie to be found by a yard ſubdivided decimally, as the common parts of quarters and nails are found by a yard vulgarly ſubdivided: but for want of a yard ſubdivided decimally, this *Reduction* may be performed by the ſeventh *Tablet* of the precedent *Table* of *Reduction*, *viz.* looking into the ſaid *Tablet*, right againſt ¾ of a yard, I find this decimal ——— .75

Alſo the decimal correſpondent to 1 nail is ——— .0625

And the decimal of ¼ of a nail is ——— .015625

The ſum of thoſe three decimals is ——— .828125

Wherefore the breadth of the Wainſcot in yards and decimal parts is ——— 3.828125

Again the decimal of half a yard is .5, wherefore the length of the Wainſcot is ——— 6.5

The length and breadth being multiplied one by the other produce the *ſuperficial content*, therefore the number of ſquare yards required is ——— 24.8828125

Wherefore I conclude that 24 ſquare yards and ſomewhat more are contained in that piece of

Wainſcot; and it is evident by the Firſt place of the *decimal*, that what is above 24 yards is more than $\frac{8}{10}$ but leſs than $\frac{9}{10}$ of a ſquare yard; or more ſtrictly, it is more than $\frac{88}{100}$, but leſs than $\frac{89}{100}$ of a ſquare yard: but by taking all the places in the decimal you have the exact anſwer to this queſtion, becauſe the common parts of *quarters*, *nails*, and *quarters* of *nails*, may be always exactly reduced into decimals, but that ſeldom happens in other things; nevertheleſs, albeit by decimal operations you cannot always hit the mark, yet you may come as near it as is poſſibly to be imagined, and that with much more eaſe than by vulgar computations in queſtions of this nature, as will appear by comparing the precedent operation with the common way of working here in your view, *viz.* the 3 *yards*, 3 *quarters* of a *yard*, 1 *nail*, and $\frac{1}{4}$ of a *nail* (which expreſs the breadth before mentioned) muſt all be reduced into quarters of *nails* by the ſixth Rule of the ſeventh Chapter; ſo there will be found 245 quarters of Nails, as you ſee by the operation.

$$\begin{array}{l}
\textit{y. q. n. q.n.} \\
3\text{---}3\text{---}1\text{---}1. \\
\quad 4 \\
\hline
\quad 15 \\
\quad\;\, 4 \\
\hline
\quad 61 \\
\quad\;\, 4 \\
\hline
245 \textit{ quarters of nails.}
\end{array}$$

Again the 6 yards and half which expreſs the length aforeſaid, muſt likewiſe be reduced into quarters of Nails by the aforeſaid *Rule*; ſo there will be found 416 quarters of nails of a yard, as you ſee by the operation.

y. *q.*
6 —— ——2
4
——
26
4
——
104
4
——
416 *quarters of Nails.*

Then Multiplying the breadth and length one by the other, to wit, 245 by 416, the product will give 101920 for the ſuperficial content of the piece of Wainſcot in ſquare quarters of nails of a yard; now theſe ſquare quarters of nails of a yard muſt be reduced to ſquare yards, and the readieſt way to perform that, is to find firſt of all how many quarters of nails of a yard are contained in one yard in length, *viz.* ſince there are 16 nails in a yard, there are conſequently 4 times 16 quarters of nails, to wit, 64 quarters of nails in a yard in length; therefore 64 multiplied by 64 produceth 4096 ſquare quarters of nails in a yard ſquare; laſtly, I ſay by the Rule of three, if 4096 ſquare quarters of nails of a yard give 1 yard ſquare, how many yards ſquare will 101920 ſquare quarters of nails give? So will the anſwer be found 24 $\frac{3616}{4096}$ yards, which is the ſame with 24 .8828125 before found by the decimal operation (for $\frac{3616}{4096}$ is equal to the decimal .8828125, as will appear by reducing them to a common denominator by the fourteenth *Rule* of the ſeventeenth *Chapter.*) Now I leave it to the Reader to judge, which of theſe two ways is the more expeditious, and ſo let him take which liketh him beſt.

Example 2. There is a ſquared piece of Timber terminated at both ends with equal long ſquares, *viz.* the breadth of the piece of Timber is 1 foot 5 Inches 3 quarters of an Inch, and 1 half quarter of an Inch; the depth or thickneſs is 1 foot 3 inches 1 quarter of an Inch, and $\frac{1}{8}$ or half a quarter of an Inch; and the length of the piece is 11 feet 10 inches, and 3 quarters; the queſtion is how many ſolid or cubical feet are contained in that piece of Timber? The Anſwer may be found by *decimal Multiplication* in manner following, *viz.* Foraſmuch as it is deſired that the ſolid content may be given in feet, the parts of a foot as well in the breadth, depth, and length, which are before expreſs'd by the accuſtomed parts of inches, quarters, and half quarters, muſt be reduced into the decimal parts of a foot, which are as eaſie to be found by a foot ſubdivided decimally, as the other common parts by a foot vulgarly ſubdivided; but for want of a foot ſubdivided decimally, this *Reduction* may be performed by the eighth *Tablet* of the precedent *Table* of *Reduction*, viz.

The decimal correſpondent to 5 Inches is —————— .416

The decimal of $\frac{3}{4}$ of an Inch is —————— .062

The decimal of half a quarter of an Inch is —————— .01

The ſum of thoſe 3 decimals is —————— .488

Wherefore the breadth of the piece of Timber is —————— 1.488

In like manner the common parts of Inches, *&c.* in the depth or thickneſs of the piece of Timber, will be reduced by the ſaid *Tablet*, into theſe decimals, *viz.*

The decimal correſpondent to 3 inches is —.25
The decimal of ¼ of an Inch is ——————.02
The decimal of half a quarter of an Inch is—.01
The ſum of theſe 3 decimals is —— ——.28
Wherefore the depth or thickneſs is —— 1 .28

Again the accuſtomed parts of Inches, *&c.* in the length of the piece of Timber will be reduced to theſe decimals, *viz.*

The decimal of 10 Inches is——————.833
The decimal of ¾ of an Inch is ——————.062
The ſum of thoſe 2 decimals is——————.895
Wherefore the length of the piece is——11 .895

Now if the breadth, depth and length be multiplied continually, the laſt product is the ſolid content required, *viz.* 1 .488 multiplied by 1 .28 produceth 1 .90464, which multiplied by 11 .895 produceth 22 .65, *&c.* Wherefore I conclude that 22 ſolid feet, half a Foot, and ſomewhat more than half a quarter of a foot are contained in that piece of Timber.

Example 3. How many *Equinoctial Degrees* are correſpondent unto 136 *days*, 21 *hours*, and 40 *minutes*? The Anſwer is found by multiplying the time given by 360, for as 1 *day* is to 360 *degrees*, ſo 136 *days*, 21 *hours*, and 40 *minutes*, to the *Equinoctial degrees* required; but firſt the 21 *hours* and 40 *minutes* muſt be reduced to decimal parts of a day, by the tenth *Tablet*, thus.

The decimal of 21 hours is——————.875
The decimal of 40 minutes is——————.02777

The ſum of theſe 2 decimals is——————.90277
Therefore the time propoundedis---136.90277
Which being multiplied by 360 produceth——————————— } 49284.99,&

Wherefore I conclude, that 49284 .99 or very near 49285 *Equinoctial degrees* are correſpondent unto 136 *days*, 21 *hours*, and 40 *minutes*, which was required by the queſtion.

CHAP. XXVII.

Diviſion by Decimal Fractions.

I. IN any of the Caſes which may happen in Diviſion, if the Dividend be greater than the Diviſor, the quotient will be either a whole number or elſe a mixt number: But when the Dividend is leſs than the Diviſor, the quotient muſt neceſſarily be a fraction; for a leſſer number is contained in a greater once at the leaſt, but a greater is not contained once in a leſſer.

II. Sometimes the Dividend, whether it be a whole number, mixt number, or decimal fraction, is to be prepar'd by annexing a competent number of Cyphers thereunto, to make room for the Diviſor: So if 32,5 were given to be divided by 17.325 the Dividend 32 .5 muſt be increaſed with cyphers at pleaſure after this manner 32 .50000 &c. Likewiſe if 1 were given to be divided by 360, the Diiſion cannot be made till the Dividend 1 be increaed with cyphers, which being annexed, the Diviend will ſtand thus 1 .000000,*&c.* Here, note that he cyphers annexed in manner aforeſaid do ſupply laces of decimal parts, and will be uſeful in diſcoering the quality of the quotient according to the ourth *Rule* of this *Chapter.*

III. After the Dividend is prepared by annexng cyphers, when occaſion requires (as in the laſt ule,) all the places thereof muſt be eſteemed as ne whole number (to wit conſiſting of units or ntegers:) and ſo is the Diviſor to be eſteemed vhether it be a decimal fraction or mixt number; or in all caſes the Diviſion muſt be performed in very reſpect according to the Rules of Diviſion f whole numbers in the ſixth Chapter. So if this nixt number 326.25 were given to be divided by his mixt number 12.3, you muſt divide in the ſame manner, as when you divide 32625 Integers by 123 Integers. Alſo if this decimal .8356 were given to be divided by this decimal .05, you are to divide in the ſame manner, as when you divide 8356 Integers by 5 Integers; and after the quotient is found he degree or place of the firſt figure which ariſeth in the quotient muſt be enquired after; *viz.* you muſt know how far ſuch firſt figure is diſtant from the place of units, to the end that the point or line which is uſed to ſeparate between the place of units (or firſt place of Integers) and the firſt place of decimals may be duly placed: This is the only knot in decimal Diviſion, and may be reſolved by the following Rule, *viz.*

A general Rule to diſcover the quality of the quotient in all caſes of Diviſion by decimal Fractions.

IV. In any of the Caſes which may happen in Diviſion of decimals, the firſt figure which ariſeth in the Quotient, will be always of the ſame place or degree with that figure or cypher of the Dividend, which at the firſt queſtion ſtandeth over, or at leaſt belongeth unto the place of units in the Diviſor. To illuſtrate this Rule I ſhall give examples in all the principal caſes; and firſt let a mixt number be given to be divided by a mixt number, *viz.* Let it be required to divide 172 .5 by 3 .746 here (according to the ſecond Rule of this Chapter) the Dividend muſt be encreaſed with cyphers at pleaſure, ſo will it ſtand thus 172 .500000, &c. then Diviſion being made according to the Rules of Diviſion of whole Numbers in Chapter 6, the Quotient ariſing will be 46049, &c.

3 .746) 172 .500000 (46049, &c.

Now it remaineth to ſeparate the Integers in this quotient from the decimal parts; to perform which, I ſubſcribe the Diviſor 3 .746 orderly underneath,

3 .746) 172 .500000 (46, 049, &c.

3 .746.

the firſt Dividual 172 .50 (being that part of the Dividend whereof the firſt queſtion muſt be asked) or at leaſt I imagine the Diviſor to be ſo ſubſcribed, and ſo I find that the figure 3 which ſtands in the place of Units in the Diviſor will be placed under 7, which is the place of tens (or ſecond place of Integers) in the Dividend; wherefore by the fourth Rule before given, I conclude that the firſt figure ariſing in the quotient muſt likewiſe ſtand in the place of tens (or ſecond place of Integers) and conſequently the next place on the right hand muſt be the place of *Units*; ſo it is evident that the ſeparating point or line muſt be placed between the figure 6 and 0 in the quotient, that done, the true quotient is found to be 46 .049. &c. to wit, 46 Integers, and $\frac{49}{1000}$ parts of an Integer, and ſomewhat more: for 46 $\frac{49}{1000}$ is leſs than the true quotient, but 46 $\frac{50}{1000}$ is greater than it, and therefore albeit, after the aforeſaid Diviſion of 172 .500000 by 3.746 is ended, there will be a remainder, to wit, 446 which ſeems to be greater, yet here it is leſs in value than $\frac{1}{1000}$ part of an unit or Integer, and if to that remainder you annex another cypher and continue the diviſion, you will proceed nearer the truth and not miſs $\frac{1}{10000}$ part of an unit of the true quotient, and in that order you may proceed infinitely near, when you cannot obtain the quotient exactly by Diviſion of Decimals.

Example 2. Suppoſe this mixt number 2.34 be given to be divided by this mixt number 52 .125 (where you may obſerve that the Dividend is leſs than the Diviſor;) firſt (as before) annex cyphers at pleaſure to the Dividend, to make room for the Diviſor, then the diviſion being proſecuted as in whole numbers, at length theſe figures will ariſe in

52 .125) 2 .3400000 (0448, &c.
• • •
52, 125

the quotient, to wit, 448: and to the end the degree or quality of the firſt figure 4 may be diſcovered, I ſubſcribe the Diviſor 52 .125 under the firſt dividual 2 .34000 (for ſo far the firſt queſtion did extend in the Diviſion) and thereby I find that the figure 2 which ſtands in the place of units in the diviſor will be ſeated under 4, which is in the ſecond place of decimals, wherefore I conclude that the firſt figure ariſing in the quotient muſt alſo ſtand in the ſecond place of decimals, and conſequently the firſt place of decimals (which is next on the left hand to the ſecond) muſt be ſupplied with a cypher; ſo that if a cypher be prefixed on the left hand of 4, and then a point placed before that cypher, the quotient will at length be diſcovered to be .0448, &c. or $\frac{448}{10000}$, and ſomewhat more that is to ſay, $\frac{448}{10000}$ is leſs than the true quotient, but $\frac{449}{10000}$ is greater than it; and if you will proceed nearer the truth, you may continue the diviſion, as is directed in the firſt Example of this Rule.

Example 3. Where a whole number is divided by a decimal fraction, *viz.* ſuppoſe 82 Integers were given to be divided by this decimal .056; After cyphers are annexed to the dividend at pleaſure, and

.056) 82. 00000 (146428, &c.

the diviſion proſecuted as in whole numbers (to wit, 8200000 being divided by 56) theſe figures 146428 will ariſe in the quotient: now to the end the degree or ſeat of 1, the firſt figure in the quotient may be known, I ſubſcribe the Diviſor .056 under the firſt dividual 82 (for ſo far did the firſt queſtion in the diviſion extend;) and becauſe the diviſor is leſs than unity, I ſupply the place of units by a cypher or o prefixed on the left hand of the point of ſeparation in the diviſor; alſo I pre-

.056) 0082.00000 (1464.28, &c.

0.056

fix cyphers before (to wit on the left hand of) the Integers in the dividend to repreſent a ſucceſſion of places of Integers (for the order of places in Integers is from the right hand towards the left;) then I find that the cypher or o which repreſents the place of units in the diviſor, doth ſtand under that cypher, which repreſents the fourth place of Integers in the *dividend* (as you ſee by the Example;) wherefore I conclude that the firſt figure ariſing in the quotient muſt alſo be ſeated in the fourth place of Integers, and conſequently the 4 firſt places in the quotient will be Integers, and the reſt a decimal, ſo that the true quotient is 1464 Integers, and $\frac{28}{100}$ parts of an Integer, and ſomewhat more, *viz.* 1464 .28 is leſs than the true quotient, but 1464 .29 is greater than it.

Example 4. Suppoſe this decimal .0125 be given to be divided by this decimal .5; after diviſion is finiſhed according to the Rules of diviſion of

.5) .0125 (25

whole numbers (to wit after 125 is divided by 5) theſe figures 25 will ariſe in the quotient; now to diſcover the degree or ſeat of 2 the firſt figure in the quotient, I ſubſcribe the diviſor .5 under the firſt dividual .012, and having (as in the laſt Example) prefixed a cypher on the left hand of the point of ſeparation in the diviſor, to denote or repreſent the place of units, I find that ſuch cypher or place of units do ſtand under the figure 1, which is ſeated in the ſecond place of decimals in the dividend, wherefore I conclude by the Rule, that the firſt figure which ariſeth in the quotient muſt alſo be in the ſecond place of decimals, and therefore prefixing a cypher to ſupply the firſt place of decimals, and putting a point before that cypher, the quotient is at length diſcover'd to be .025 or $\frac{25}{1000}$.

.5).0125(.025
0.5

Example 5. Suppoſe this decimal .8564 be given to be divided by this .008, firſt I annex cyphers to the dividend at pleaſure, then proſecuting the diviſion as in whole numbers, to wit, dividing .856400 by 8, the quotient ariſing is 107.050, now to diſcover the degree or place of 1, the firſt figure in the quotient, I ſubſcribe the diviſor .008 under the firſt dividual .8, then I prefix a cypher to ſet forth, or ſupply the place of units in the diviſor, alſo I prefix cyphers to repreſent places of integers in the dividend; that done, I find that the cypher or 0 which ſupplieth

.008).856400(107.050

.008)000.85640(107.05
0008

plieth the place of units in the diviſor, doth ſtand under the Cypher which is ſeated in the third place of Integers in the dividend; wherefore I conclude by the Rule, that the firſt figure ariſing in the Quotient muſt be alſo in the third place of Integers, and conſequently the three firſt places in the Quotient will be Integers, and the reſt a decimal; ſo that the true Quotient is 107 .05 or $107\frac{5}{100}$.

Example. 6. Let it be required to divide this decimal fraction .73952 by this .32; firſt dividing 73952 by 32 as if they were whole numbers, the figures ariſing in the quotient will be 2311. Now to diſcover the quality or value of the ſaid figures I ſubſcribe the *Diviſor* .32 under the firſt *dividual* .73, then prefixing a Cypher as well on the left hand of the *dividend*, as of the *diviſor* ſo ſubſcribed (or imagined to be ſubſcribed) as aforeſaid, to repreſent the place of units in each of them, I find the cypher or 0, which ſupplieth the place of units in the *Diviſor*, to ſtand under the 0 which repreſents the place of units in the *dividend*; wherefore I conclude by the preceeding fourth *Rule*, that the firſt figure ariſing in the *quotient* will ſtand in the place of units, and conſequently the following places of the *quotient* will be a decimal fraction, ſo that the true *quotient* is 2 .311 or $2\frac{311}{1000}$.

.32)0 .73952 (2.311

0 .32

The reaſon of the foregoing fourth Rule will appear from the following Conſiderations.

I. If the product of the Multiplication of two numbers be divided by one of them, the quotient is the ſame with the other number: As, if 269 .0625, the product of 14 .35 multiplied by 18 .75, be divided by 14 .35; the quotient will give 18 .75.

II. If the Diviſor be multiplied by the firſt figure in the Quotient, the Product is the firſt number to be ſubtracted from the Dividend (being the ſame with the laſt particular Product in the Multiplication of the two Numbers that produced the Dividend;) and every particular place of that Product is of the ſame degree with that figure or cypher of the Dividend, which ſtands over ſuch particular place when the ſubtraction is made; For a figure of one degree (or place) cannot be ſubtracted from a figure of a different degree: As in the laſt mentioned Example, the work whereof is here in view; the Diviſor 14 .35 being taken as in a whole number and multiplied by 1, the firſt figure in the quotient, produceth 1435, which muſt be conceived to conſiſt of the ſame degrees as are in 269 .0 in the Dividend, from which the ſaid Product is to be ſubtracted, and therefore the ſaid Product 1435 is really but 143 .5, as you may ſee by the laſt particular Product, in the Multiplication of the mixt number 14 .35 by 18 .75.

$$\begin{array}{r|l}
14 & 35 \\
18 & 75 \\
\hline
 & 7175 \\
10 & 045 \\
114 & 80 \\
143 & 5 \\
\hline
14.35)269 & 0625.\ (18.75 \\
143 & 5 \\
\hline
125 & 56 \\
114 & 80 \\
\hline
10 & 762 \\
10 & 045 \\
\hline
 & 7175 \\
 & 7175 \\
\hline
 & 0
\end{array}$$

III. And therefore to diſcover the degree of the firſt figure in the Quotient, is nothing elſe but to find out the degree of that figure, which multiplying the figure or cypher in any particular place of the Diviſor, will produce the ſame degree as that figure or cypher in the Dividend is of, which ſtands over, or at leaſt belongs unto ſuch particular place of the Diviſor, at the firſt queſtion ; becauſe the degree produced muſt be ſubtracted from the like degree above it.

IV. Now among many Rules that might be given to diſcover the degree of the firſt figure in the quotient, and conſequently the degrees of all the reſt, the preceding fourth Rule of this Chapter is ſufficient, namely, The firſt figure which ariſeth in the quotient, is always of the ſame place or degree with that figure or cypher in the Dividend, which at the firſt queſtion ſtands over, or at leaſt belongs unto the place of units in the Diviſor: The reaſon is, becauſe if a figure ſtanding in the units place of the Diviſor, be multiplied by (or doth multiply) a figure of the ſame degree with that degree in the Dividend, which at the firſt queſtion belongs to the ſaid units place of the Diviſor, the firſt place in the Product ſhall be of that degree alſo, whether it be of Integers or decimal Parts; and conſequently the reſt of the places in the ſaid Product ſhall be of the ſame degrees with their correſpondent degrees (or places) in the Dividend, as they ought to be, to the end that due Subtraction may be made (according to Obſerv. 2.)

So in the Example before given, the firſt figure 1 in the quotient, ſhall be of the degree or place of Tens, becauſe if the figure 4 ſtanding in the units place of the Diviſor 14.35, be multiplied by Ten, to wit, the degree which the figure 6 in the Dividend is of that belongs to the ſaid 4 at the firſt queſtion, it will produce four Tens, to be ſubtracted from the ſaid ſix Tens: In like manner if a figure in the place of units be multiplied by units, the firſt place in the Product ſhall be units; if by tenth parts of an unit, or Integer, the firſt place in the Product ſhall be Tenths, *&c.*

Having explained all neceſſary Rules in Diviſion

concernin

concerning decimal fractions. I ſhall give a taſte of their excellent uſe, by the two following qœſtions and then conclude this Chapter.

Queſt. 1. A Merchant bought of Gold *Plate* 356 *ounces*, 13 *Penny-weight*, and 15 *grains* for 1160 *Pounds ſterling*, the queſtion is what he paid for an *ounce?* *Anſwer*, 3*l*—5*s*—½*d*. very near. The operation by *decimals* may be after this manner, *viz.*

By the ſecond *Tablet* of *Reduction* the *decimal* of 13 *Penny-weight* it— } .65

The *decimal* of 15 *grains* is ——.03125

The *Sum* of thoſe 2 *decimals* is——.68125

Wherefore the quantity of *Plate* in *ounces* and *decimal parts* of an *ounce* is ——————————— } 356.68125

Then by the *Rule* of Three, I ſay, if 356.68125 *ounces* coſt 1160 *pounds*, what 1 *ounce?* Here 'tis evident that if I divide 1160 by 356.68125, the *quotient* will give the value of an *ounce*, to wit, 3.252 *pounds*, or 3 *pounds*, 5 *ſhillings* and ½*d*. very near.

356.68125) 1160.0000000 (3.252, &c.

Queſt. 2. Suppoſe the length of the *Tropical Year* (or the ſpace of time wherein the Sun running through the whole *Ecliptick* Circle conſiſting of 360 *degrees*, is returned to the ſame *Equinoctial* or *Solſtitial* point from whence he departed) to conſiſt of 365 *days*, 5 *hours*, and 49 *minutes*, the queſtion is to know the Suns mean or equal motion for 1 *day*, to wit, what part of 360 *degrees* the Sun moveth in a whole day? The operation by *decimals*, thus.

By the tenth *Tablet* of *Reduction* the *decimal* correſpondent to 5 hours is——————————— .2083333

The *decimal* of 49 *minutes* is———— .0340277

The *ſum* of thoſe *decimals* is———— .2423610

Wherefore the *time* given in *days* and *decimal parts* of a *day* is 365 .2423610

Then by the rule of three, if 365 .242361 *days* give 360 *degrees* (or a total circumference) what will 1 *day* give? Here if I divide 360 by 365 .242361, the *quotient* will give the diurnal motion required; which will be found very near .98564, *&c.* or $\frac{98564}{100000}$ parts of a degree, which *decimal* being reduced into the common *Sexagenary parts* (by the fourth *Rule* of the 26 *Chapter*) will give $59'$——$8''$, *&c.* and ſuch is the Suns diurnal motion very near, according to the aforeſaid ſuppoſition of the length of the *Tropical Year*.

I ſhall here add the vulgar *Sexagenary* reſolution of this queſtion, that by comparing both ways together, the excellency of *decimal Arithmetick* in Calculations of this Nature may be the more perſpicuous.

The aforeſaid queſtion being ſtated according to the *Rule* of Three will ſtand thus,

days hours	*degrees*	*day.*
If 365 : 5 : 49——	360——	——1

The firſt term in the Rule muſt be reduced into *minutes* (by the ſixth *Rule* of the ſeventh *Chapter*;) ſo there will be found 525949 *minutes*.

D. *H.* ′
365——5——49
24

1465
730

8765
60

525949 *minutes.*

Likewiſe the third term 1 *day* muſt be reduced into *minutes*, which will be found to be 1440, as you ſee by the following operation.

1 *Day* or 24 *Hours.*
60

1440 *minutes.*

Then multiply the third term by the ſecond, to wit, 1440 by 360, the product is 518400, which being divided by the firſt term 525949 (according to the note in the ninth *Rule* of the 16th *Chapter*) the *quotient* will give $\frac{518400}{525949}$ parts of a degree, which fraction being reduced into the accuſtomed *Sexagenary* parts (by the ninth *Rule* of the ſeventeenth *Chapter*) will give as before $59' \; 8''$, &c. for the Suns mean diurnal motion; now which of theſe two ways is the more expeditious, I leave to him who is vers'd in both to determine.

CHAP. XXVIII.

The Rule of Three Direct in Fractions.

I. TO repeat ſuch things as have already been declared in reference to the definition of this Rule, as alſo to the due placing of the 3 given numbers, would be ſuperfluous; and if reſpect be had to the Rules of *Multiplication* and *Diviſion* in *Fractions* delivered in the 20, 21, 26 and 27 Chapters, the working of the Rule of three direct in fractions as well vulgar as decimal, is the ſame with that in whole numbers, *viz.* multiply the ſecond number by the third (or the third by the ſecond,) and divide the product by the firſt number, ſo the quotient is the fourth number ſought; to wit, the anſwer of the queſtion.

Otherwiſe thus in Vulgar Fractions.

Multiply the Denominator of the firſt number by the Numerator of the ſecond, alſo multiply that Product by the Numerator of the third number, and reſerve this laſt Product for a new Numerator; again multiply the Numerator of the firſt number by the Denominator of the ſecond, alſo multiply this Product by the Denominator of the third number, ſo ſhall this laſt product be a new Denominator; laſtly, the new fraction (whoſe Numerator and Denominator is found as aforeſaid) is the fourth number ſought, which, if it be a proper fraction, may (if occaſion require) be reduced into the known parts of the Integer (by the ninth Rule of the ſeventeenth Chapter;) if an improper fraction, it is to be reduced into its equivalent whole number or mixt number, by the thirteenth Rule of the ſeventeenth Chapter.

Example, If $\frac{3}{4}$ of a yard of Velvet be ſold for $\frac{2}{3}$ of a pound *ſterling*, what ſhall $\frac{5}{6}$ of a yard coſt? *Anſwer* $\frac{40}{54}$ *l.* or 14 *s.* 9 $\frac{7}{9}$ *d.* For according to the *Rule* I multiply the Denominator 4 by the Numerator 2, and the *product* is 8, this 8 again multiply by the Numerator 5, and the *product* gives 40 for a new Numerator: Moreover, I multiply the Numerator 3 by the Denominator 3, and the *product* which is 9 I again multiply by the Denominator 6, ſo the laſt *product* is 54 for a new Denominator; wherefore I conclude that $\frac{40}{54}$ is the fourth uumber ſought, which if it be reduced (according to the ninth Rule of the ſeventeenth Chapter) gives 14 *s.* 9$\frac{42}{54}$ *d.* (or 9 $\frac{7}{9}$ *d.*) for the *Anſwer* of the queſtion.

$$\begin{array}{ccccccc} y. & & l. & & y, & & l. \\ \frac{3}{4} & \text{——} & \frac{2}{3} & \text{——} & \frac{5}{6} & \text{——} & (\frac{40}{54} \end{array}$$

II. When any of the three given numbers is a whole number or mixt number, ſuch number muſt firſt of all be reduced into an improper fraction (by the tenth or eleventh *Rule* of the ſeventeenth *Chapter*) to the end that all the three given numbers may be 3 fractions: Moreover, If after ſuch Reduction, the firſt and third numbers be not fractions of Integers of the ſame particular denomination, ſuch of the ſaid numbers which is of the leſſer denomination, muſt be reduced to a fraction of the greater (by the ſixteenth *Rule* of the ſeventeenth *Chapter*;) which preparations being performed, the reſt of the Work is to be proſecuted according to the firſt *Rule* of this *Chapter.* An *Example* of this ſecond *Rule* here followeth. If a quantity of *Ambergreece* weighing $\frac{5}{7}$ *lb. Troy*, be ſold for 60 *l. ſterling*, what are 19 $\frac{5}{8}$ *grains* worth at that rate? *Anſwers*, $\frac{65940}{552960}$ *l.* or 2 *s.* 4 $\frac{119}{192}$ *d.*

This queſtion being ſtated according to the 7 *Rule* of the 8 *Chapter* will ſtand thus,——

lb. *l.* *gr.*
1 $\frac{5}{7}$ —— 60 —— 19 $\frac{5}{8}$

which 3 numbers will be reduced (by the tenth and eleventh *Rules* of the ſeventeenth *Chapter*) into theſe improper *fractions.*——————

lb. *l.* *gr.*
$\frac{12}{7}$ —— $\frac{60}{1}$ —— $\frac{157}{8}$

But ſince the third number $\frac{157}{8}$ *grains Troy* is not a fraction of an Integer of the ſame name with the firſt (which is a fraction of a pound *Troy*,) it muſt be reduced into a fraction of a pound *Troy*, thus, $\frac{157}{8}$ *gr.* is $\frac{157}{8}$ of $\frac{1}{24}$ of $\frac{1}{20}$ of $\frac{1}{12}$ of a pound *Troy*, which compound fraction will be reduced (by the 16 *Rule* of the 17 *Chapter*) into this ſingle fraction, to wit, $\frac{157}{46080}$ *lb. Troy* and ſo the 3 numbers will at length ſtand thus in the *Rule.*

$\frac{12}{7}$ *lb.* —— $\frac{60}{1}$ *l.* —— $\frac{157}{46080}$ *lb.*

Then working as in the firſt *Example* of this *Chapter*,) the anſwer will be found $\frac{65940}{552969}$ *l.* which being reduced (according to the 3 and 4 *Rules* of the 17 *Chapter*) is found equal unto 2 *s.* 4 $\frac{119}{192}$ *d.*

Another *Example.* When the $\frac{2}{5}$ of $\frac{3}{4}$ a *Ship* is valued at 147 *l.*—11 *s.*—3 *d.* how much is the whole Ship worth? *Anſw.* 491 *l.*—17 *s.*—6 *d.*

Note, when in any queſtion whatſoever a compound fraction, to wit, a fraction of a fraction, is one of the given numbers, ſuch compound fraction muſt firſt of all be reduced to a ſingle fraction (by the 16 *Rule* of the 17 *Chapter*;(ſo here the *compound fraction* $\frac{2}{5}$ of $\frac{3}{4}$ being reduced into a ſingle fraction gives $\frac{6}{20}$ or $\frac{3}{10}$; then ſay if $\frac{3}{10}$ be worth 147*l.* 11*s.* 3*d.* what is 1 or the whole *Ship* worth?

Ship *l.* *s.* *d.* *Ship*
$\frac{3}{10}$——147: -11:——3 ———1

After due reduction is made by converting the 147*l* 11*s.* 3*d.* into pence, and that number of pence, as alſo the third number 1. into improper fractions, the 3 numbers will ſtand in the Rule thus,

Ship *pence* *Ship*
$\frac{3}{10}$————$\frac{35415}{1}$————$\frac{1}{1}$

Laſtly, Proceeding as in the firſt Rule of this Chapter, the fourth number will be found to be $\frac{354150}{3}$*d.* which being reduced firſt by the 13 *Rule* of the 17 *Chapter*, and then by the 7 *Rule* of the 7 *Chapter*, the *Anſwer* at length is 491*l.*——17*s.*——6*d.*

An Example of the Rule of three direct in *Decimals* may be this that follows. If 19 *ounces*, 3 *penny weight*, and 5 *grains* of Gold, be worth 62*l.*—10*s.*—6*d.* what is the value of 1 $\frac{1}{2}$ *ounce*? *Anſw.* 4*l.*—17*s.*——10 $\frac{3}{4}$*d.* very near.

By the 2. *Tablet* in the *Table* of *Reduction* in the 23 Chapter, the *decimal fraction* correſpondent to 3 *penny weight*, is——————— .15

Alſo, the *decimal* of 5 *grains* is-——.010416

The ſum of thoſe 2 *decimals* is——.160416

Wherefore the firſt number in the Rule of three is———— *oz.* 19.160416

Again, by the firſt *Tablet* of the aforementioned *Table* the *decimal* of 10 *ſhillings* is———— .5

Alſo the *Decimal* of 6 *pence* is——025

The ſum of theſe two *decimals* is——525

Wherefore the ſecond number in the rule of three is ———— *l.* 62 .525

Moreover by the ſaid *Tablet* 2. the *decimal* of ½ of an *ounce* or 10 *penny weight* is 5, wherefore the third number in the Rule of three is——— *oz.* 1 .5

So that after the ſaid Reduction is finiſht the 3 given numbers will ſtand in the Rule thus:

oun.	*l.*	*oun.*
19.160416——	---62.525——	-1 .5

Laſtly, multiplying the ſecond number by the third, and dividing the product by the firſt number, (according to the Rules of Multiplication and Diviſion of Decimals delivered in the 26 and 27 Chapters) the fourth number will be this, to wit, 4.8948, &c. that is four pound *ſterling*, and $\frac{8948}{10000}$ parts of a pound, which *decimal* being reduced according to the fourth *Rule* of the 26 *Chapter*) gives 17*s.*——10*d.*——3*far.*

The proof of the Rule of three direct in Fractions is the ſame as in whole numbers, reſpect being had to the Rules of Multiplication in Fractions.

CHAP. XXIX.

The Inverſe Rule of Three in Fractions.

1. After a queſtion belonging to this Rule is duly ſtated(according to the ſeventh rule of the eighth Chapter) and prepared if need require, according to the ſecond Rule of the 28 Chapter; the operation will be the ſame as in the Rule of three Inverſe in whole numbers, reſpect being had to the Rules of Multiplication and Diviſion in Fractions, *viz.* multiply the firſt number by the ſecond, and divide the Product by the third; the quotient is the fourth number ſought, to wit, the anſwer of the queſtion.

Or thus, in Vulgar Fractions.

Multiply the Denominator of the third fraction by the Numerator of the ſecond, alſo multiply that Product by the numerator of the firſt fraction, and reſerve the laſt product for a new Numerator: again multiply the Numerator of the third fraction by the Denominator of the ſecond; alſo multiply this Product by the Denominator of the firſt fraction, ſo is the laſt Product a new Denominator; laſtly, this new fraction is the fourth number ſought, or anſwer of the queſtion.

Examples

Example, if of cloth, which is, 1 $\frac{3}{4}$ yard in breadth 3 $\frac{1}{2}$ yards in length will make a Cloak, how much in length of ſtuff which is $\frac{5}{8}$ yards in breadth will make a Cloak of the ſame bigneſs with the former? *Anſwer* 9$\frac{4}{5}$ yards.

The 3 numbers being duly placed will ſtand thus——— } *brea.* *leng.* *brea.*
1 $\frac{3}{4}$*y.*—3$\frac{1}{2}$*y.*—$\frac{5}{8}$*y.*

Then (after the firſt and ſecond numbers are reduced into improper fractions) the three Numbers will ſtand thus——————— } $\frac{7}{4}$——$\frac{7}{2}$——$\frac{5}{8}$

Laſtly, 8, 7 and 7 being multiplied continually give 392 for a numerator; alſo 5, 2 and 4 being multiplied continually give 40 for a denominator, whereby this improper fraction $\frac{392}{40}$ ariſeth, which (by the thirteenth rule of the ſeventeenth Chapter) will be found to be 9 $\frac{32}{40}$, or (the fraction being reduced into its leaſt terms) 9 $\frac{4}{5}$, which is the *Anſwer* of the queſtion.

Ex. 2. Suppoſe when Wheat is at 2*l.*--00*s.*--6*d.* the Quarter, the penny white loaf ought to weigh 8 ounces and 1 $\frac{5}{29}$ *penny weight* of *Troy* weight; what ought it to weigh when Wheat is at 36 ſhillings the Quarter? *Anſwer*, 9 ounces and 1 $\frac{37}{116}$ *peny weight.*

The 3 given numbers being duly placed in the rule and reduced will ſtand thus. } *pence* *p. w.* *pence*
$\frac{486}{1}$: $\frac{4674}{29}$: $\frac{432}{1}$

And if the operation be proſecuted according to the rule before given, the *Anſwer* will be found 181 $\frac{3996}{12528}$ penny weight, or 9 ounces, 1 $\frac{37}{116}$ penny weight.

CHAP.

CHAP. XXX.

The double Rule of Three in Fractions.

I. THe *Double Rule of Three* is ſo called, becauſe it is compoſed of two ſingle Rules, and may either be reſolved at one Work by the Rule compound of 5 numbers, or elſe by two diſtinct ſingle Rules of three; which latter way, to ſuch as underſtand the Rule of three in fractions, is (as I conceive) leſs troubleſome in the ſtating, and (in the method whereby I intend to proſecute it) the ſame in operation with the former. This I ſhall manifeſt firſt in whole numbers, then in fractions.

Example 1. If I pay 28 ſhillings for the carriage of 3 C. weight for 50 miles, how much ought I to pay for the carriage of 17 C. for 84 miles? *Anſwer,* 13*l.*——6*s.*——6*d.* $\frac{18}{25}$.

Of the 5 given numbers I make choice of three ſuch which will make a ſingle rule of three, and ſay,

C. *ſhil.* *C.*
If 3——28——17

Which Rule I find (by the third rule of the ninth Chapter) to be direct, and therefore I multiply the third number 17 by the ſecond 28, and the product which is 476 I place as a numerator over the diviſor as denominator. Then with this fraction (whether it happen to be a proper or improper fraction) and the remaining two numbers in the queſtion, which have not yet been uſed, I orm a ſecond rule of Three, and ſay, $\frac{476}{3}$

miles *ſhill.* *miles.*
If $\frac{50}{1}$ ——— $\frac{476}{3}$ ——— $\frac{84}{1}$

Which being a Rule of Three direct, I work as a rule of three in fractions, according to the firſt rule of the 28 Chapter, and ſo find the fourth number to be $\frac{39984}{150}$*s.* or 13*l.* —6*s.*—6$\frac{1}{2}\frac{8}{5}$*d.*

Or the firſt ſingle Rule being varied, the operation will be thus,

miles *C.* *miles* *C.*
1. *By a Rule Inverſe,* 50 —3— 84 —— ($\frac{150}{84}$

C. *ſh.* *C.* *ſh.*
2. *By a Rule Direct,* $\frac{150}{84}$: $\frac{28}{1}$: $\frac{17}{1}$: ($\frac{39984}{150}$

Otherwiſe thus,

C. *m.* *C.* *m.*
1. *By a Rule inverſe,* 3 —— 50 —— 17 — ($\frac{150}{17}$

m. *ſh.* *m.* *ſh.*
2. *By a Rule direct,* $\frac{150}{17}$: $\frac{28}{1}$: $\frac{84}{1}$: ($\frac{39984}{150}$

Thus you ſee the two ſingle rules to be varied three manner of ways in reſolving the queſtion propounded, and each way produceth the ſame *Anſwer*; the like diverſity may be found in all queſtions reſolvable by the double rule of three, or rule compound of 5 numbers.

Example 2. If 40$\frac{3}{5}$*l.* in $\frac{2}{3}$ of a year, gain 2$\frac{1}{2}$*l.* what will 100 *l.* gain after that rate in $\frac{7}{2}$ of a year? *Anſw.* $\frac{2500}{774}$*l.* or 5 *l.*——7*s.*——9 $\frac{3}{29}$*d.*

By 2 Single Rules of Three, thus,

l. l. l. l.
1. *By a Rule Direct,* $\frac{203}{5} : \frac{5}{2} :: \frac{100}{1} : (\frac{2500}{406}$.

year l. year l.
2. *By a Rule Direct,* $\frac{2}{3} : \frac{2500}{406} :: \frac{7}{12} : (\frac{52500}{9744}$.

Or by theſe two ſingle Rules.

year l. year l.
1. *By a Rule Direct,* $\frac{2}{3} : \frac{5}{2} :: \frac{7}{12} : (\frac{105}{48}$

l. l. l. l.
2. *By a Rule Direct,* $\frac{203}{5} : \frac{105}{48} :: \frac{100}{1} : (\frac{52500}{9744}$

Otherwiſe thus,

l. year l. l.
1. *By a Rule Inverſe,* $\frac{203}{5} : \frac{2}{5} :: \frac{120}{1} : (\frac{406}{1500}$

l. year l. l.
2. *By a Rule Direct,* $\frac{406}{1500} : \frac{5}{2} :: \frac{7}{12} : \frac{52500}{9744}$

Thus by 2 ſingle Rules of Three, varied three ſeveral ways, you ſee the *Anſwer* of the queſtion to be $\frac{52500}{9744}$ *l.* to wit, 5*l.*—7*s.*—9 $\frac{3}{20}$*d.*

CHAP. XXXI.

The Rule of Falſe in Fractions.

I. WHen a queſtion propounded cannot readily be applied to the *Rule of Three*, or any of the vulgar Rules in *Arithmetick*; the beſt refuge for ſuch as are not acquainted with *Algebra* is the Rule of *two falſe Poſitions*, which, for that it hath already been handled in *whole Numbers*, I ſhall the more briefly touch upon in *Fractions.*

II. When a number is ſought by a queſtion, you are to feign or ſuppoſe ſome number taken by gueſs to be the number ſought, and to make trial whether that feigned number will anſwer the conditions in the queſtion or not, by comparing the number reſulting at the end of the Work, with the given number reſulting from the true number ſought; and if you find both thoſe reſults to be the ſame, then is the number which you firſt took by gueſs, the true number or anſwer of the queſtion; but if the number reſulting from the ſuppoſitious number be either greater or leſs than the given reſult, with which it ought to be compared (to ſee whether you have hit the mark or not) ſuch exceſs or defect muſt be noted for the Error of the firſt Poſition, to wit, an Exceſs muſt be ſignified by this note †; and a Defect by this —

III. In like manner a ſecond number muſt be feigned, and after trial is made therewith, to ſee whether it will perform the conditions preſcribed in the queſtion, by comparing the reſults as aforeſaid,

ſaid, the error of this ſecond poſition, if two much, is to be noted by +, if too little by —, as before.

IV. After the errors of both poſitions are diſcovered, the two numbers before ſuppoſed or feigned to be the number ſought, muſt be multiplied by the altern errors, that is, the firſt Poſition by the ſecond Error, and the ſecond Poſition by the firſt Error; then if the notes of the Errors be unlike, to wit, one of them +, and the other —, the ſum of the ſaid Products is to be taken for a dividend, and the ſum of the Errors for a Diviſor: but if the notes of the Errors be both alike, to wit, both of them +, or both —, the difference of the ſaid Products is to be taken for a Dividend, and the difference of the Errors for a Diviſor; laſtly, the Quotient ariſing from the Diviſion (made by the ſaid Dividend and Diviſor) gives the true number ſought, or anſwer of the queſtion, if it be ſolvable by the *Rule of Falſe.* Theſe Rules, are the ſame in ſubſtance with thoſe delivered in the 15 Chapter, and may be farther illuſtrated by the following Queſtions.

Queſt. 1. A Gentleman hired a Servant for a Year for 6 pounds *ſterling*, and a livery Cloak valued at a certain rate, but it happened that $\frac{7}{12}$ of the Year being expired they fell at variance, and the Gentleman put away his Servant, giving him the Cloak together with 50 Shillings in Money, which was the Servants full due for the time of his Service, the queſtion is to find what the Cloak was valued at? *Anſw.* 2 *l.*——8 *s.*——0 *d.*

1. I ſuppoſe the Cloak to be valued at 3 pounds, and then ſeek how much thereof was due to the

y. l. y.
1.——3——$\frac{7}{12}$——($\frac{7}{4}$l.

Servant, ſaying, if one Year give 3 *l.* how much $\frac{7}{12}$ of a Year? *Anſw.* $\frac{7}{4}$ *l.*

y. l. y.
1——6——$\frac{7}{12}$——($\frac{7}{2}$ l.

2. I likewiſe find what parts of the 6 Pounds was due to the Servant at the end of $\frac{7}{12}$ of the Year, ſaying, if 1 Year give 6 Pounds, how much $\frac{7}{12}$ of the Year? *Anſwer*, $\frac{7}{2}$ *l.*

3. For as much as the Cloak together with the Money which the Servant received, ought to be equal to the part of the Cloak, together with the part of the 6 Pounds wages due to him at the end of $\frac{7}{12}$ of the Year, therefore 3 *l.* (the ſuppoſed value of the Cloak) together with 2 $\frac{1}{2}$ *l.* (the Money which the Servant received) ſhould be equal to $\frac{7}{4}$ of a Pound (the value of part of the Cloak due to the Servant at the end of $\frac{7}{12}$ of the Year) together with $\frac{7}{2}$ *l.* (the wages due for the ſame time) that is to ſay, $\frac{11}{2}$ *l.* (the ſum of 3 *l.* and 2 $\frac{1}{2}$) ſhould be equal to $\frac{21}{4}$ *l.* (the ſum of $\frac{7}{4}$ *l.*) and $\frac{7}{2}$ *l.* but it is greater by $\frac{1}{4}$, wherefore the firſt Poſition for the value of the Cloak being 3 Pounds, the Error is found to be $\frac{1}{4}$ too much.

4. I make a ſecond Suppoſition gueſſing the value of the Cloak to be 2 Pounds, and Proceeding in every reſpect as with the firſt Suppoſition I find the *Errour* to be $\frac{1}{6}$ too little, ſo that the two *Poſitions* with their *Errours* will be as you ſee:

Pos.		*Er.*
3	+	$\frac{1}{4}$.
2.	—	$\frac{1}{6}$.

Now in regard the *errours* are *fractions*, I may take in their ſtead whole numbers in the ſame proportion, to wit, multiplying the *Numerator* of the firſt fraction (or firſt *errour*) by the *Denominator* of the ſecond, I take the Product which is 6 inſtead of the firſt *errour* $\frac{1}{4}$, likewiſe multiplying the *Numerator* of the ſecond fraction by the *Denominator* of the firſt, I take the Product which is 4 inſtead of the ſecond *errour* $\frac{1}{6}$, Or inſtead of the ſaid 6 and 4 I may take 3 and 2 which are in the ſame proportion with 6 and 4, or with $\frac{1}{4}$ and $\frac{1}{6}$.) Then multiplying the *poſitions* and new *errours* croſs-wiſe, and adding the Products together (becauſe the ſigns are unlike) the ſum is 12 for a *Dividend*, and the ſum of the *errours* 3 and 2 is 5 for a *Diviſor*, ſo the Quotient will be found to be $2\frac{2}{5}$ *l.* ſo much therefore was the value of the Cloak, as will eaſily appear if trial be made with $2\frac{2}{5}$ *l.* in the ſame manner as with the firſt feigned number.

Pos.		*Er.*		
3	+	$\frac{1}{4}$	6	3
2	—	$\frac{1}{6}$	4	2
6				
6				

5)12($2\frac{2}{5}$ *poun.*

Queſt. 2. *Vitruvius* (*in lib.* 9. *cap.* 3.) reporteth that King *Hiero* having given commandment for the making of a *Crown* of pure *Gold*, was informed that the Workman had detained part of the *Gold*, and mixt the reſt with as much *Silver* as he had ſtole of *Gold*; the King being much diſpleaſed at the deceit, recommended the examination of the Buſineſs to the famous *Archimedes* of *Syracuſa*, who without defacing the *Crown* diſcovered the Cheat in this manner; *viz.* Experience telling him that a quantity of *Gold* would poſſeſs leſs room or ſpace than the ſame quantity of *Sil-*

ver, and conſequently that a mixt maſs of *Gold* and *Silver* of the ſame quantity would take up ſome mean ſpace between the two former, he made a maſs of pure *Gold* of the ſame weight with the *Crown*, likewiſe another maſs of *Silver* of the ſame weight, then having put the *Crown*, as alſo the other two Maſſes ſeverally into a Veſſel filled up to the brim with Water, he diligently reſerved the Water flowing over into another Veſſel, and from thoſe 3 ſeveral quantities of Water ſo expell'd, he found out the quantity of *Gold* and of *Silver* in the *Crown*: But foraſmuch as *Vitruvius* delivers not the practical operation, I ſhall here ſhew the ſame after the manner of *Cardanus*, *Gemma Friſius*, and other *Arithmeticians*.

Let us therefore ſuppoſe the weight of the *Crown*, as alſo of the two ſeveral Maſſes to have been 5 *l.* Suppoſe alſo that by putting of the Maſs of *Gold* into the Veſſel, 3 *l.* of Water was expell'd; by putting in of the *Crown*, $3\frac{1}{4}$*l.* and by putting in of the Maſs of *Silver*, $4\frac{1}{2}$*l.* The queſtion therefore is to know how much *Gold* and how much *Silver* the *Crown* was compoſed of. This may be reſolved after this manner. Suppoſe 3 *l.* of *Gold* to be in the *Crown*, then there remained 2 *l.* of *Silver*, now ſay by the Rule of 3, if 5 *l.* of *Gold* expel 3 *l.* of Water, how much 3 *l.* of *Gold*? *Anſwer*, $1\frac{4}{5}$ *l.* Alſo if 5 *l.* of *Silver* expel $4\frac{1}{2}$ *l.* of Water, how much 2 *l.* of *Silver*? *Anſwer*, $1\frac{4}{5}$ *l.* of Water, add therefore the Water of the *Silver* and of the *Gold* together, to wit, $1\frac{4}{5}$ and $1\frac{4}{5}$, ſo there will ariſe $3\frac{3}{5}$ *l.* of Water: This ought to have been $3\frac{1}{4}$*l.* (for ſo much overflowed

5—3—3—($1\frac{4}{5}$
5—$4\frac{1}{2}$-2—($1\frac{4}{5}$

flowed by putting in of the *Crown*;) but it is too much by $\frac{7}{20}$, wherefore $\frac{7}{20}$ is to be noted with ✝ for the *errour* of the *firſt Poſition* 3*l.* Again, feign another quantity of *Gold* to have been in the *Crown*, to wit, 2*l.* therefore there remained 3*l.* of *Silver*; then ſay if 5*l.* of *Gold* expel 3*l.* of Water, how much 2*l.* of *Gold?* *Anſw.* 1 $\frac{1}{5}$ *l.* of Water: Alſo if 5 *l.* of *Silver* expel 4 $\frac{1}{2}$ *l.* of Water, how much 3 *l.* of *Silver?* *Anſw.* 2 $\frac{7}{10}$, then add 1 $\frac{1}{5}$ *l.* unto 2 $\frac{7}{10}$, the ſum will be 3 $\frac{9}{10}$ *l.* of Water: this ought to have been 3 $\frac{3}{4}$ *l.* but it is too much by $\frac{13}{20}$. wherefore $\frac{13}{20}$ is to be noted with ✝ for the *errour* of the *ſecond Poſition* 2 *l.* Here becauſe the *errours* are *fractions* having a *common Denominator*, I take their *Numerators* 7 and 13 inſtead of the *errours*; then multiplying croſs-wiſe, to wit, 3 by 13 the Product is 39, alſo 2 by 7 the Product is 14, which ſubtracted from the former Product 39 (becauſe the *errours* are like) leaves 25 for a *Dividend*; alſo the difference between the *errours* 7 and 13 is 6 for a *Diviſor*; Laſtly, dividing 25 by 6, the Quotient is 4 $\frac{1}{6}$; ſo much *Gold* therefore was in the *Crown*, and conſequently (becauſe the weight of the *Crown* was 5 *l.*) there was $\frac{5}{6}$ *l.* of *Silver* which may be proved thus: Say, if 5*l.* of *Gold*, expel 3*l.* of Water, how much 4 $\frac{1}{6}$ *l.* of *Gold?* *Anſwer*, 2 $\frac{1}{2}$ *l.* of Water: Again, if 5*l.* of *Silver* ex-

5——3——2——(1 $\frac{1}{5}$

5——4 $\frac{1}{2}$——3——(2 $\frac{7}{10}$

Pos.		*Er.*	
3	✝	$\frac{7}{20}$	7
2	✝	$\frac{13}{20}$	13
39			
14			

6) 25 (4 $\frac{1}{6}$ *lb.* of *Gold.*

ple $4\frac{1}{2}$ of Water, how much $\frac{5}{6}$ of *Silver?* *Anſw.* $\frac{3}{4}$ *l.* of Water, which being added to $2\frac{1}{2}$ *l.* the ſum is $3\frac{1}{4}$ *l.* of Water, to wit, as much as flowed over when the *Crown* was put into the Veſſel.

Here note, that in making a trial of this nature, there is no neceſſity that the Maſs of Gold or of Silver be of the ſame weight with the *Crown*, or whatſoever thing is to be examined, but of what notable part of weight you pleaſe.

Note alſo; that for the more eaſie diſcovering of the Dividend and Diviſor by the notes of + and —— according to the fourth Rule of this Chapter, the following Verſe may be a help, to wit,

Addito diſſimiles, ſubtrahitoque pares.

Or thus,

Notes being unlike, Addition make;
If like, leſſer from greater take.

The Reader may ſee more queſtions to exerciſe the *Rule* of *Falſe* in the tenth Chapter of the *Appendix*, and the demonſtration thereof in the ninth Chapter of the ſame:

CHAP. XXXII.

The Extraction of the Square (or Quadrate) Root.

1. THE Extraction of the Square-root is that, by which having a number given, we find out another number, which being multiplied by it ſelf, produceth the number given.



II. In the Extraction of the Square-root, the number propounded is always conceived to be a ſquare number, that is, a certain number of little ſquares comprehended within one intire great ſquare, and the root or number required is the ſide of that great ſquare, as will readily appear by this Diagram, where you ſee 25 little ſquares contained within one great ſquare; now if the ſaid content 25 be given, and the ſide or root of the ſquare containing the ſaid 25 little ſquares is required, the invention of ſuch ſide or root is called the Extraction of the ſquare-root; which root muſt be ſuch, that if it be ſquared, that is, multiplied by it ſelf, the Product muſt be equal to the ſquare content firſt given: So 5 is the ſqure root of 25, for 5 times 5 is 25. Likewiſe this ſquare number 49 being propounded, his root is 7.

III. Square numbers are either ſingle or compound.

A ſingle ſquare number.

IV. A ſingle ſquare nnmber is that, which being produced by the multiplication of one ſingle figure by it ſelf, is always leſs than 100: So 25 is a ſingle ſquare number produced by 5; likewiſe 4 is a ſquare number produced by 2.

V. All the ſingle ſquare numbers together with their reſpective roots are expreſſed in the Table following.

Squares.	1	4	9	16	25	36	49	64	81
Roots.	1	2	3	4	5	6	7	8	9

Here in the uppermoſt rank of the Table are placed in the ſingle ſquare numbers of every particular figure, and in the other their reſpective roots; and therefore if it were demanded, What is the ſquare-root of 36, the anſwer will be 6. So the ſquare-root of 16 is 4; the ſquare-root of 9 is 3, &c. And contrarily the ſquare of the root 6 is 36: Alſo the ſquare of 3 is 9.

VI. When a ſquare number is given, that exceeds not 100, and yet is none of the ſquare numbers mentioned in the Table, for his root you are to take the root of the ſquare number that being leſs, yet comes neareſt unto it: So 45 being given, the root that belongs unto it is 6, and 10 being given, his correſpondent root is 3.

VII. A compound ſquare number is that, which being produced by a number (that conſiſts of more places than one) multiplied by it ſelf, is never leſs than 100: So 1024 is a compound ſquare number produced by the multiplication of 32 multiplied by it ſelf.

A compound ſquare number.

VIII. To prepare any ſquare number given for extraction, put a point over the firſt place thereof on the right hand (being the place of Units;) then proceeding towards the left hand, paſs over the ſecond place, and put another point over the third place, alſo paſſing over the fourth place put another point over the fifth, and ſo forward in ſuch manner that between every two points which are next one to the other, one place will be intermitted: So if the ſquare root of 1024 be required, the firſt point is to be placed over 4, and the ſecond over 0 as you ſee, and ſo many points as are in that manner placed, of ſo many figures the root demanded will conſiſt.

$\dot{1}0\dot{2}4$

IX. Having thus prepared your number, you may ſee it diſtributed by the points into ſeveral ſquares: So in the laſt *Example*, 10 is the firſt ſquare and 24 the ſecond; likewiſe if this number 144 were propounded for extraction, after points are duly placed according to the laſt Rule, you will ſee 1 to be the firſt ſquare and 44 rhe ſecond.

$\dot{1}4\dot{4}$

X. Having drawn a crooked line on the right hand of the number propounded for extraction (after the ſame manner as is uſually done in Diviſion to denote the place of the Quotient,) find the root of the firſt ſquare, and place it in the Quotient; So I find, by the ſixth Rule aforegoing, 3 to be the correſpondent root of 10; wherefore I write 3 in the Quotient, and then the Work will ſtand as you ſee.

```
 ·  ·
1024 (3
```

XI. Subſcribe the ſquare of the figure placed in the Quotient under the firſt ſquare of the number given, as you ſee in the Margin.

```
 ·  ·
1024 (3
9
```

XII. Having drawn a line under the ſquare (of the figure placed in the Quotient, ſubſcribed as aforeſaid,) ſubtract the ſame out of the firſt ſquare of the number propounded, and place the remainder orderly underneath the line; ſo the ſquare of 3 which is 9 being ſubtracted from 10, the remainder is 1, and the Work will ſtand as you ſee in the Margin.

```
 ·  ·
1024 (3
 9
———
 1
```

XIII. To the ſaid remainder bring down the next ſquare of the number propounded, that is write down the figures or cyphers ſtanding in the two following places of the number propounded on the right hand of the ſaid remainder, ſo the ſquare 24 being placed next to the remainder 1, there will be found this number 124 which may be called the *Reſolvend.*

```
 ·  ·
1024 (3
 9
———
 124
```

XIV. Double the root being the number placed in the Quotient, and place the ſaid double on the left hand of the Reſolvend, like a Diviſor: So the double of 3 is 6, which being placed before a crooked line on the left

```
 ·  ·
1204 (3
 9
———
6) 124
```

left hand of the Reſolvend 124, the work will ſtand as you ſee.

XV. Let the whole Reſolvend, Except the firſt place thereof on the right hand (being the place of units) be always eſteemed as a Dividend, then demanding how often the Diviſor before found, is contained in the ſaid Dividend, and obſerving in that behalf the Rules before taught in Diviſion, write the anſwer in the Quotient, and alſo on the right hand of the Diviſor, to wit, between the Diviſor and the crooked line: So if you ask how often the Diviſor 6 is found in the Dividend 12, the anſwer is 2, wherefore I write 2 in the Quotient, and alſo after the Diviſor 6, as you ſee in the Margin.

```
     ·  ·
    1024 (32
     9
   ---------
 62)124
```

XVI. Multiply all the number which ſtandeth on the left hand of the Reſolvend, (to wit, before the crooked line) by the figure laſt placed in the Quotient, and write the Product orderly underneath the Reſolvend (to wit, units under units, tens under tens, *&c.*) then having drawn a line under the ſaid Product, ſubtract it from the Reſolvend, and ſubſcribe the remainder under the line: So 62 being multiplied by 2, the Product is 124, which if I ſubtract out of the Reſolvend 124, the remainder is 0; and thus the whole Work being finiſhed, the ſquare root of 1024 (the number propounded) is found to be 32.

```
     ·  ·
    1024 (32
     9
   ---------
 62)124
    124
   ---------
      0
```

Note, 1. When the Product before mentioned exceeds the *Reſolvend* placed above it, the work is erroneous, and then you are to reform it by placing a leſſer figure in the Quotient.

Note 2. For every one of the particular ſquares (diſtinguiſhed by the points) except the firſt on the left hand, a *Reſolvend* is to be ſet apart, by bringing down to the remainder the congruent particular ſquare, as is directed in the 13 *Rule*; and as often as a *Reſolvend* is ſet apart, ſo often a new *Diviſor* is to be found by doubling or multiplying by 2 all the root in the quotient (conſiſting of what number of places ſoever.)

Note 3. The work of the 10, 11, and 12 *Rules* for finding of the firſt figure in the *root*, is but once uſed in the extraction of the root of a number conſiſting of what number of places ſoever; but the Work of the 13, 14, 15, and 16 Rules is to be repeated for the finding of every place in the root, except the firſt.

The practice of theſe 3 *Notes* will be ſeen in the following *Examples*.

Example 1. Let it be required to extract the ſquare root of 43623.

```
•  • •
43623 (2
4
————————
0
```

Having diſtributed the number propounded into ſeveral ſquares by points, as is directed in the eighth Rule of this Chapter, I demand the ſquare root of 4 the firſt ſquare, which I find by the 5 rule of this Chapter to be 2; wherefore placing 2 in the quotient, and the ſquare thereof, which is 4, under the firſt ſquare 4, I draw a line, and ſubtracting 4 from 4 the remainder is 0, which I ſubſcribe underneath the line. This is always the firſt Work, which is no more repeated in the whole Extraction (as was intimated in the third *Note* aforegoing.)

Then bringing down the next ſquare, which is 36, and placing it next after the remainder 0, the *Reſolvend* is 36; and doubling the root 2 in the quotient, the Product is 4 for a *Diviſor* (by the 13 and 14 Rules) and the *Dividend* will be 3 (by the 15 Rule;) wherefore I demand how often the *Diviſor* 4 is contained in the *Dividend* 3, and not finding it once contained in it, I place 0 in the quotient, and alſo next after the *Diviſor* 4; and becauſe the Product of 40 multiplied by 0 (the laſt Character in the quotient) is 0, the *Reſolvend* 36, from which the ſaid Product ought to be deducted, remains the ſame without alteration; therefore I bring down 23 the next ſquare, and place it after the remainder 36, ſo will 03623 be a new *Reſolvend*, then doubling the whole root in the quotient, which is 20, the *Diviſor* will be 40 (according to the ſecond *Note* before mentioned,) and the *Dividend* will be 362 (to wit, all the *Reſolvend* except the firſt place on the right hand by Rule 15.) wherefore I demand how often the *Diviſor* 40 is contained in the *Dividend* 362, or how often 4 in 36, and though it be 9 times in it, yet (according to the firſt *Note* aforegoing) I can take but 8, (for if I ſhould take 9, and proceed according to the 15

```
    .  .  .
    43623 (20
    4
-------------
40) 036
```

```
    .  .  .
    43623
       .
    4
-------------
40) 03623
```

and 16 Rules, a number would ariſe greater than the *Reſolvend*, from which ſuch number ariſing ought to be ſubtracted ;) wherefore I write 8 in the quotient, and alſo after the Diviſor 40; this done, I multiply 408 (the number on the left hand of the *Reſolvend*) by 8 the figure laſt placed in the quotient, and the Product, to wit, 3264 I ſubſcribe under, and ſubtract from the *Reſolvend* 3623, ſo there will remain 359, thus the work being finiſhed I find 208 to be the number of units contained in the root ſought; and becauſe after the extraction is ended there happens to be a remainder, to wit, 359, I conclude that the root ſought is greater than the ſaid 208, but leſs than 209, yet how much it is greater than 208, no Rules of Art hitherto known will exactly diſcover although we may proceed infinitely near, as in the next Rule will be manifeſt.

```
   . . .
  43623(208
  4
---------
408)03623
     3264
    -----
      359
```

XVII. To find the fractional part of the root very near, a competent number of pairs of cyphers, to wit, 00, 0000, 000000, or 00000000, &c. are to be annexed to the number firſt propounded, then eſteeming the number propounded with the cyphers annexed to be but one entire number, the extraction is to be made according to the precedent Rules, and look how many points were placed over the number firſt given, ſo many places of Integers will be in the root, the reſt of the root towards the right hand will be the Numerator of a decimal fraction, which Numerator conſiſteth of ſo many places as there were points over the cyphers annexed: So if 43623 were given as before, to find the root thereof (according to this rule) annex cyphers in this manner, and then if you extract it according to the Rules aforegoing, you

```
 .  .  .  .  .  .
43623.000000 (208 .861, &c,
```

will find the root ariſing in the quotient to be 208 .861, that is 208 $\frac{861}{1000}$; and becauſe after the extraction is finiſht there happens to be a remainder, I conclude that 208 $\frac{862}{1000}$ is leſs than the true or exact root, but 208 $\frac{862}{1000}$ is greater than it; ſo that by annexing three pairs of Cyphers to the number propounded, you will not miſs $\frac{1}{1000}$ part of an unit of the true *root*; alſo by annexing 4 pairs of cyphers, you will not miſs $\frac{1}{10000}$ part of an unit, and in that order you may proceed infinitely near, when you cannot obtain the true root. The whole operation of the ſaid *Example* here followeth.

```
          .  .  .   .  .  .
         43623.000000 (208. 861, &c.
         4              The root
  408) 03623
        3264
  4168) 35900
        33344
  41766) 255600
         250596
  417721) 500400
          417721
           82679
```

Again, if 10 were propounded to be extracted, you muſt prepare it thus,

10.00000000000000 (

And then the root thereof being extracted will be —— } $3\frac{1622776}{10000000}$, &c.

Which (according to the third Rule of the 22 Chapter) may be written thus —— } 3. 1622776, &c.

See here part of the Work in the extraction of the *Root* of 10, which may give you a light and underſtanding of the reſt.

```
      ·  ·  ·  ·  ·  ·  ·  ·
    10.00000000000000 (3. 16227, &c.
          ·  ·  ·  ·  ·

   9
 ---------
 61 ) 100
       61
 ---------
 626 ) 3900
       3756
 ------------
 6322 ) 14400
        12644
 --------------
 63242 ) 175600
         126484
 ------------------
 632447 ) 4911600
          4427129
 ------------------
           484471
```

XVIII. The

The Proof.

XVIII. The extraction of the ſquare root is proved by multiplying the root by it ſelf, for that done, the Product (in ſuch caſe, when there is no remainder after the extraction is finiſhed) will be equal to the number whoſe ſquare root was enquired; ſo in the firſt *Example* of this *Chapter*, the root 32 being multiplied by it ſelf, produceth 1024 the number propounded: But when after the extraction is finiſhed, there happeneth to be a remainder, and that the root is found as near as you pleaſe, in a mixt number of Integers and Decimal parts (by annexing cyphers, as in the 17 Rule) then ſuch mixt number being multiplied by it ſelf muſt produce a mixt number leſs than the number firſt propounded for extraction, yet ſo near unto it, that if the figure ſtanding in the laſt place of the Numerator of the Decimal fraction in the root be made greater by 1, and then the mixt number ſo increaſed be multiplied by it ſelf, the Product muſt be greater than the number firſt propounded: So in the *Example* of the 17 *Rule*, if 208. 861 be multiplied by it ſelf, it produceth 43622. 917, &c. which is leſs than the propounded number 43623, but if 208. 862 be multiplied by it ſelf, the Product will be 43623. 335, &c. which is greater than 43623.

To extract the ſquare root of a Fraction.

XIX. The ſquare root of a Fraction is found in this manner, *viz.* extract the root of the Numerator (by the precedent Rules of this Chapter) which root ſhall be a new Numerator. Alſo the root of the Denominator is to be taken for a new Denominator: So the new Fraction ſhall be the ſquare root of the Fraction firſt propounded ed. Thus the *ſquare root* of $\frac{9}{16}$ is $\frac{3}{4}$, *viz.* the *root* of 9 is 3 for a new *numerator*, alſo the *root* of 16 is 4 for a new *denominator.* In like manner the *ſquare root* of $\frac{1}{4}$ is $\frac{1}{2}$ But here note diligently, that if the *Fraction* whoſe *ſquare root* is required be not in its leaſt terms, it muſt firſt of all be reduced by the 4th *Rule* of the 17th *Chap.* before any extraction be made; for oftentimes it happens that the *Fraction* firſt given hath not a perfect *root*, but when ſuch *Fraction* is reduced into its leaſt terms, the *root thereof* may be extracted: So in this *Fraction* $\frac{8}{18}$, each term is *incommenſurable* to its *ſquare root*, *viz.* neither 8 nor 18 hath a *ſquare root* expreſſible by any true or rational number; but the ſaid $\frac{8}{18}$ being reduced to its leaſt terms $\frac{4}{9}$, the *root* of this may be extracted, for the *root* of 4 is 2 for a new *Numerator*; alſo the *root* of 9 3 for a new *Denominator*, ſo that $\frac{2}{3}$ is found to be the *ſquare root* of $\frac{4}{9}$, (equivalent unto $\frac{8}{18}$.)

XX. When either the Numerator or Denominator of a Fraction hath not a perfect ſquare root, ſuch root is uſually expreſt by prefixing this Character, $\surd$ or $\surd$q. before the Fraction given: So the *ſquare root* of $\frac{13}{16}$ is ſignified thus, $\surd\,\frac{13}{16}$, or thus $\surd$q. $\frac{13}{16}$, becauſe the *root* of $\frac{13}{16}$ cannot be expreſt by any true or rational number whatſoever, yet it may be found very near, as in the next *Rule.*

To extract the ſquare root near, of a fraction incommenſurable to its ſquare root.

XXI. The *ſquare root* of a *Fraction* which is incommenſurable to its *root*, may be found near, in this manner, *viz.* reduce the fraction propoſed into a decimal by the third Rule of the 23 Chapter: The more places are in the decimal, the nearer will the root be found, but the decimal muſt conſiſt of an even number of places, *viz.* either of two, four, ſix, eight, or ten, *&c.* places; then extract the *ſquare root* of that decimal, as if it were a whole number, according to the Rules aforegoing, which *root* found ſhall be a decimal expreſſing near the *ſquare root* of the fraction propoſed.

So if the *ſquare root* of $\frac{13}{16}$ be required near, reduce the ſaid $\frac{13}{16}$ into a decimal (by the 3d *Rule* of the 23d *Chapter)* which will be found .81250000, *&c.* Then extracting the *ſquare-root* thereof as if it were a whole number, it will be found .9013 very near.

XXII. The *ſquare root* of a mixt number commenſurable to its *root* is found in the ſame manner as in the 19th Rule of this Chapter, the mixt number being firſt reduced into an improper fraction by the 10th Rule of the 17th Chapter.

To extract the ſquare root of a mixt number.

So the *ſquare root* of $34\frac{33}{64}$ will be found $5\frac{7}{8}$ *viz.* $34\frac{33}{64}$ being reduced into the improper Fraction $\frac{2209}{64}$, the *ſquare root* of the Numerator 2209 will be 47 for a new Numerator; alſo the *ſquare root* of the *Denominator* 64 is 8, for a new *Denominator*; ſo is found $\frac{47}{8}$, which by (the 13th Rule of the 17th Chapter) is $5\frac{7}{8}$ the *ſquare root* ſought. And here the ſame Caution is to be obſerved as in the 19th Rule of this Chapter; *viz.* The fractional part of the mixt number, or the improper fraction equivalent unto the mixt number, muſt be in the *leaſt terms* before any extraction be made.

To find the ſquare root near, of a mixt number incommenſurable to its root.

XXIII. When the mixt number given is incommenſurable, to its *ſquare root*, prefixing this Character before it, *viz.* √ or √ q. So the *ſquare root*, of 7 $\frac{2}{3}$ will be thus expreſſed, √ 7 $\frac{2}{3}$ or √ q. 7 $\frac{2}{3}$: But if you deſire to find the *ſquare root* near of a mixt number *incommenſurable* to its *root*, reduce the fractional part of the mixt number into a Decimal of an even number of places, as in the 21 Rule of this Chapter, and annex the Decimal ſo found unto the whole part of the mixt number; then eſteeming the ſaid whole number and Decimal as one entire number, extract the *ſquare root* thereof according to the aforegoing Rules of this Chapter, and from the root found, cut off always to the right hand, ſo many places as there are points over the Decimal annexed; which number ſo cut off ſhall be a Decimal, ſhewing the fractional part of the root, and that on the left hand ſhall be the whole part of the root; ſo the *ſquare root* of 7 $\frac{2}{3}$ will be found 2. 7688 very near.

CHAP. XXXIII.

The Extraction of the Cube Root.

I. THE Extraction of the Cube Root is that, by which having a number given, we find another number, which being firſt multiplied by it ſelf, and then by the Product, produceth the number given.

II.

A Cubical number.

II. In the Extraction of the Cube root, the number propounded is always conceived to be a Cube number, that is, a certain number of little Cubes, comprehended within one entire great Cube, and the root or number required is the ſide of that great Cube: what a Cube is may be well expreſs'd by a Die, which indeed is a little Cube it ſelf; wherefore if you place four Dice in a ſquare form, that is, laying two and two in a rank, you ſhall have a ſquare containing four Dice, upon which if you yet erect ſuch another ſquare of Dice, you ſhall have a great entire Cube comprehending two times 4, that is 8 Dice or little Cubes; and here 8 is the Cube number given, and 2 is the root, or number required: In like manner if you rank 25 Dice in a ſquare form, *viz.* laying 5 in a rank, you have a ſquare containing 25 Dice, now upon this ſquare of Dice if you erect four other ſuch ſquares one upon another, you ſhall have a great entire Cube comprehending 5 times 25, this is 125 little Cubes, and in this caſe 125 is the Cube number propounded, and 5 the root or number required.

III. A Cube number is either ſingle or compound.

A ſingle Cube number.

IV. A ſingle Cube number is that, which being produced by the multiplication of one ſingle figure firſt by it ſelf, and then by the product is always leſs than 1000. So 125 is a ſingle Cube number produced by 5 multiplied firſt by it ſelf, and then by 25 the product; for 5 times 5 is 25, and 5 times 25 is 125.

V. All the ſingle Cube numbers, and ſquare numbers, together with their reſpective roots, are expreſſed in the Table following.

Cubes.	1	8	27	64	125	216	343	512	729
Squares.	1	4	9	16	25	36	49	64	81
Roots.	1	2	3	4	5	6	7	8	9

Here in the uppermoſt rank of the Table are placed the ſingle Cube numbers of the particular figures 1,2,3,4,5,6,7,8,9. in the next the ſquares of thoſe figures, and in the loweſt rank the figures themſelves being the reſpective roots of the Cubes and Squares in the uppermoſt ranks; and therefore the *Cube root* of 125 being demanded, the Anſwer is 5, and the *Cube root* of 216 being required, the Table will give you 6, and ſo of the reſt.

VI. When a Cube number is given, that exceeds not 1000, and yet is none of the Cube numbers mentioned in the Table, for this root you are to take the root of the Cube number that being leſs, yet comes neareſt unto it: ſo 125 being given, the root that belongs unto it is 5.

A Compound Cube number.

VII. A compound Cube number is that, which being produced by a number (that conſiſts of more places than one) firſt multiplied by it ſelf, and then by the Product is never leſs than 1000. So 157464 is a *Compound Cube number*, being produced by 54 multiplied firſt by its ſelf, and then by 2916 the Product, for 54 times, 54 is 2916, and then 54 times 2916 is 157464, *the Compound Cube number propounded.*

VIII. To prepare a Cube number for extraction, put a point over the firſt place thereof towards the right hand (to wit the place of Units;) then paſſing over the ſecond and third places, put another point over the fourth, and paſſing over the fifth and ſixth put another point over the ſeventh, and in that order (to wit two places being intermitted between every two adjacent points) place as many points as the number will permit: So 157464 being given, you are to place the points as in the Margin, and ſo many points as are in that manner placed, of ſo many figures the *root* demanded will conſiſt.

$15\dot{7}46\dot{4}$

IX. Having thus prepared your number, you may ſee it diſtributed by the points into ſeveral Cubes: ſo in the ſame example 157 is the *firſt Cube*, and 464 the *ſecond*. In like manner if this Number 7464 were propounded for extraction, after points are duly placed as before, you will ſee 7 to be the *firſt Cube*, and 464 the *ſecond*.

$15\dot{7}46\dot{4}$

$\dot{7}46\dot{4}$

X. Having drawn a crooked line on the right hand of the number propounded to ſignifie a quotient, find the Cube root of the firſt Cube and place it in the quotient: So I finding (by the ſixth Rule of this *Chapter*) 5 to be the correſpondent root of 157, I write 5 in the quotient, and then the work will ſtand as you ſee in the Margin.

$15\dot{7}46\dot{4}$ (5

XI. Subſcribe the Cube of the root placed in the quotient, under the firſt Cube of the number given: So 125 being the Cube of 5 the root (by the fifth

$15\dot{7}46\dot{4}$ (5
125

fifth Rule of this Chapter) I write it under 157 the firſt Cube of the number given, as you ſee in the example.

XII. Draw a line under the Cube ſubſcribed as aforeſaid (to wit, the Cube of the root placed in the quotient) and ſubtract this Cube from the firſt Cube of the number propounded, placing the remainder orderly underneath the line: So 125 the Cube of 5 being ſubtracted from 157, the remainder is 32, and the work will ſtand as you ſee.

$$\begin{array}{l} 15\dot{7}46\dot{4}\ (5 \\ 125 \\ \hline \ 32 \end{array}$$

XIII. To the ſaid remainder bring down the next Cube of the number propounded (to wit, the figures or cyphers that ſtand in the 3 next Places) placing the ſaid Cube next after, to wit, on the right hand of the remainder, ſo the next Cube 464 being placed after the remainder 32, there will be found this number 32464, which may be called the *Reſolvend.*

$$\begin{array}{l} 15\dot{7}46\dot{4}\ (5 \\ 125 \\ \hline 32464\ \textit{reſol.} \\ \hline \end{array}$$

XIV. Having drawn a line underneath the *Reſolvend*, ſquare the root in the quotient, that is, multiply it by it ſelf, and ſubſcribe the triple of the ſaid ſquare or product under the reſolvend in ſuch manner, that the firſt place (to wit, the place of units,) of the ſaid triple ſquare may ſtand directly under the third place (or place of hundreds) in the reſolvend: So the ſquare of the root 5 is 25, the triple whereof is 75, which I ſubſcribe under the Reſolvend

$$\begin{array}{l} 15\dot{7}46\dot{4}\ \ (5 \\ 125 \\ \hline \ 32464\ \textit{reſol.} \\ \hline 75 \end{array}$$

vend in ſuch manner, that the figure 5 which is in the firſt place (to wit, the place of units) in the triple Product 75, may ſtand under 4, which is ſeated in the third place of the reſolvend, as you ſee in the Margin.

XV. Triple the root or number in the quotient, and ſubſcribe this triple number in ſuch manner, that the firſt place thereof (to wit, the place of units) may ſtand directly under the ſecond place (to wit the place of tens) in the Reſolvend: ſo the triple of the root 5 is 15, which I ſubſcribe in ſuch manner, that the figure 5 which is in the firſt place (to wit the place of units) in the ſaid triple number, doth ſtand directly under 6, which is ſeated in the ſecond place of the Reſolvend, and the Work will ſtand as in the Margin.

```
 .  .
157464 ( 5
125
-----------------
 32464 Reſolv.
-----------------
   75
    15
```

XVI. The triple ſquare of the root, and the triple of the root being placed one under the other, as is directed in the 14 and 15 Rules aforegoing, draw a line underneath, and add them together in ſuch order as they are ſeated, and let the ſum be eſteemed as a Diviſor: So the triple ſquare 75, and the triple number 15, being added together as they are ranked in the Work, the ſum will be 765 for a Diviſor.

```
 .  .
157464 ( 5
125
-----------------
 32464 Reſolv.
-----------------
   75
    15
-----------------
   765 Diviſor.
-----------------
```

XVII. Let the whole Reſolvend, except the firſt place thereof towards the right hand (to wit, the place of units) be eſteemed as a Dividend, then demanding how often the firſt figure (towards the left hand) of the Diviſor is contained in the correſpondent part of the Dividend, and obſerving in that behalf the Rules before taught in Diviſion, write the Anſwer in the Quotient: So if I ask how often 7 (the firſt figure of the Diviſor towards the left hand) is contained in 32 (the correſpondent part of the Dividend placed above) the Anſwer will be 4; wherefore I write 4 in the Quotient, as you ſee in the Example.

157464 (54
125
32464 *Reſolv.*
75
15
765 *Diviſor.*

XVIII. Having drawn another line under the Work, multiply the triple ſquare before ſubſcribed (as is directed in the fourteenth Rule) by the figure laſt placed in the quotient, and ſubſcribe this Product under the ſaid triple ſquare; (to wit, units under units, tens under tens, *&c.*) So 75 being multiplied by 4, the Product is 300, which I ſubſcribe under 75 (the triple ſquare) and the work will ſtand as you ſee in the Margin.

157464 (54
125
32464 *Reſolv.*
75
15
765 *Diviſor.*
300

XIX. Multiply the figure laſt placed in the quotient firſt by it ſelf, and then the Product by the triple number before ſubſcribed (as is directed in the 15th Rule of this Chapter;) this done, ſubſcribe the laſt Product under the ſaid triple number (to wit, units under units, tens under tens, &c.) ſo 4 being ſquared or multiplied by it ſelf, the Product is 16, which being multiplied by the triple number 15, the Product is 240, this therefore I ſubſcribe under the aforeſaid triple number 15, and the Work will ſtand as you ſee.

```
 .  .
157464 (54
125
----------------
32464 Reſolvend.
----------------
  75
   15
----------------
  765 Diviſor.
----------------
  300
   240
```

XX. Subſcribe the Cube of the figure laſt placed in the quotient, under the reſolvend, in ſuch manner that the firſt place of this Cube (to wit, the place of units,) may ſtand under the place of units in the reſolvend: So 64 being the Cube of 4, I write it under the reſolvend 32464, in ſuch manner that the figure 4, which is in the place of units in the *Cube* 64, may ſtand under the figure 4 which is ſeated in the place of units of the reſolvend: Obſerve the Work in the Margin,

```
 .  .
157464 (54
125
----------------
32464 Reſolvend.
----------------
  75
   15
----------------
  765 Diviſor.
----------------
  300
   240
    64
```

XXI. Drawing yet another line under the work, add the three laſt numbers together in the ſame order as they are ſeated, and ſubtract the ſum of them from the Reſolvend, placing the remainder orderly underneath: So the ſum of the three laſt numbers, as they are ranked in the Work, is 32464, which if you ſubtract out of the reſolvend 32464, the remainder is 0. Thus the whole Work being finiſhed, the Cube root of 157464 (the number propounded) is found to be 54.

157464 (54
125
32464 *Reſolvend.*
75
15
765 *Diviſor.*
300
240
64
32464
0

Note 1. When the ſum of the three laſt numbers before mentioned is greater than the reſolvend, the Work is erroneous, and then you are to reform it by placing a leſſer figure in the quotient.

Note 2. For every one of the particular Cubes (diſtinguiſhed by the points) except the firſt Cube on the left hand, a reſolvend is to be ſet apart, by bringing down to the remainder the next Cube (as is directed in the 13 Rule) And as often as a reſolvend is ſet apart, ſo often is a new Diviſor to be found, by adding the *triple* of all the *root* in the *quotient* (conſiſting of what number of places ſoever,) to the *triple* of the *ſquare* of ſuch *root*, after they are orderly placed according to the 14 and 15 *Rules.*

Note 3. The Work of the 10, 11 and 12 *Rules* for finding of the firſt figure in the *root* is but once uſed in the extraction of the root of any number whatſoever, but the Work of all the following *Rules* is to be uſed for the finding of every place in the *root*, except the firſt.

The practice of theſe 3 *Notes* will be ſeen in the following *Examples.*

Example 2. Let it be required to extract the *Cube root* of 8302348.

Having diſtributed the number given into ſeveral *Cubes* by points, as is directed in the eighth *Rule* of this *Chapter*, I demand the *Cube root* of 8 (the firſt *Cube* on the left hand) which I find by the fifth *Rule* of this *Chapter* to be 2, wherefore placing 2 in the quotient, and 8 the Cube thereof under 8 the firſt Cube, I draw a line, and ſubtracting 8 out of 8 the remainder is 0, which I ſubſcribe under the line. This is always the firſt Work, and is no more repeated in the whole extraction (as was intimated in the 3 Note aforegoing;) then bringing down the next Cube (to wit, the figures ſtanding in the three following places of the number propounded) which is 302, I place it after the remainder 0, ſo is 302 the *reſolvend*; this done, having drawn a line underneath the *reſolvend*, I ſeek for the triple of the ſquare of the root, *viz.* the *root* in the *quotient* is 2, which multiplied by it ſelf produceth the ſquare 4, the triple whereof is 12, this I ſubſcribe under the *reſolvend*, in ſuch manner that the figure 2

.   .   .
8302348 (2
8
———
0

in the units place of this triple ſquare 12, may ſtand directly under the figure 3, which is ſeated in the third place of the *Reſolvend*, (to wit, the place of hundreds) according to the 14th Rule aforegoing; Again, I triple the root 2, which produceth 6, and ſubſcribe this triple number 6 under the ſecond place (or place of tens) in the *Reſolvend*, to wit, under 0 (according to the 15th *Rule* of this *Chapter*;) then drawing a line under the Work, and adding together the ſaid two numbers laſt ſubſcribed, as they are ranked, the ſum of them is 126 for a Diviſor (according to the 16th *Rule* aforegoing.)

8302348 (2
8

0302 *Reſolvend.*

12
 06

126 *Diviſor.*

That done, eſteeming 30, to wit, all the places except the firſt or place of units in the *Reſolvend*, as a *Dividend*, I demand how often the *diviſor* 126 is contained in 30, and not finding it once contained therein, I write 0 in the *quotient*; and now becauſe the ſum of the three numbers which ought to have been Produced (according to the 18, 19, and 20 *Rules* of this *Chapter*) by the multiplication of 0 (which was laſt placed in the *Quotient*) amounts to 0; the *Reſolvend* 302, out of which the ſaid ſum ſhould have been ſubtracted, remains the ſame without alteration, wherefore having drawn a line under the Work, I write down a new the old *Reſolvend* 302, and bringing down the next *Cube* 348, I annex it to the ſaid

302; ſo there will be a new *reſolvend*, to wit, 302348.

Then ſquaring the root 20 (that is multiplying of it by it ſelf) the Product is 400, which I triple or multiply by 3, and ſubſcribe the Product 1200 underneath the new reſolvend in ſuch manner, that the place of units in this triple quadrate 1200 may ſtand under the place of hundreds, or third place of the reſolvend 302348, to wit, under 3 (according to the 14th *Rule*.) Again, I ſubſcribe the triple of the *root* 20, which is 60, in ſuch manner that the place of units in this triple root 60 may ſtand under the place of tens or ſecond place of the reſolvend, to wit, under 4, then adding together the two numbers laſt ſubſcribed, to wit, 1200 and 60, in ſuch order as they are ranked in the Work, the ſum is 12060 for a Diviſor.

```
.   .   .
8302348 (202
8
------------
0302 Reſolvend.
------------
12
 06
------------
126 Diviſor.
------------
302348 Reſolvend.
------------
1200
  60
------------
12060 Diviſor.
------------
2400
 240
  08
------------
242408 Ablatitium
------------
 59940
```

Again, eſteeming the whole reſolvend, except the firſt place (or place of units) as a dividend, to wit, 30234, I demand how often 1 (the firſt figure of the diviſor towards the left hand) is contained in 3 the correſpondent part of the Dividend; and though it be three times contained in it, yet (according to the firſt *Note* at the end of the 21 *Rule* of this *Chapter*) I dare take but 2, for if I ſhould take 3, and proceed according to the 18, 19, 20, and 21 *Rules* of this *Chapter*, a Number would ariſe greater than the reſolvend (from which ſuch number ariſing ought to be ſubtracted) wherefore I write 2 in the *Quotient*.

Then multiplying the triple ſquare 1200 before ſubſcribed, by 2 (the figure laſt placed in the quotient,) the Product is 2400, which I ſubſcribe under the ſaid 1200 (to wit, units under units, and tens under tens, &c.) Alſo multiplying the triple root 60 before ſubſcribed, by 4 (the quadrate of 2 the figure laſt placed in the quotient) the product is 240, which I ſubſcribe under the ſaid triple root 60; laſt of all I ſubſcribe 8 the *Cube* of the ſaid new *root* 2, under the place of units or firſt place of the *Reſolvend*, to wit, under 8, and having added together thoſe three Numbers laſt ſubſcribed, to wit, 2400, 240 and 8 as they ſtand in ranks in the Work, the ſum of them is 242408, which being ſubducted from the *reſolvend* 302348, there will remain 59940. Wherefore the Work being finiſhed, I find 202 to be the number of units contained in the *Cube root* of 8302348 the number propounded: And becauſe after the extraction is ended there happens to be a remainder, to wit, 59940, I conclude that the *Cube root* ſought is greater than the ſaid 202, but leſs than 203; yet how much it is greater than 202, no Rules of Art hitherto known will exactly diſcover, although we may proceed infinitely near, as by the next *Rule* will be manifeſt.

XXII. To find the fractional part of the root very near ternaries of Cyphers, to wit, 000, 000000, or 000000000, &c. are to be annexed to the number firſt propounded; then eſteeming the number propounded with the cyphers annexed to be but one entire number, the Extraction is to be made according to the preceding *Rules* of this *Chapter*, and look how many points were placed over the number firſt given, ſo many of the foremoſt places in the Quotient are the Integers or Units contained in the Cube root ſought, and the reſt of the places in the quotient are to be eſteem'd as the Numerator of a Decimal Fraction, which Numerator conſiſteth of ſo many places as there were points over the cyphers firſt annexed: So if 8302348 were given as before, to find the *Cube root* thereof (according to this Rule) annex cyphers in this manner.

$$\dot{8}30\dot{2}34\dot{8},00\dot{0}00\dot{0}\,($$

And then if you proſecute the extraction according to the Rules foregoing, you ſhall find the *Cube root* ſought to be 202. 48, &c. that is, 202 $\frac{48}{100}$ and more; wherefore you may conclude that 202 $\frac{48}{100}$ is leſs than the true *root*, but 202 $\frac{49}{100}$ is greater than it: So that by annexing two ternaries of cyphers, to wit, 6 cyphers, to the number propounded, you will not miſs $\frac{1}{100}$ part of an unit of the true *root*, alſo by annexing 3 ternaries of cyphers, to wit, 9 cyphers, you will not miſs $\frac{1}{1000}$ part of an unit of the true *root*, and in that order you may proceed infinitely near, when you cannot obtain the true *root*. The whole operation of the ſaid Example here followeth, where you may obſerve, that for the more certain and eaſie placing, as well of the numbers which conſtitute the ſeveral Diviſors, as of thoſe which conſtitute the ablatitious Numbers to be ſubtracted from the ſeveral and reſpective Reſolvends, down-right lines are drawn between the particular *Cubes* of the number propounded, firſt diſtinguiſhed by points as before.

8	302	348	000	000	(202.48, &c.
8					
0	302				Reſolvend
1	2				
	06				
1	26				Diviſor
	302	348			Reſolvend
	120	0			
		60			
	120	60			Diviſor
	240	0			
	2	40			
		08			
	242	408			Ablatitium
	59	940	000		Reſolvend
	12	241	2		
		6	06		
	12	247	26		Diviſor
	48	964	8		
		96	96		
			64		
	49	061	824		Ablatitium
	10	878	176	000	Reſolvend
	1	228	972	8	
			60	72	
	1	229	033	52	Diviſor
	9	831	782	4	
		3	886	08	
				512	
	9	835	668	992	Ablatitium
	1	042	507	008	

In like manner the *Cube root* of 2 will be found to be near equal to 1.25992, &c. that is, $1\,,\overset{2}{\circ}\overset{5}{\circ}\overset{9}{\circ}\overset{9}{\circ}\overset{2}{\circ}$ and more.

The Proof. XXIII. The extraction of the Cube root is proved by multiplying the root cubically, to wit, the root being firſt multiplied by it ſelf, and then the product multiplied by the root, the number ariſing or laſt Product (in caſe there be no remainder after the extraction is finiſhed) will be equal to the number propounded: ſo in the firſt example of this *Chapter*, the *Cube root* 54 being multiplied firſt by it ſelf produceth 2916, which being multiplied again by 54 produceth 157464, to wit, the number whoſe *Cube root* was inquired. But when after the Extraction is finiſhed, there happeneth to be a remainder, and that the *root* is found as near as you pleaſe in *Integers* and *Decimal parts* (by annexing cyphers as in the 22 *Rule* of this *Chapter*,) then ſuch mixt number expreſſing the *root*, being multiplied cubically, muſt produce a mixt number leſs than the number firſt propounded, yet ſo near unto it, that if the figure ſtanding in the laſt place of the *decimal fraction* in the *root* be made greater by 1, and the mixt number ſo increaſed be multiplied cubically, the Product muſt be greater than the number firſt propounded: So in the Example of the 22 *rule* of this *Chapter*, if 202.48 be multiplied cubically, it produceth 8301305.49, &c. which is leſs than the propounded number 8302348, but if 202.49 be multiplied cubically, there will ariſe 8302535.49, &c. which is greater than the ſaid given number.

XXIV. The Cube root of a Fraction is found in this manner, *viz.* extract the Cube root of the

Numerator (according to the aforegoing Rules,) which root reſerve for a new Numerator; alſo the Cube root of the Denominator ſhall be a new Denominator; laſtly this new Fraction ſhall be the Cube root of the Fraction firſt propounded: ſo the *Cube root* of $\frac{8}{27}$ is $\frac{2}{3}$, for the *Cube root* of 8 is 2 for a new *Numerator*, alſo the *cube root* of 27 is 3 for a new *Denominator*. In like manner the *cube root* of $\frac{1}{8}$ is $\frac{1}{2}$. But here note diligently, that the *fraction* whoſe *cube root* is required, muſt be in its leaſt terms before any Extraction be made; for oftentimes it happens that the *fraction* firſt given hath not a perfect root, albeit, when ſuch *fraction* is reduced into its leaſt terms, the root thereof may be extracted: ſo in this *Fraction* $\frac{16}{54}$ neither the *numerator* nor *denominator* hath a perfect *cube root*, yet the ſaid $\frac{16}{54}$ being reduced to its leaſt terms $\frac{8}{27}$, (by the fourth Rule of the 17 Chapter) the *cube root* of this may be extracted, for the *cube root* of 8 is 2 for a new *numerator*, alſo the *cube root* of 27 is 3 for a new *denominator*, ſo that the *cube root* of $\frac{8}{27}$ (which is equal to $\frac{16}{54}$) is found to be $\frac{2}{3}$.

To extract the Cube root of a Fraction.

XXV. The Cube root of a fraction which hath not a perfect Cube root may be found near in this manner, *viz.* reduce the Fraction given into a Decimal Fraction, by the third Rule of the 23 Chapter, the more places are in the Decimal, the nearer will the root be found, but the Decimal muſt conſiſt of ternaries of places, to wit, either of three, ſix, nine, or twelve, &c. places; then extract the Cube root of the Numerator of that Decimal, as if it were a whole number (according to the Rules before given,) which root found ſhall be a Decimal expreſſing near the Cube root of the Fraction propounded.

So if the *cube root* of $\frac{2}{3}$ were required, I reduce the ſaid $\frac{2}{3}$ into a *decimal*, whoſe *numerator* may conſiſt of *ternaries* of places, to wit into this, *666666666666* &c. then extracting the *cube root* thereof, I find 8735, which is very near the *cube root* of $\frac{2}{3}$.

XXVI. The Cube root of a mixt number commenſurable to its root may be found in the ſame manner as in the 24 Rule of this Chapter, the mixt number being firſt reduced into an improper Fraction (by the 10 Rule of the 17 Chapter.)

So the *cube root* of $12\frac{19}{27}$ will be found to be $2\frac{1}{3}$, *viz.* reducing $12\frac{19}{27}$ into this improper fraction $\frac{343}{27}$, the cube root of $\frac{343}{27}$ will be found $\frac{7}{3}$ or $2\frac{1}{3}$. And here the ſame caution is to be obſerved as in the 24 *Rule* of this *Chapter*, *viz.* the fractional part of the mixt number, or the improper fraction equivalent unto the mixt number, muſt be expreſſed by a *Numerator* and *Denominator* in the leaſt terms before any extraction be made.

XXVII. When the mixt number, whoſe Cube root is required, hath not a perfect cube root, this character, √c. is uſually prefixed before ſuch mixt number; ſo the *cube root* of $2\frac{3}{4}$ is thus expreſſed, √c. $2\frac{3}{4}$. Likewiſe √c. $\frac{5}{8}$ denotes the *cube root* of $\frac{5}{8}$, which is a fraction, whoſe *cube root* is inexpreſſible by any true or rational number: But if you deſire to know the *cube root* near of a mixt number which hath not a perfect *cube root*, reduce the fractional part of the mixt number into a *decimal* (as in the 25 *Rule* of this *Chapter*) and annex the *decimal* ſo found unto the Integers of the mixt number; then eſteeming the ſaid Integers with the *decimal* ſo annexed as one entire number, extract the *cube root* thereof and from the *root* found cut off always to the right hand ſo many places as there were points over the ſaid *decimal* annexed, which places ſo cut off ſhall be the fractional part of the *root*, and thoſe remaining on the left hand ſhall be the Integers of the *root*: So the *cube root* of $2\frac{3}{8}$ will be found 1 .334, and more.

XXVIII. I might here proceed to ſhew the extraction of the *roots* of the *Biquadrate* (or fourth *Power*) the fifth *Power*, &c. but their operations being exceeding tedious, and hardly intelligible without the knowledge of *Algebra*, I ſhall only in this place touch upon the Extraction of the *Biquadrate root*, becauſe it may be extracted by the Rules delivered in the 32 *Chapter*, and referr the more curious Arithmetician for further ſatisfaction in this matter, to my Treatiſe of the Elements of *Algebra*.

To extract the Biquadrate *root.*

XXIX. A quadrate or ſquare number multiplied by it ſelf produceth a *Biquadrate* number: So 4 multiplied by it ſelf produceth the *Biquadrate* 16. Therefore if a number be propounded and the *Biquadrate root* thereof be required, firſt extract the *quadrate* or *ſquare root* of the number propounded, and then extract the *ſquare root* of that *root* for the *Biquadrate root* ſought. Thus if 20736 be a number propounded, the *Biquadrate root* thereof will be found 12: For the *ſquare root* of 20736 is 144, and the *ſquare root* of 144 is 12. When the number given hath not a perfect *Biquadrate root*, you are to annex *quaternaries* of *Cyphers*, to wit, either 4, 8, 12, or 16, &c. *cyphers*, and then proceed as before; ſo will you find the *root* near, whoſe fractional part will be a decimal. Thus the *Biquadrate root* of 7 will be found near 1. 62.

CHAP. XXXIV.

The Relation of Numbers in Quantity.

I. THus far ſingle Arithmetick: Comparative Arithmetick inſues, which is wrought by numbers, as they are conſidered to have Relation one to another.

Boetius Ar. l. 1. *cap.* 21. II. This Relation conſiſts in quantity, or quality.

III. Relation in quantity is the reference or reſpect that the numbers themſelves have one unto another: As when the compariſon is made between 6 and 2, or 2 and 6: 5 and 3, or 3 and 5.

IV. Here the Terms or Numbers propounded are always two, whereof the firſt is called the Antecedent, and the other the Conſequent: So in the firſt Example, 6 is the Antecedent, and 2 the Conſequent: And in the ſecond, 2 is the Antecedent, and 6 the Conſequent.

V. Relation in Quantity conſiſts either in the difference, or elſe in the rate or reaſon that is found betwixt the Terms propounded.

Difference. VI. The difference of two numbers is the remainder, which is left after ſubtraction of the leſs out of the greater: So 6 and 2 being the terms propounded, 4 is the difference betwixt them: for if you ſubtract 2 out of 6, the remainder is 4.

Rate or Reaſon.

VII. The rate or reaſon betwixt two numbers is the quotient of the Antecedent divided by the Conſequent: So if it be demanded what Rate or Reaſon 6 hath to 2, I anſwer, triple Reaſon: For if you divide 6 the Antecedent, by 2 the Conſequent, the Quotient is 3, 2 being contained juſt 3 times in 6. In like manner is there ſubtriple Reaſon betwixt 2 and 6, for if you divide 2 by 6, the Quotient is $\frac{2}{6}$, or (which is all one) $\frac{1}{3}$, becauſe 6 being not once found in 2, there remains 2 for the Numerator, 6 the Diviſor being the Denominator of the Fraction given you in the Quotient, according to the 9th *Rule* of the 16 *Chapter* aforegoing.

VIII. This Rate or Reaſon of Numbers is either equal or unequal.

Equal Reaſon.

IX. Equal Reaſon is the Relation that equal numbers have unto one another: as 5 to 5, 6 to 6, 7 to 7, &c.

X. Here the one being divided by the other, the Quotient is always an Unit: For if it be demanded how often 5 is in 5, the anſwer is 1.

Unequal Reaſon.

XI. Unequal Reaſon is the relation that unequal Numbers have one unto another: and this is either of the greater to the leſs, or of the leſs to the greater.

XII. Unequal Reaſon of the greater to the leſs, is when the greater Term is Antecedent: as of 6 to 2, 5 to 3, and the like.

XIII. Here the Quotient of the Antecedent divided by the Conſequent is always greater than an Unit: So 6 divided by 2, the Quotient is

3, and 5 divided by 3, the quotient is $1\frac{2}{3}$.

XIV. Unequal reaſon of the leſs to the greater, is when the leſſer Term is Antecedent: as of 2 to 6, 3 to 5, &c.

XV. Here the quotient of the Antecedent divided by the conſequent is always leſs than an unit: So 2 divided by 6, the quotient is $\frac{2}{6}$ or $\frac{1}{3}$; and 3 divided by 5, the quotient is $\frac{3}{5}$.

XVI. Each of theſe kinds of unequal Reaſon is again ſubdivided into five other kinds or varieties, whereof the three firſt are ſimple, and the other two are mixt.

XVII. The ſimple kinds of unequal reaſon are, 1. Manifold. 2. Superparticular. 3. Superpartient.

Manifold Reaſon.

XVIII. Manifold Reaſon of the greater to the leſs is, when the Conſequent is contained in the Antecedent divers times without any part remaining: As 4 to 2, 8 to 4, 16 to 8, which is called Double reaſon, becauſe the leſs is contained twice in the greater; ſo 6 to 2 is triple reaſon, 8 to 2 fourfold reaſon, &c.

XIX. Here the quotient of the Antecedent divided by the conſequent is always a whole number: So 8 divided by 2, the quotient is 4.

Submanifold.

XX. The oppoſite of this kind, *viz.* of the leſs to the greater, is called ſubmanifold: Examples hereof are 2 to 4, 4 to 8, 8 to 16, &c. Likewiſe 2 to 6, 2 to 8, 2 to 10, &c.

Superparticular.

XXI. Superparticular is, when the Antecedent contains the conſequent once, and beſides an aliquot part of the conſequent;

quent; that is, an half, a third, a fourth, or a fifth part, &c. of the conſequent, as 3 to 2, 4 to 3, 5 to 4, 6 to 5, and the like; here three divided by 2, the quotient is $1\frac{1}{2}$, and 4 being divided by 3 the quotient is $1\frac{1}{3}$. In like manner 5 divided by 4, the quotient is $1\frac{1}{4}$, and 6 divided by 5 the quotient is $1\frac{1}{5}$; wherefore I ſay 2 and half 2 (that is 1) conſtitute 3: So likewiſe 3 and one third part of 3 (*viz.* 1.) conſtitute 4, and ſo of the reſt.

XXII. Here the quotient of the Antecedent divided by the Conſequent is a mixt number, whoſe whole part, as alſo the Numerator of the fraction annexed, is always an unit: As is obſervable in the examples laſt mentioned.

Subſuperparticular.

XXIII. The oppoſite reaſon of this kind is Subſuperparticular, as 2 to 3, 3 to 4, 4 to 5, 5 to 6, &c,

Superpartient.

XXIV. Superpartient is, when the Antecedent contains the Conſequent once, and beſides divers parts of the Conſequent: As 5 to 3, 7 to 5, 7 to 4, 8 to 5, 9 to 5, 11 to 7, &c. here 5 divided by 3, the quotient is $1\frac{2}{3}$, and therefore 5 contains 3 once, and $\frac{2}{3}$ of 3; for 3 and two thirds of 3 (*viz.* 2) conſtitute 5.

XXV. Here the quotient of the Antecedent divided by the conſequent is a mixt number, whoſe whole part being an unit, hath always for the Numerator of the fraction annexed unto it a number compoſed of more units than one: So the conference being made betwixt 5 and 3, and 5 the Antecedent being divided by 3 the conſequent, the quotient is $1\frac{2}{3}$.

Subſuperpartient.

XXVI. The oppoſite of this reaſon is Subſuperpartient: Examples hereof are 3 to 5, 5 to 7, 4 to 7, 5 to 8, 5 to 9, 7 to 11, and the like.

XXVII. The mixt kinds of unequal reaſon are manifold Superparticular, and manifold Superpartient.

Manifold Superparticular.

XXVIII. Manifold Superparticular reaſon is, when the Antecedent contains the conſequent divers times, and beſides an aliquot part of the conſequent: as 5 to 2, 10 to 3, 17 to 4, 21 to 5, and the like.

XXIX. Here the quotient of the Antecedent divided by the conſequent is a mixt number, whoſe whole part conſiſting of more units than one, hath always an unit for the Numerator of the Fraction annexed unto it; ſo 5 divided by 2 the quotient is $2\frac{1}{2}$, and 21 divided by 5, the quotient is $4\frac{1}{5}$.

Submanifold ſuperparticular.

XXX. The oppoſite of this Reaſon is Submanifold Superparticular; as 2 to 5, 2 to 7, 3 to 7, 4 to 9, &c.

Manifold Superpartient.

XXXI. Manifold Superpartient is, when the antecedent contains the conſequent divers times, and beſides divers parts of the conſequent; as 8 to 3, 17 to 5, 19 to 4, 28 to 5, &c.

Submanifold Superpartient.

XXXII. Here the quotient of the Antecedent divided by the Conſequent is a mixt Number, whoſe whole part as alſo the Numerator of the Fraction annexed unto it, is always a Number compoſed of more units than one: So 8 divided by 3, the quotient is $2\frac{2}{3}$, and 28 divided by 5, the quotient is $5\frac{3}{5}$.

XXXIII. The oppoſite here is Submanifold Superpartient: as 3 to 8, 5 to 17, 4 to 19, 5 to 28, and the like.

And theſe are the ſeveral kinds of varieties of the Rates or Reaſons that are found amongſt Numbers, ſo that no two Numbers whatſoever can be named, but the Rate or Reaſon betwixt them is comprehended under one of theſe five kinds.

CHAP. XXXV.

The Relation of Numbers in Quality, where of Arithmetical and Geometrical Proportion.

I. RElation in quality (otherwiſe called Proportion) is either the reference or reſpect that the Reaſons of Numbers have one unto another, or elſe which the differences of numbers have one to another.

Vide Euclid. l. 3. *d.* 5. & *Alſted. Arith. c* 5.

II. Therefore here the Terms propounded ought always to be more than two, for otherwiſe there cannot be a compariſon of Reaſons or Differences in the Plural Number.

III. This proportion is either Arithmetical, or Geometrical.

IV. Arith-

Arithmetical Proportion.

IV. Arithmetical proportion is, when divers numbers differ according to an equal difference, as 2, 4, 6, 8, 10, &c. here 2 is the common difference betwixt 2 and 4, 4 and 6, 6 and 8, 8 and 10, &c. So 1, 2, 3, 4, 5, 6, 7, &c. differ by *Arithmetical* Proportion, 1 being the common difference betwixt them.

V. Arithmetical Proportion is either continued or interrupted.

1. Continued.

VI. Arithmetical Proportion continued is, when divers numbers are linked together by a continual progreſſion of equal differences. Such are the examples laſt propounded, as alſo theſe 1, 3, 5, 7, 9, 11, 13, &c. And 100000, 200000, 300000, 400000, &c.

VII. In a rank of numbers that differ by Arithmetical Proportion continued, the ſum of the firſt and laſt Terms being multiplied by half the number of the Terms, the Product is the total ſum of all the Terms: So it being demanded, how many ſtrokes the Clock ſtrikes betwixt Midnight and Noon; the Terms of the Progreſſion in this queſtion are Twelve, *viz.* 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12. for in that order the Clock ſtrikes, wherefore if I multiply 13 the ſum of 12 and 1 (the firſt and laſt Terms) by 6 (being half the number of the Terms) the Product is 78, which is the total ſum of all the Terms propounded being added together.

VIII. Or thus, Multiply the number of the Terms by the half ſum of the firſt and laſt Terms, and then likewiſe the Product will give you the total of all the Terms: So 13, 11, 9, 7, 5, 3, being given, their total is 48, for 8 the half ſum of 13 and 3, the firſt and laſt Terms being multiplied by 6, the number of the Terms, the Product is 48.

IX. Three numbers being given, that differ by Arithmetical proportion continued, the mean being doubled, is equal to the ſum of the extreams: So 5, 6, 7, being given, 6 being doubled, is equal to the ſum of 5 and 7, the two extreams.

X. Arithmetical proportion may be continued either upwards or downwards.

Upwards.

XI. Upwards, when the terms of the Progreſſion increaſe, as theſe, 2, 4, 6, 8, 10, 12, *&c.* or theſe 1, 2, 3, 4, 5, 6, &c. And this laſt rank is more particularly termed *Natural Progreſſion.*

XII. Here when the firſt term is alſo the common difference of the terms, the laſt term being divided by the number of the terms, the quotient will give you the firſt term of the rank: Again in this caſe the firſt term multiplied by the number of the terms produceth the laſt term: So this rank 3, 6, 9, 12, 15, 18, 21, being propounded, wherein 3 is both the firſt term, as alſo the common difference of the terms; I ſay 21 the laſt term being divided by 7 the number of the Terms, the quotient is 3 the firſt term; contrariwiſe 3 the firſt term multiplied by 7 produceth 21 the laſt term.

XIII. Arithmetical proportion continued downwards is, when the terms of the progreſſion decreaſe: Such as are 35, 32, 29, 26, 23, 20: And 40, 35, 30, 25, 20, 15, 10, 5.

Downwards.

XIV. Here when the laſt term is alſo the common difference of the terms, the firſt term being divided by the Number of the terms, the quotient will give you the laſt term: Again, the laſt term multiplied by the number of the terms, produceth the firſt term of the rank.

This Rule is in the inverſe of the 12th Rule aforegoing.

For Example, this rank 40, 35, 30, 25, 20, 15, 10, ,5 being propounded, in which 5 is both the laſt term, and likewiſe the common difference of the terms, I ſay, 40 the firſt term being divided by 8 the number of the terms, the quotient is 5 the laſt term: On the other ſide 5 the laſt term being multiplied by 8, the product is 40 the firſt term.

XV. Arithmetical Proportion interrupted is, when the Progreſſion is diſcontinued: as in theſe numbers 2, 4, 8, 10; here 2 and 4 being compared with 8 and 10 differ according to Arithmetical proportion, but ſo do not 4 and 8 differ, for 2 is the common difference betwixt 2 and 4, 8 and 10, whereas the difference betwixt 4 and 8, is 4. In like manner 8, 14, 17, 23, differ by Arithmetical proportion interrupted.

2. *Interrupted.*

XVI. Four nmmbers being given, that differ by Arithmetical proportion either continued or interrupted, the ſum of the two means is equal to the ſum of the two extreams: So 5, 6, 7, 8, being given, the ſum of 6 and 7, the two mean numbers, is equal to the ſum of 5 and 8, the two extreams: and 8, 14, 17, and 23, being propounded the ſum of 14 and 17 being added together is equal to the ſnm of 8 and 23.

XVII. Geometrical proportion is, when divers numbers differ according to like Rate or Reaſon: that is, when the reaſons of numbers, being compared together are equal. So 1, 2, 4, 8, 16, 32, &c. which differ one from another by double reaſon, are ſaid to differ by *Geometrical* proportion, for as 1 is half 2, ſo 2 is half 4, 4 half 8, 8 half 16, 16 half 32, &c.

Geometrical Proportion.

XVIII. Geometrical proportion is either continued or interrupted.

1 *Continued.*

XIX. Geometrical proportion continued is, when divers numbers are linked together by a continued progreſſion of the like reaſon: Of this ſort is the example laſt given: For as 1 is to 2, ſo is 2 to 4, 4 to 8, 8 to 16, 16 to 32, &c. So likewiſe the numbers 3, 9, 27, 81, 243, 729, &c. differ by *Geometrical* proportion continued, *viz.* by triple reaſon, each of them being contained three times in the next number that follows it.

XX. In numbers continually proportional from 1, the firſt number from 1 is the root or firſt power, the ſecond is the ſquare or ſecond power, the third the cube or third power, the fourth the Biquadrate or fourth power, the fifth the fifth power, the ſixth the ſixth power, &c. So in this rank of numbers, 1, 3, 9, 27, 81, 243, 729, &c. 3 is the root, 9 the ſquare, 27, the cube, 81 the Biquadrate, 243 the fifth power, 729 the ſixth power, &c.

XXI. The root being multiplied by it ſelf produceth the ſquare, which being again multiplied by the root produceth the cube, and ſo each Proportional being multiplied by the root, produceth the

Mean proportionals.

proportional next above it, and then the numbers comprehended betwixt 1, and the laſt number produced are called mean proportionals: So in this rank of proportional numbers, 1, 2, 4, 8, 16, 32, &c. 2 the root being multiplied by it ſelf produceth 4 the ſquare, which being again multiplied by 2 produceth 8 the cube, then 8 being multiplied by 2, the product is 16 the biquadrate, and ſo of the reſt in their order, and here 2, 4, 8, and 16, are the mean proportionals in the rank propounded.

Continual means. Briggius Arith. Log. c.6.

XXII. If you multiply the root by it ſelf, and conſequently the ſubſequent numbers by themſelves, the numbers intercepted betwixt 1 and the number laſt produced may not unfitly be called continual means: So 2 being given for the root, multiplied by it ſelf, the product is 4, which being again multiplied by it ſelf produceth 16, then 16 in like manner ſquared produceth 256, which likewiſe multiplied by it ſelf produceth 65536, I ſay then that 2, 4, 16, and 256 are continual means betwixt 1 and 65536.

XXIII. The continual means comprehended betwixt any number given and 1, are diſcovered by a continued extraction of the ſquare roots; for example, 65536 being given, the root thereof extracted is 256, whoſe root is 16; then the root of 16 is 4, and the root of 4 is 2; ſo that at laſt I find 256, 16, 4, and 2 to be continual means intercepted betwixt 65536 and 1 as before.

XXIV. In numbers that increaſe by Geometrical proportion continued, if you multiply the laſt term by the quotient of any one of the terms divided by another term, which being leſs is next unto it, and then deducting the firſt term out of that product, divide the remainder by a number that is an unit leſs than the quotient, the laſt quotient will give you the total of all the terms propounded in the progreſſion; ſo this rank 2, 6, 18, 54, 162, 486, 1458, being propounded, wherein the proportionals differ by ſubtriple proportion, I firſt take 2 and 6 the two firſt terms, and dividing 6 by 2, I find the quotient 3, wherefore multiplying 1458 the laſt term, by 3 the quotient, the product is 4374, out of which if I deduct 2 the firſt term, the remainder is 4372, which being divided by 2 (*viz.* a number which is an unit leſs than three the quotient) the laſt quotient gives me 2186, which is the total ſum of the proportionals propounded.

XXV. Three proportionals being given, the ſquare of the mean is equal to the product of the extreams: So 4, 8, and 16 being propounded, 8 times 8 being 64, is equal to 4 times 16, which is likewiſe 64.

2. *Interrupted.*

XXVI. Geometrical Proportion interrupted is, when the progreſſion of like reaſon s diſcontinued, in ſuch ſort that four numbers being given, the like reaſon is not found betwixt the ſecond and third, that is betwixt the firſt and ſecond, and the third and fourth; of this ſort are theſe numbers 2, 4, 16, 32. here as 2 is to 4, ſo is 16 to 32, for they differ by double reaſon; but as 2 is to 4, ſo is not 4 to 16, for 4 and 16 differ by fourfold reaſon, 4 beng contained 4 times in 16: So likewiſe 4, 8, 8, 16, differ according to Geometrical proportion interrupted.

XXVII. The numbers of Multiplication and Diviſion are proportional; for in Multiplication, as 1 is to the Multiplicator, ſo is the Multiplicand to the Product, or as 1 is to the Multiplicand, ſo is the Multiplicator to the Product: Again, in Diviſion, as the Diviſor is to 1, ſo is the Dividend to the Quotient: Or as the Diviſor is to the Dividend, ſo is 1 to the Quotient.

XXVIII. Four proportional Numbers whatſoever being given, the product of the two means is equal to the Product of the two extreams: So 2, 4, 16, 32, being propounded, 4 times 16 (which is 64) is equal to 2 times 32, which is likewiſe 64.

Here endeth the firſt Book, which containeth all that is abſolutely neceſſary, for the full underſtanding of *common or practical Arithmetick.* Such as deſire to ſee how the ſame is performed by artificial, or borrowed numbers, called *Logarithms*, may peruſe Mr. *Wingate's Second Book*, being a diſtinct *Treatiſe* of *Artificial Arithmetick.*

AN
APPENDIX,

CONTAINING

Choice Knowledge in *Arithmetick*, both *Practical* and *Theoretical*; the Contents whereof are expreſs'd in the following Page.

Compoſed by *John Kerſey.*

Teacher of the

MATHEMATICKS,

Vox audita perit, litera ſcripta manet.

THE CONTENTS OF THE APPENDIX.

CHAP.

1. Of Contractions in the *Rule of Three.*

2. Of Rules of Practice by *aliquot Parts.*

3. Of Exchanges of *Coins*, *Weights* and *Meaſures.*

4. Practical queſtions about *Tare*, *Tret*, *Loſs*, *Gain*, *Barter*, *Factorſhip*, and meaſuring of *Tapeſtry.*

5. Of *Intereſt of Money*, and the conſtruction of *Tables* to value *Annuities*, &c.

6. A *demonſtration* of the *Rule of Three.*

7. A *demonſtration* of the *Double Rule of Fellowſhip.*

8. A *demonſtration* of the *Rule of Alligation:* Where alſo of the *compoſition* of *Medicines.*

9. A *demonſtration* of the *Rule* of *Falſe.*

10. A Collection of *choice Queſtions* to exerciſe all the parts of *vulgar Arithmetick*, to which alſo are added various *practical Queſtions*, about the *Menſuration* of *Superficial Figures* and *Solids*, with the *Gauging* of *Veſſels.*

11. *Sports* and *Paſtimes.*

An Explication of ſuch Notes or Characters, which for brevity ſake are uſed in this APPENDIX.

THis + is a Note of *Addition*, ſignifying that the number which followeth ſuch ſign is to be added to the number preceding it; ſo 3 + 4 implieth that 4 is to be added to 3: Sometimes alſo when no number is placed next after the ſaid note, it implieth that the number preceding is not exactly expreſt; ſo the *ſquare root* of two is 1 .414 + or 1 .414, *&c.* that is, $1\frac{414}{1000}$ and ſomewhat more.

This — is a ſign of *Subtraction*, ſignifying that the number which followeth ſuch ſign is to be ſubtracted from the number preceding it; ſo 6 — 2 ſignifieth the difference between 6 and 2, or 2 to be ſubtracted from 6.

This *x* is a ſign of *Multiplication*, ſignifying that the number which precedeth ſuch ſign is to be multiplied into, or by the number following the ſign; So 3 *x* 4 implieth that 3 is to be multiplied by 4; likewiſe by 3 *x* 4 *x* 8 is underſtood the *continual multiplication* of the numbers 3, 4, and 8; *viz.* 3 is to be multiplied by 4, and the product is to be multiplied by 8. Sometimes alſo the ſaid ſign hath reference to as many of the preceding or following numbers as have a little line placed over them; ſo $3\, x\, \overline{2\, x\, 6}$ or $\overline{2\, x\, 6}\, x\, 3$ ſignifieth that 3 is to be multiplied by the *ſum* of 2 and 6. Likewiſ wiſe $8 - 5 \times 3$, or $3 \times \overline{8 - 5}$ implieth that 3 is to be multiplied by the *difference* between 8 and 5: Moreover if A and B repreſent two numbers, then $A \times B$ or $A\,B$ implieth the product of the multiplication of thoſe numbers: Likewiſe $\overline{B - C} \times A$ ſignifieth the product ariſing from the multiplication of the exceſs of the number B above the number C, by (or into) the number A. Again, if A B and A C repreſent two lines, then ▭ $AB \times AC$ implieth a rectangular Figure or long ſquare made of the lines A B and A C.

Numbers placed as you ſee in the Margin, denote a *Diviſor*, a *Dividend*, and a *Quotient*, to wit, 3 the *Diviſor*, 18 the *Dividend*, and 6 the *Quotient*; the like is to be underſtood of other numbers ſo placed.

3) 18 (6

Numbers placed after the manner of a *fraction* denote a *Quotient*, which ariſeth from dividing the *Numerator* by the *Denominator*; ſo $\frac{2 \times 5 \times 6}{3 \times 4}$ is equal to the *Quotient*, which ariſeth from dividing the *Product* of the *continual Multiplication* of 2, 5 and 6 by the *Product* of 3 multiplied by 4.

Four numbers placed as you ſee in the Margin are *Geometrical proportionals*, *viz.* As 2 is to 4, ſo is 6 to 12: or if 2 give 4, then 6 will give 12. Sometimes alſo they are placed thus, 2 .... 4 .... 6 .... 12.

2 . 4 :: 6 . 12

This $=$ is a note of *equality* or *equation*; ſo by $3 + 4 = 5 + 2$ is ſignified that the ſum of 3 and 4 is equal to the ſum of 5 and 2: Alſo $7 - 3 = 9 - 5$ ſignifieth that the *difference* between 7 and 3 is equal to the *difference* between 9 and 5; that is, 7 leſſened by 3 leaves the ſame remainder, as 9 leſſened by 5. Alſo $4 \times 3 = 12$ implieth that the *product* of the multiplication of 4 by 3 is equal to 12.

$>$ This is a ſign of *majority*, ſignifying that the number on the left hand of ſuch ſign is *greater* than the number on the right hand thereof, ſo $5 > 3$ implieth that 5 is *greater* than 3.

$<$ This is a ſign of *minority*, ſignifying that the number on the left hand of ſuch ſign is *leſs* than the number on the right hand thereof; ſo $3 < 5$ implieth that 3 is *leſs* than 5.

This Character $\surd$ or $\surd$ q. ſignifies the ſquare root of the number which follows it, ſo $\surd\,144$ implies the ſquare root of 144, to wit 12.

Alſo this $\surd$ c. ſignifies the cube root of the number which follows it: So $\surd$ c. 1728 ſignifies the cube root of 1728, which cube root will be found to be 12.

AN APPENDIX.

CHAP. I.

Of Contractions in the Rule of Three.

SUch as are well verſ'd in the parts of *Arithmetick*, which have been fully laid open in the precedent Book, and are mindful of the *Notes* or *Symbols* before explained, will find no difficulty in the 1, 2, 3, 4, 5, and 10 Chapters of this *Appendix*, wherein divers compendious operations no leſs delightful than uſeful are methodically handled, and the reſt will be as eaſie to ſuch as are but meanly acquainted with *Geometrical demonſtration*.

II. To repeat the brief ways of *Multiplication* ſet forth in the 10, 11, and 12 *Rules* of the fifth *Chapter*, or thoſe of *Diviſion*, in the 11, 15, and 16 *Rules* of the ſixth *Chapter* aforegoing, would be a ſuperfluous work, and therefore I ſhall preſuppoſe the *Reader* to be throughly acquainted with them, as alſo with competent knowledge in the operations of Fractions both *vulgar* and *decimal.*

III. It will be no ſmall advantage to the Practical *Arithmetician*, to have by heart not only the common Table of *Multiplication*, but this alſo in the *Margin*, to the end that when a number is given to be multiplied or divided by 12, (which happens in the *Reduction* of *ſhillings* to *pence* and the converſe) the *product* or *quotient* may be written down in one line only, as in the Examples following.

$$\left.\begin{array}{c}2\\3\\4\\5\\6\\7\\8\\9\end{array}\right\} \times 12 = \left\{\begin{array}{r}24\\36\\48\\60\\72\\84\\96\\108\end{array}\right.$$

$$\begin{array}{r} 3472 \\ 12 \\ \hline 12)\,41664\,(3472 \end{array} \qquad\qquad \begin{array}{r} 4736 \\ 12 \\ \hline 12)\,56832\,(4736 \end{array}$$

IV. When a whole number is given to be divided by a Diviſor, which is equal to the product of the Multiplication of two ſingle figures, inſtead of dividing by that Diviſor you may firſt divide by one of thoſe ſingle figures, and then divide the quotient by the other, ſo will the laſt quotient be the ſame as if the Diviſion had been finiſht by the Diviſor firſt given: Thus if 3466 *farthings* be given to be reduced to *ſhillings* becauſe $8 \times 6 = 48$, I firſt divide 3466 by

8, ſo there will ariſe 433 for a new *Dividend*, and 2 *farthings* remain; then I divide the ſaid 433 by 6, ſo there will ariſe--$72\frac{1}{6}$, or 72 *ſhillings* 2 *pence*, which with the 2 *farthings* remaining of the firſt *Diviſion* make in all 72 *s.* : $2\frac{1}{2}$ *d.* which is the very *quotient*, when 3466 farthings are divided by 48. Note that you are to reſerve a farthing for every unit remaining of the firſt Diviſion by 8, and 2 pence for every unit remaining of the ſecond Diviſion by 6. The reaſon of the operation is evident, for $\frac{1}{6}$ of $\frac{1}{8} = \frac{1}{48}$.

8) 3466
s. d.
6)433 (72 : $2\frac{1}{2}$

In like manner, if 7136 pence are given to be reduced into pounds, becauſe 240 *d.* = 1 *l.* alſo 6 x 40 =240, therefore if 7136 pence be firſt divided by 6, the quotient will give 1189 ſix pences, and 2 pence remain; then if 1189 be divided by 40, (that is by 4, after 9 the laſt place of the Dividend towards the right hand is cut off) the quotient will be 29 *l.* and there will remain 29 ſix pences, or 14*s.* 6*d.* which together with the 2 pence remaining of the firſt Diviſion, and the ſaid 29 *l.* makes in all 29 *l.* : 14 *s.* : 8*d.* which is the ſame with the quotient, when 7136 pence are divided by 240, for $\frac{1}{40}$ of $\frac{1}{6} = \frac{1}{240}$.

6) 7136
l. s. d.
40) 118|9 (29 : 14 : 8

Again, ſuppoſe 3463 pence are given to be reduced into Shillings, foraſmuch as 4 x 3 = 12, I firſt divide 3463 by 4, ſo there will ariſe 865 for a new Dividend and 3 pence remain: Then I divide the ſaid 865 by 3, ſo there will ariſe $288\frac{1}{3}$ or 288 *s.* 4 *d.*

4) 3463
       s. *d.*
3) 865 (288..7

4 *d.* which with the 3 pence before remaining make 288 *s.* 7 *d.* which is the ſame with the Quotient, when 3463 pence are divided by 12, for $\frac{1}{3}$ of $\frac{1}{4}=\frac{1}{12}$.

V. In the Rule of Three as well direct as inverſe when the Diviſor with either of the other two given numbers may be ſeverally divided by ſome common meaſure, without leaving any remainder, the Quotients may be taken for new terms, and proceeding in like manner as often as is poſſible, the operation according to the tenth Rule of the eighth Chapter, or the ſecond Rule of the ninth Chapter, will be much contracted: So if it be demanded what 52 yards of Cloth will coſt at the rate of 21 *l.* for 14 Yards; the *Anſwer* will be found 78 pounds, in manner following.

y. *l.* *y*
14... 21... 52
2... 3... 52
1... 3... 26..(78

In the firſt rank you may obſerve, that the Diviſor 14 and the ſecond term 21, being ſeverally divided by their common meaſure 7, the three new terms (in the ſecond rank) will be 2, 3, 52. Again, in the ſecond rank, the diviſor 2 and the third term 52 being ſeverally divided by their common meaſure 2, the three new terms (in the third rank) will be 1, 3, 26. Laſtly, working with theſe according to the *Rule* of *Three direct*, the *Anſwer* to the queſtion (or fourth term) will be found to be 78.

Another Example, If 21 men will finiſh a work in 16 Days, what time muſt be allowed to 12 men for the finiſhing of ſuch a work? *Anſwer*, 28 days.

men *days* *men*
21 . . . 16 . . . 12
7 . . . 16 . . . 4
7 . . . 4 . . . 1 (28 *days*.

In the firſt rank you may obſerve, that the Diviſor 12 (for the Rule is inverſe) and the firſt term 21 being ſeverally divided by their common meaſure 3, the three new terms (in the ſecond rank) will be 7, 16, 4. Again in the ſecond rank, the Diviſor 4, and the ſecond term 16, being ſeverally divided by their common meaſure 4, the three new terms in he third rank will be 7, 4, 1. Laſtly, working with theſe as the Rule of three inverſe requires, the *Anſwer* to the queſtion (or fourth term) will be ound 28.

VI. In the Rule of three, as well direct as inverſe, when the Diviſor and either of the other two terms re fractions having a common denominator, the aid denominators may be rejected, and their numerators retained as new terms: So if it be demanded what is the value of $\frac{7}{8}$ of an *Ell*, when $\frac{3}{8}$ of n *Ell* are worth 66 pence, the *Anſwer* will be ound 154 pence, and the work will ſtand as you ee.

$\frac{3}{8}$ . . 66 . . $\frac{7}{8}$
3 . . 66 . . 7
1 . . 22 . . 7 (154

Another Example. If 3 $\frac{3}{4}$ *yards* of Scarlet cloth coſt 8 *l.* 15 *s.* what is the price of one *yard* at that rate? *Anſwer* 2 *l.* 6 *s.* 8 *d.*

$\frac{15}{4}$ . . . $\frac{35}{4}$ . . . 1
15 . . . 35 . . . 1
3 . . . 7 . . . 1 . . (2 $\frac{1}{3}$ *l*

VII. In the Rule of Three as well direct as inverſe when the Diviſor only is a fraction, either of the other two terms may be reduced to a fraction of the ſame Denominator, and then the Denominators may be rejected, as before in the ſixth Rule. Alſo when one of the three given terms is a fraction, and is not the Diviſor, the Diviſor may be reduced to a fraction of the ſame Denominator with the fraction firſt given, and then the common Denominators may be likewiſe cancelled.

An *Example* of the firſt Caſe may be this, if $\frac{7}{8}$ of a *yard* coſt 14 *s.* what is the price of 1 *yard*? *Anſwer* 16 *ſhillings.*

yard. *ſhill.* *yard.*
$\frac{7}{8}$ . . . 14 . . . 1
$\frac{7}{8}$ . . . 14 . . . $\frac{8}{8}$
7 . . . 14 . . . 8 . . (16 *ſhill.*

An *Example* of the ſecond Caſe; if of ſtuff which is $\frac{3}{4}$ of a yard in breadth, 7 yards in length will make a Garment; how much of that ſtuff which is one yard in breadth will be ſufficient for the ſame purpoſe? *Anſwer* 5 $\frac{1}{4}$ *yards.*

Rules of 3 Inverſe. $\begin{cases} \frac{3}{4} \dots 7 \dots 1 \\ \frac{3}{4} \dots 7 \dots \frac{4}{4} \\ 3 \dots 7 \dots 4 \; (\,.. \frac{21}{4} \text{ or } 5\frac{1}{4} \end{cases}$

CHAP. II.

Rules of Practice by Aliquot parts.

I. AN *Aliquot part* takes its name from the *Latine* word *aliquoties*, for (according to *Euclid*) an *aliquot part* is of a greater number ſuch a part, which being taken (*aliquoties* or) certain times doth preciſely conſtitute that greater number; ſo 3 is an *aliquot part* of 12, for 3 taken 4 times doth exactly make 12, without any exceſs or defect; in like manner 4 is an *aliquot part* of 20, becauſe 4 taken 5 times doth preciſely make 20; but 7 is not an *aliquot part* of 20, for 7 taken twice doth want of 20, and being taken thrice doth exceed 20; this kind of *part* laſt mentioned is by *Euclid* called *pars aliquanta*, of which there will be no uſe in this place.

II. When the *Rule of Three direct* hath 1 or an *Integer* for the firſt Term, it is commonly called the *Rule of Practice*, either from the great uſe and practice thereof in common affairs, or elſe for that queſtions of this nature, may be reſolved by operations more ſpeedy and practical than thoſe of the *Rule of Three.*

III. The choiceſt of theſe *Rules* of *Practice* may be reduced to 5 Caſes, *viz.*

When the price of 1 *or an Integer conſiſts,*
1. *Of ſhillings under* 20.
2. *Of pounds and ſhillings.*
3. *Of pence under* 12.
4. *Of ſhillings and pence.*
5. *Of pounds, ſhillings, pence, with parts of a penny.*

All which caſes, with others of the like nature, are handled in their order.

IV. Any even number of ſhillings is either $\frac{1}{10}$ of a pound (that is 2 *ſhillings*,) or elſe is compoſed of $\frac{1}{10}$ *l.* (to wit 2 *s.*) taken certain times: So 8 *s.* is compoſed of $\frac{1}{10}$ *l.* (or 2 *ſhillings*) taken four times, in like manner 18 *s.* is compoſed of $\frac{1}{10}$ *l.* taken nine times.

V. When the price of 1, or an Integer of what name ſoever, 2 is ſhillings, the price of as many Integers as one will of that name is diſcoverable at firſt ſight, to wit by accounting the double of the figure which ſtands in the firſt place (towards the right hand) of the ſaid number of Integers, as ſhillings, and the reſt of the ſaid number as pounds: ſo 345 yards at two ſhillings the yard will coſt 34 *l.* 10 *s.* for the double of 5 is 10, which I write down apart as ſhillings, then eſteeming the remaining figures towards the left hand, to wit 34, as an entire number of pounds, the *Anſwer* will be 34 *l.* 10 *s.* This contraction is nothing elſe, but dividing the number

yard	*ſhill.*	*yards*
1 ...	2 ...	345

Anſwer 34 *l.* 10 *s.*

ber of Integers, whoſe price is required by 10 More *examples* hereof are theſe;

oz. *ſhill.* *oz.*
1 ... 2 ... 2057

l.
Anſw. 205 .. 14

yard *ſhill.* *yards.*
1 ... 2 ... 120

l. *s.*
Anſw. 12 .. 0

VI. When the given price of 1 or Integer is any even number of ſhillings greater than two ſhillings, multiply the number of Integers, whoſe price is required, by half the given number of ſhillings, with this caution, that the double of the figure which ariſeth in the firſt place of the product be written apart as ſhillings, and the reſt of the product as pounds: So if it be demanded what 218 yards at 8 ſhillings the yard will amount unto, the *Anſwer* will be found 87*l.* 4*s.* for I multiply 218 by 4 (which is half 8 the given number of ſhillings) ſaying, 4 times 8 is 32, here the double of 2 (to wit, of that figure which is to poſſeſs the firſt place in the product) is 4, which I ſet apart as ſhillings, keeping 3 in mind for the three tens, again 4 times 1 is 4, which

y. *s.* *y.*
1 ... 8 ... 218
4

87 : 4

with 3 in mind makes 7; laſtly, 4 times 2 makes 8, ſo I conclude that the *Anſwer* to the queſtion is 87 *l.* 4 *s.* The reaſon of this contraction is evident from the fourth and fifth Rules aforegoing. More examples of this Rule are theſe following.

yard *s.* *yard.*
1 . . . 14 . . . 436

l. *s.*
Anſw. 305 . . 4

---

yard *s.* *yard.*
1 . . . 18 . . . 230

l. *s.*
Anſw. 207 . . . 0

VII. Any odd number of ſhillings is either compos'd of $\frac{1}{10}$ *l.* (or 2 *s.*) and of $\frac{1}{20}$ *l.* (or 1 *s.*) or elſe it is compoſed of $\frac{1}{10}$ *l.* (or 2 *s.*) taken certain times, and of $\frac{1}{20}$ *l.* (or 1 *s.*) So 3 *s.* is compos'd of 2 *s.* and 1 *s.* Alſo 7 *s.* is compoſed of 2 *s.* taken three times and of 1 *s.* Likewiſe 13 *s.* is compoſed of 2 *s.* taken ſix times, and of 1 *s.*

VIII. When the given price of 1 or an Integer is an odd number of ſhillings, work for the greateſt even number of ſhillings contained in that odd number, according to the fifth or ſixth Rule aforegoing; then for the odd ſhilling remaining, take $\frac{1}{20}$ of the number of Integers, whoſe price is required (by the 16th Rule of the ſixth Chapter of the preceding Book.) Theſe two reſults added together give the Anſwer to the queſtion;

queſtion: So if it be demanded what 2344 ounces at 13 *s.* the ounce will coſt, the anſwer will be found 1523 *l.* 12 *s.* For if (according to the ſixth Rule of this Chapter) I multiply 2344 by 6, (to wit, by half the remainder, when one is abated from 13 the given number of ſhillings) there will ariſe 1406 *l.* 8 *s.* Then taking $\frac{1}{20}$ of 2344, there will ariſe 117 *l.* 4 *s.* which being added to the former product gives 1523 *l.* 12 *s.* for the *anſwer* to the queſtion.

oz. *ſhill.* *oz.*
1 .. 13 .. 2344
6
l. *s.*
1406 .. 8
117 .. 4
l. *s.*
Anſw. 1523 .. 12

Note, When 5 ſhillings is the given price of 1 or an Integer, the briefeſt way will be to take $\frac{1}{4}$ of the number of Integers, whoſe value is required, for ſuch quotient will give the pounds and ſhillings, which anſwer the queſtion: ſo 2347 ounces at 5 *s.* the ounce amount unto 586 *l.* 15 *s.* for $\frac{1}{4}$ of 2347 is $586\frac{3}{4}$ or 586 *l.* 15 *s.* But when the given price of 1 is any other odd number of ſhillings, this eighth Rule will be as compendious as any other whatſoever.

More Examples of this Rule are theſe following.

yard *ſhill.* *yard.*
1 .... 19 739
l. *s.*
665 ... 2
36 ... 19
Anſw. 702 ... 1

yard *ſhill.* *yards.*
1 . . . 17 . . . 345

l. *s.*
276 . . . 0
17 . . . 5

Anſw. 293 . . . 5

IX. When the given price of 1 or an Integer conſiſts of pounds and ſhillings, firſt multiply the number of Integers whoſe price is required, by the number of pounds in the ſaid given price, and ſubſcribe the product as pounds; then proceed with the ſhillings in the ſaid given price, according to the ſixth or eighth Rule of this Chapter, and having ſubſcribed that which ariſeth under the aforeſaid product of pounds, add them all together for the anſwer of the queſtion: So if it be demanded what 328 hundred weight will amount unto at 2 *l.* 17 *s.* *per* C. (or one hundred weight) the *anſwer* will be found to be 934 *l.* 16 *s.* as by the operation is evident.

C. *l.* *s.* *C.*
1 . . . 2 : 17 . . . 328

l. *s.*
656 . . 0
262 . . 8
16 . . 8

Anſw. 934 : 16

More *Examples* to illuſtrate this *Rule* are theſe following:

C. *l.* *s.* *C.*
1...7 : 12...504

l. *s.*
3528..0
302..8

Anſw. 3830..8

C. *l.* *s.* *C.*
1...5 : 7...129

l. *s.*
645..
38..14
6.. 9

Anſw. 690.. 3

X. Any number of pence under 12 is either an Aliquot part of a ſhilling, or elſe compos'd of Aliquot parts thereof; ſo 3 pence is an Aliquot part, to wit, $\frac{1}{4}$ of a ſhilling. Likewiſe 4 is $\frac{1}{3}$ of 12; moreover 5 pence are compos'd of 2 Aliquot parts, to wit, of 3 pence, which is $\frac{1}{4}$ of a ſhilling, and of 2 pence which is $\frac{1}{6}$ of a ſhilling; all which will readily appear by the following Table.

Pence

Pence.	*Aliquot parts of a Shilling.*
1	$\frac{1}{12}$ (or $\frac{1}{3}$ of $\frac{1}{4}$)
$1\frac{1}{2}$	$\frac{1}{8}$
2	$\frac{1}{6}$
3	$\frac{1}{4}$
4	$\frac{1}{3}$
5	$\frac{1}{4} + \frac{1}{6}$
6	$\frac{1}{2}$
7	$\frac{1}{4} + \frac{1}{3}$
8	$\frac{1}{3} + \frac{1}{3}$
9	$\frac{1}{2} + \frac{1}{4}$
10	$\frac{1}{2} + \frac{1}{3}$
11	$\frac{1}{2} + \frac{1}{3} + \frac{1}{4}$

XI. When the given price of 1 or an Integer is an Aliquot part of a ſhilling, divide the number of Integers whoſe value is required by the denominator of ſuch aliquot part; ſo will the quotient be the number of ſhillings which anſwer the queſtion, which number of ſhillings (when there is occaſion) may be reduced to pounds by the brief way of dividing by 20: So if it be required to know what 2686 ounces at 4 pence the ounce will amount unto; the anſwer will be found 44 *l.* 15 *s.* 4 *d.* for ſince 4 *d.* is an aliquot part, to wit, $\frac{1}{3}$ of a ſhilling, I divide 2686 by 3, ſo will the quotient be 895$\frac{1}{3}$ *s.* or 895 *s.* 4 *d.* which ſhillings being divided by 20, give 44 *l.* 15 *s.* 4 *d.* for the anſwer to the queſtion, as you ſee by the following operation.

oz. *d.* *oz.*
1 . . . 4 . . . 2686

s. *d.*
20) 89|5 . . 4
Anſw. 44 . . 15 . . 4

More Examples of this Rule are theſe following.

yard *d* *yard.*
1 . . . 6 . . . 759

s. *d.*
20) 37|9 . . 6
Anſw. 18 19 6

---

yard *d* *yards.*
1 . . . 1 . . . 204

Anſw. 17 *ſhillings.*

XII. When the given price of an Integer is compos'd of aliquot parts of a ſhilling, divide the number of Integers, whoſe price is required, by the ſeveral denominators of the aliquot parts contained in the given number of pence, then add the quotients together, and the ſum ſhall be the number of ſhillings which anſwer the queſtion : So if it be demanded what 2347 yards of linen cloth will coſt at 9 pence the yard, the anſwer will be found 88 *l.* 0 *s.* 3 *d.* For ſince 9 *d.* is compos'd of 6 *d.* and 3 *d.* to wit, of aliquot parts $\frac{1}{2}$ and $\frac{1}{4}$ of a ſhilling, I firſt divide 2347 by 2 (the denominator of the aliquot part $\frac{1}{2}$) ſo there ariſeth 1173$\frac{1}{2}$, or 1173 *s.* 6 *d.* Again, dividing the ſaid 2347 by 4 (the denominator of the other aliquot part) there will ariſe 586$\frac{3}{4}$, or 586 *s.* 9 *d.* which two quotients being added together give 1760 *s.* 3 *d.* or 88 *l.* 0 *s.* 3 *d.* which is the anſwer of the queſtion. More Examples to illuſtrate this Rule are theſe.

yard *d.* *yards.*
1 ... 9 ... 2347

s. *d.*
1173 : 6
586 : 9

20) 176|0 : 3
l. *s.* *d.*
Anſw. 88 : 0 : 3

yard *d.* *yards*
1 ... 8 ... 782

s. *d.*
260 ... 8
260 ... 8

20) 52|1 ... 4
l. *s.* *d.*
Anſw. 26 .. 1 .. 4

oz. d. oz.
1 . . . 11 . . . 540

180
180
135

20) 49|5 *s.* *d.*
Anſw. 24 . . . 15 : 0

XIII. When the given price of an Integer conſiſts of ſhillings and pence, firſt multiply the number of Integers whoſe value is required by the ſaid given number of ſhillings, and ſubſcribe the product as ſhillings, then divide the ſaid number of Integers by the ſeveral denominators which are correſpondent to the aliquot parts contained in the given number of Pence, and ſubſcribe the quotient or quotients underneath the aforeſaid product of ſhillings, all which being added together give the number of ſhillings which anſwers the queſtion: So if it be demanded what 347 yards of cloth will coſt at the rate of 7*s.* 10*d.* the yard, the anſwer will be found 135*l.* 18*s.* 2*d.* for firſt 347 being multiplied by 7 (the given number of ſhillings) produceth 2429 ſhillings, then dividing 347 by 2 and 3 ſeverally, (becauſe 10 *d.* is com-

yard *s.* *d.* *yards.*
1 . . 7 : 10 . . 347

s. *d.*
7x347= | 2429 :
2) 347(. . | 173 : 6
3) 347(. . | 115 : 8

20) 271|8 : 2
l. *s.* *d.*
Anſw. 135 : 18 : 2

pos'd of $\frac{1}{2}$ and of $\frac{1}{3}$ a ſhilling) the quotients will be 173$\frac{1}{2}$ and 115$\frac{2}{3}$, that is 173 *s.* 6 *d.* and 115 *s.* 8 *d.* Laſtly, the ſum of all is 2718 *s.* 2 *d.* or 135 *l.* 18 *s.* 2 *d.*

More Examples of this kind are theſe.

yard   *s.*   *d.*   *yards,*

1 ... 17 : 9 ... 540

$$\begin{array}{r|l} & \textit{s.} \\ 17 \times 540 = & \left\{\begin{array}{l} 3780 \\ 540. \end{array}\right. \\ \left.\begin{array}{r} 2 \\ 4 \end{array}\right\} \begin{array}{l} 540(.. \\ 540(.. \end{array} & \begin{array}{r} 270 \\ 135 \end{array} \end{array}$$

20) 958|5

l.   *s.*   *d.*

Anſw. 479 : 5 : 0

y.   *s.*   *d.*   *y.*

1 ... 14 : 6 ... 313

$$\begin{array}{rl} & \textit{s.} \\ 14 \times 313 = & \left\{\begin{array}{l} 1252 \\ 313. \end{array}\right. \\ 2\,)\,313\,(. & 156 : 6 \end{array}$$

20) 453|8

Anſw. 226 : 18 : 6

XIV. When the price of an Integer conſiſts of ſhillings and pence, and that ſuch ſhillings and pence jointly conſidered do make an aliquot part of a pound, it will oftentimes be a briefer way than that in the laſt Rule, to divide the number of Integers, whoſe value is required, by the denominator of ſuch aliquot part, ſo will the quotient give the anſwer to the queſtion in pounds and known parts of a pound. Thus if it be demanded what 767 yards will coſt at the rate of 6 *s.* 8 *d.* the yard, the anſwer will be found 255 *l.* 13 *s.* 4 *d.* For ſince 6 *s.* 8 *d.* is an aliquot part, to wit, $\frac{1}{3}$ of a pound, I divide 767 by 3. ſo there ariſeth in the quotient $255\frac{2}{3}$, or 255 *l.* : 13 *s.* : 4 *d.* which is the anſwer of the queſtion. *Note* that the *Aliquot parts* of a pound convenient for this Rule are theſe expreſt in the following *Table.*

```
y.     s.  d.
1 . . . 6 : 8 . . 767
------------------------------
                   l.   s.  d.
3 ) 767 ( 255 . . 13 : 4
```

ſh.	*d*	*Aliquot parts of a pound.*
6	8	$\frac{1}{3}$
3	4	$\frac{1}{6}$
2	6	$\frac{1}{8}$
1	8	$\frac{1}{12}$
1	4	$\frac{1}{15}$
1	3	$\frac{1}{16}$

XV. When the given price of 1 or an Integer conſiſts of pounds, ſhillings and pence, reduce the ſaid pounds and ſhillings all into ſhillings, then proceed according to the 12 Rule of this Chapter : So 517 *C.* at 3 *l.* : 17 *s.* 5 *d. per C.* will be found to amount unto 2001 *l.* 4 *s.* 5 *d.* for having reduced 3 *l.* 17 *s.* into 77 *s.* I multiply 517 by 77, and write down the particular products; then for the 5 pence which is compos'd of the aliquot parts $\frac{1}{4}$ and $\frac{1}{6}$ of a ſhilling, I take $\frac{1}{4}$ and $\frac{1}{6}$ of 517, and ſubſcribe the quotients orderly underneath the aforeſaid products: Laſtly, adding all together the ſum 40024 *s* 5 *d.* or 2001 *l.* 4 *s.* 5 *d.* for the anſwer of the queſtion.

C	*l.*	*s.*	*d.*	C.
1 ...	3 :	17 :	5 ...	517

$$77 \times 517 = \left\{\begin{matrix} 3619 \\ 3619 \end{matrix}\right.$$

4) 517 (.. 129 : 3*d.*
6) 517 (.. 86 : 2

20) 4002|4 : 5
Anſw. 2001 : 4 : 5

More *Examples* of this *Rule* are theſe following.

C.	*l.*	*s.*	*d.*	C.
1 ...	5 :	13 :	8 ...	108

s.

$$113 \times 108 = \left\{\begin{matrix} 324 \\ 108 \\ 108 \end{matrix}\right.$$

3) 108 (.. 36
36

20) 1227|6
l. *s.* *d.*
Anſw. 613 : 16 : 0

C.

C. *l.* *s.* *d.* *C.*
1 . . . 2 : 10 : 6 . . . 84

---

50 × 84 = 4200.
2) 84 (42
l. *s.* *d.*
20) 424|2 (212 : 2 : 0

C. *l.* *s.* *d.* *C.*
1 . . . 1 : 12 : 4¼ . . . 306

---

s. *d.*
32 × 306 = { 612
918.
3) 306(. . 102
48) 306(. . 6 : 4½

---

20) 990|0 : 4½

l. *s.* *d.*
Anſw. 495 : 0 : 4½

Note, When the given price of an Integer con- ſts of certain Pence together with ½ *d.* or ¾ *d.* it will e convenient to take due *aliquot parts* of the num- er of Integers propounded for all the given price f an Integer, except 1 *d.* and the ſaid ½ *d.* or ¾ *d.* hen for that Peny, and ½ *d.* take ⅛ of the ſaid In- egers propounded, and if there be yet a farthing, ake ⅙ of the ſaid *quotient* which ariſeth by taking ; both which quotients give the value in ſhillings orreſpondent to 1 ¾ *d.* This will be evident by the llowing *Examples.*

yard *d.* *yards*
1 ... $8\frac{3}{4}$ .. 326

s. *d.*
3) 326(.. | 108 .. 8
4) 326(.. | 81 .. 6
8) 326(.. | 40 .. 9
6) 40(.. | 6 .. 8
6) 9(.. | 0 .. $1\frac{1}{2}$

20) 23|7 .. $8\frac{1}{2}$
l. *s.* *d.*
Anſw. 11 : 17 : $8\frac{1}{2}$

s. *d.*
1 .. 3 : $6\frac{1}{2}$ .. 720

s.
3 × 720 = | 2160
4) 720(:: | 180
6) 720(:: | 120
8) 720(:: | 90
l. *s.* *d.*
20) 255|0 (127 : 10 : 0

XVI. When the price of an Integer is given, and the price of many Integers of the ſame name together with $\frac{1}{4}$ or $\frac{1}{2}$ or $\frac{3}{4}$ of an Integer is required, the value of thoſe Integers may be firſt found by ſome of the precedent Rules, and then for the price of $\frac{1}{2}$ of an Integer, take $\frac{1}{2}$ of the given price of an Integer; likewiſe for $\frac{1}{4}$ of an Integer, take of the ſaid given price; alſo for $\frac{3}{4}$ of an Integer take the compoſed of $\frac{1}{2}$ and $\frac{1}{4}$ of the ſaid given price: So if it be demanded what 34 *C.* 3 *qu.* (to wit, 34 hundred weight, and $\frac{3}{4}$ of an hundred weight) of Sugar will coſt at 4 *l.* 16 *s.* 3 *d. per C.* the *Anſwer* will be found 167 *l.* 4 *s.* 8 $\frac{1}{4}$ *d.* as by the ſubſequent operation is manifeſt.

C. *l.* *s.* *d.* *C.* *q.*
1 . . . 4 : 16 : 3 . . . 34 : 3.

s. *d.*

$$96 \times 34 = \begin{cases} 204 \\ 306 \end{cases}$$

4) 34 (. . 8 . . . 6

the Quotients for $\begin{cases} \frac{1}{2}\ C. \\ \frac{1}{4}\ C. \end{cases} \begin{cases} 48 \ldots 1\frac{1}{2} \\ 24 \ldots 0\frac{3}{4} \end{cases}$

20) 334|4 . . . 8 $\frac{1}{4}$.
l. *s.* *d.*
Anſw. 167 . . . 4 . . 8 $\frac{1}{4}$

An Example of *Avoirdupois* greater Weight, where the quantity whoſe Price is ſought conſiſts of entire Hundred weights, quarters of an Hundred, and of ſome number of Pounds, which is not an aliquot Part of 28 or $\frac{1}{4}$ *C.*

C.	l.	s.	d.		C.	q.	lb.
1	5	15	7¾	. . . .	218	3	24

	115 *x* 218=	1090 218 218		
	2) 218 (. .	109	*d.*	*far.*
	8) 218 (. .	27	3	0
	⅙ of 27 *s.* 3 *d.*	4	6	2
The quotients arising for	½ *C.*	57	9	3½
	¼ *C.*	28	10	3¾
	14 *lb.*	14	5	1¾ +
	7 *lb.*	7	2	2¾ +
	3 *lb.*	3	1	0 +
	20)	2532|2	3	2 +
		l.	*s.*	*d.*
	Anſw.	1266	2	3½ +

The example laſt mentioned being (of thoſe queſtions which ordinarily happen in trade) one of the hardeſt to be reſolved by the *Rule of Practice*, I ſhall touch upon the aforegoing operation, where you may obſerve the price of 218 *C*. 3 *qu.* to be found after the manner of former Examples; then the 14 *lb.* part of the 24 *lb*. in the queſtion, I take ½ of the price of ¼ *C.* Likewiſe for 7 *lb.* I take half the price of 14 *lb.* and ſo there yet remains 3 *lb.* whoſe price is found by taking $\frac{3}{7}$ of the price of 7 *lb viz* the price of 7 *lb* being very near 7 *s* 2½ *d.* or 86 ½ *d.* I multiply 86 ½ by 3, and divide the quotient by 7, ſo there ariſeth 37 *d.* or 3 *s.* 1 *d*, very near. Laſtly, all being added together, the ſum is found to be very near 25322 *s.* 3 ½ *d.* or 1266 *l.* 2 *s.* 3½ *d.*

Note, That a quarter of a Farthing (or $\frac{1}{16}$ of a Peny) is the ſmalleſt Money expreſt in the example; and where any thing ariſeth leſs than a quarter of a Farthing, it is omitted, but it is ſuppoſed to follow this Note +, for which ſurpluſages ſome reſpect ought to be had in adding all together: Now albeit, in reſolving queſtions after this practical manner there will be ſome error, yet the loſs for the moſt part will be leſs than a Farthing, which is inconſiderable.

XVII. When the price of 1 or an Integer conſiſts of divers Denominations, as Pounds, Shillings, Pence; and the price of a certain number of Integers, which exceeds not a ſingle figure, is required, work as in the following Example, *viz.* If it be required to find what 8 *C.* muſt coſt at 3 *l.* 13 *s.* 7½ *d. per C.* it is evident that 8 *C.* muſt coſt 8 times 3 *l.* 13 *s.* 7 ½ *d.* therefore I multiply ½ by 8, ſaying, 8 half Pence make 4 Pence, which I reſerve in mind; again, 8 times 7 Pence make 4 *s*, 8 *d.* (to wit, 8 ſix Pences make 4 *s.* and there are 8 Pence beſides) to which adding 4 pence in mind, there will ariſe 5 *s.* which I reſerve in mind, and ſubſcribe a cypher under the place of Pence; again, I ſay 8 times 13 Shillings make 5 *l* 4 *s.* (to wit, 8 Angels make 4 *l.* and 8 times 3 *s.* make 1 *l* 4 *s.*) to which adding 5 *s.* in

C. *l.* *d.* *C.*
1 . . 3 : 13 : 7½ . . 8
8

Anſw. 29 : 9 : 0

in mind, the ſum will be 5 *l.* 9 *s.* wherefore I ſubſcribe 9 *s.* (the exceſs above the pounds) under the ſhillings, and keep 5 *l.* in mind; laſtly, I ſay 8 times 3 pounds make 24 pounds, which with 5 pounds in mind make 29 pounds; ſo that the total product or anſwer of the queſtion is found to be 29 *l.* 9 *s.*

More Examples of this kind are theſe.

C. *l.* *s.* *d.* *C.*
1 . . 17 : 15 : $5\frac{1}{4}$ . . . 7
7

Anſw. 124 : 8 : $0\frac{3}{4}$

C. *l.* *s.* *d.* *C.*
1 . . . 18 : 12 : $6\frac{3}{4}$ . . . 8
8

Anſw. 149 : 00 : 6

XVIII. When the price of 1 *lb.* weight is known, and the price or value of 1 *C.* (to wit 112 *lb.*) is required, the anſwer may ſometimes be given more ſpeedily than by any of the former Rules, by this Rule which follows, *viz.* Find the number of farthings contained in the given price of 1 *lb.* weight, then take twice the number of ſhillings, and once that number of groats, and having added them together, the ſum will give the value of 1 *C.* to wit, 112 *lb.* weight: So if it be demanded what 1 *C.* or 112 *lb.* weight of Cheeſe will coſt at the rate of $3\frac{1}{4}$ pence the pound weight, the anſwer will be 1 *l.* 10 *s.* 4 *d.*

; For according to the ſaid Rule, the number of farthings contained in 3 ¼ d. (the price of 1 pound weight) is 13,

	l.	*s.*	*d.*
therefore the double of 13 ſhillings is..	1	6	0
13 Groats make ..	0	4	4
Therefore the ſum (which is the price of 1 *C.* or 112 *lb.* weight) is . : .	1	10	4

The reaſon of this Rule is evident, for if 1 *lb.* weight coſt 13 farthings, then 112 *lb.* muſt neceſſarily coſt 112 times 13 farthings, or (which is the ſame) 13 times 112 farthings; but 13 times 112 farthings are equal to twice thirteen ſhillings together, with once thirteen groats, becauſe 112 farthings are compoſed of twice 48 farthings (or two ſhillings) and of 16 farthings (or one groat;) wherefore the truth of the ſaid Rule is evident.

Another Example, when Sugar is at 5 ½ *d.* the pound weight, what is the value of 1 *C* (or 112 *lb.* weight?) *Anſw.* 2 *l.* 11 *s.* 4 *d.* For in 5 ½ *d.* are contained 22 *farthings*, therefore

	l.	*s.*	*d.*
the double of 22 *ſhillings* is . ,	2	4	0
22 groats make . .	0	7	4
Which added together give the price of 1 *C.* or 112 *lb.* to wit . .	2	11	4

XIX. When the gain of (or allowance for) 100 Integers conſiſts of ſome number of pounds not exceeding 10, the gain of as many like Integers and known parts of an Integers as one will, may be found very briefly by the following method, *viz.* If 100 *l.* gain 3 *l.* what is the

Compendious ways of computing Intereſt and Factors allowances.

gain of 246 *l.* 18 *s.* 10 *d*) *Anſwer* 7 *l.* 8 *s.* 1 $\frac{98}{100}$ *d.* Firſt, I multiply 246 *l.* 18 *s.* 10 *d.* by 3 (the ſecond term) after the manner delivered in the 17 Rule of this Chapter, and write down the Product which is 740 *l.* 16 *s.* 6 *d.* Then I divide the ſaid Product by 100 (the firſt term in this *Rule of Three*) in this manner, *viz.* I divide 740 pounds by 100, which is performed by cutting off towards the right hand

```
l.     l.     l.    s.   d.
100 . . 3 . . 246 : 18 : 10
                          3
---------------------------
      l. 7|40 : 16 : 06
           |12
      ----------
      s. 8|16
          |12
      ----------
      d. 1|98
```

the two laſt places of 740, ſo the Quotient gives 7 Pounds and there will be a remainder of 40 Pounds, which 40 Pounds I reduce into Shillings, ſo there will ariſe 800 *s.* to which adding the 16 *s.* which ſtand in the place of Shillings, the ſum will be 816 Shillings; theſe are all to be divided by 100 (by cutting off two places as before,) ſo the Quotient will give 8 Shillings, and there will remain 16 Shillings, which being reduced to Pence, and unto them 6 Pence being added (to wit, the 6 Pence which ſtands in the place of Pence) there will ariſe 198 Pence; theſe alſo are to be divided by 100 (by cutting off two places to the right hand as before,) ſo the quotient gives 1 penny, and there will remain 98 pence; ſo the exact quotient or anſwer of the queſtion is found to be 7 *l.* 8 *s.* 1 $\frac{89}{100}$ *d.*

More Examples of this Rule are theſe following.

```
 l.      l.       l.    s.   d.
100 ... 6 ... 793 : 12 : 7
                          6
            ------------------
         l. 47 | 61 : 15 : 6
               | 20
         ------------
         s. 12 | 35
               | 12
         ----------
         d.  4 | 26
```

```
 l.      l.      l.    s.   d.
100 ... 8 ... 43 : 14 : 3
                        8
            ----------------
         l. 3 | 49 : 14 : 0
              | 20
         ----------------
         s. 9 | 94
              | 12
         ------------
        d. 11 | 28
```

After the ſame manner may this following queſtion and ſuch like be reſolved, *viz.* When 100 Ells of Linen cloth coſt 30 *l.* 18 *s.* 9 *d.* what is the price of 1 Ell? *Anſwer* 6 *s.* 2 *d.* 1 *farth.*

Ells. *l.* *s.* *d.* *Ell.*
100 : 30 : 18 : 9 :: 1
20

Shil. 6|18
|12

Pence 2|25
|4

Farth. 1|00

XX. When the given gain of (or allowance for) 100 Integers conſiſts of ſome number of Pounds not exceeding 10, together with ſome Aliquot part or parts of a Pound, the operation will be little different from the laſt mentioned Examples, as may appear by the reſolution of the ſubſequent queſtion, *viz.* What muſt be allowed for 2156 *l.* 13*s.* 4*d.* at the rate of 6 *l.* 15 *s.* for 100 *l.*? *Anſwer* 145 *l.* 11 *s.* 6 *d.* thus found; firſt I multiply the ſaid 2156 *l.* 13 *s.* 4 *d.* by 6 (the number of Pounds in the given allowance 6 *l.* 15 *s.*) after the manner of the laſt examples, and ſubſcribe the Product which is 12940 *l.* underneath the line as you ſee, then ſince 15 *s.* are equal to ½ *l.* together with ¼ *l.* I take ½ of 2156 *l.* 13 *s.* 4 *d.* which is 1078 *l.* 6 *s.* 8 *d.* likewiſe ¼ of the ſaid 2156 *l.* 13 *s.* 4 *d.* to wit, 539 *l.* 3 *s.* 4 *d.* and having ſubſcribed theſe quotients underneath the Product firſt found, and added them all together, I find 14557 *l.* 10 *s.* 0 *d.* for the total product, with which I proceed as in the former Examples, and ſo at length the *Anſwer* is found to be 145 *l.* 11 *s.* 6 *d.* View diligently the operation *l.* . . *l.* . . *l.* : *s.* : *d.*

100 . . $6\frac{3}{4}$ . . 2156 : 13 : 4

$6\frac{3}{4}$

12940 : 00 : 0

1078 : 06 : 8

539 : 03 : 4

l. 145|57 : 10 : 0

|20

s. 11|50

|12

d. 6|00

CHAP. III.

Concerning Exchanges of Coins, Weights, and Meaſures.

1. THE rate or proportion between *Coins*, *Weights*, &c. of different kinds being known, either from ſome good Author, or rather by experience; it will not be difficult, to ſuch as underſtand the *Rule of Three*, to know how to exchange a given quantity of one kind, for a quantity of the ſame value in another kind. But ſince in ſome caſes the common way of working may be much contracted,

tracted, I ſhall endeavour to ſhew the moſt compendious ways to perform this buſineſs.

II. In exchanging of things of different kinds (whether they be *Coins* or *Weights*, &c.) when two things of different kinds are compared together, the queſtion may be reſolved by one ſingle *Rule of Three*, as will be evident by the ſubſequent Examples; *viz.*

Queſt. 1. How many *Riders* at 21 *s.* 2 ½ *d. ſterling* the piece, ought to be received for 251 *l.* 6 *s.* 4 ½ *d.* of *ſterling* Money? *Anſwer*, 237 *Riders.* For the firſt and third terms in the *Rule of Three*, which ariſe from this queſtion, being converted into half pence, the proportion will be thus,

509 . 1 :: 120633 . 237

Queſt. 2. If 100 *Ells* of *Antwerp* make 75 *yards* of *London*, how many yards of *London* meaſure will 27 *Ells* of *Antwerp* make? *Anſwer* $20\frac{1}{4}$ *yards.*

$100 . 75 :: 27 . 20\frac{1}{4}.$

III. When more than two different *Coins*, *Weights*, *Meaſures*, &c. are compared together, *viz.* when one kind of *Coin* is compared with a ſecond of another kind; that ſecond with a third; the third with a fourth; the fourth with a fifth, *&c.* two different caſes are ordinarily raiſed from ſuch compariſon; *viz.*

It may be required to know.
- 1. *How many pieces of the firſt Coin are equal in value to a given number of pieces of the laſt Coin:* Or,
- 2. *How many pieces of the laſt Coin are equal in value to a given number of pieces of the firſt kind of Coin?*

An

An Example of the firſt Caſe.

If 35 ells of *Vienna* make 24 ells at *Lyons*; 3 ells of *Lyons* 5 ells of *Antwerp*; and 100 ells of *Antwerp* 125 ells at *Franckfort*; how many ells of *Vienna* are equal unto 50 ells at *Franckfort*? *Anſwer*, 35 ells of *Vienna*.

For the more eaſie underſtanding of the reſolutions of this queſtion and others of like nature, Let *a* repreſent an ell at *Vienna*; *b* an ell at *Lyons*; *c* an ell at *Antwerp*, and *d* an ell at *Franckfort*; then may the given terms in the queſtion be ſtated in the following order.

$$\begin{array}{rr} \textit{Suppoſitions}\left\{\begin{array}{r} \\ \\ \\ \end{array}\right. & \begin{array}{r} 35\,a = 24\,b \\ 3\,b = 5\,c \\ 100\,c = 125\,d \end{array} \\ \textit{The queſtion} & 50\,d = \;?\,a \end{array}$$

Which order of placing the ſaid given numbers (or terms) being obſerved, it appears that if 35 *a* be accounted to ſtand in the firſt place: 24 *b* in the ſecond; 3 *b* in the third; 5 *c* in the fourth; 100 *c* in the fifth, *&c.* then all the terms which ſtand in odd places, to wit, the firſt, third, fifth, and ſeventh places, will neceſſarily fall under the firſt row or column on the left hand, and all the terms which ſtand in even places, to wit, in the ſecond, fourth, and ſixth places, will fall under the latter column.

Theſe things premiſed, all queſtions which fall under Caſe 1 before-mentioned, may be reſolved by this Rule, *viz.*

Rule I.

Multiply all the given terms which ſtand in odd places (to wit, in the firſt column) according to the Rule of continual Multiplication, and reſerve the laſt Product for a Dividend: Again, multiply continually all the terms which ſtand in even places, ſo ſhall the Product be a Diviſor, and the Quotient ariſing from the ſaid Dividend and Diviſor ſhall be the anſwer of the Queſtion.

So in the laſt mentioned Queſtion, if all the Numbers in the firſt Column, to wit, 35, 3, 100, and 50 be multiplied continually, the Product will be 525000 for a Dividend; alſo if all the Numbers in the latter Column, *viz.* 24, 5 and 125 be multiplied continually, the laſt Product will be 15000 for a Diviſor, and the Quotient ariſing from the ſaid Dividend and Diviſor will be 35, which is the number of Ells of *Vienna* required.

35	24
3	5
100	125
50	

525000 : 15000) 525000 (35

The reaſon of the ſaid Rule I, will be manifeſt by ſolving the queſtion propounded by three ſingle Rules of Three, thus,

$$\text{I. } 24\,b.\ 35\,a :: 3\,b.\ \frac{35 \times 3}{24}\,a\ (= 5\,c.$$

$$\text{II. } \frac{5}{1}\,c.\ \frac{35 \times 3}{24}\,a :: \frac{100}{1}\,c.\ \frac{35 \times 3 \times 100}{5 \times 24}\,a\ (= 125\,d.$$

$$\text{III. } \frac{125}{1}\,d.\ \frac{35 \times 3 \times 100}{5 \times 24}\,a :: \frac{50}{1}\,d.\ \frac{35 \times 3 \times 100 \times 50}{125 \times 5 \times 24}\,a.$$

which fourth proportional laſt found, to wit, $\frac{35 \times 3 \times 100 \times 50}{125 \times 5 \times 24}$ being well viewed and compared with the before mentioned order of placing the terms given in the queſtion gives the very Rule I. before expreſt in words.

An Example of the latter of the two Caſes before mentioned

If 10 *lb.* of *Avoirdupois* weight at *London* be equal to 9 *lb.* of *Amſterdam*; 45 *lb.* at *Amſterdam.* 49 *lb.* at *Bruges*; and 98 *lb.* at *Bruges* equal to 116 *lb.* at *Dantzick*; how many *lb.* of *Dantzick* are equal to 112 *lb.* of *Avoirdupois* weight at *London?* *Anſwer*, 129. 92 *lb.* of *Dantzick.*

That the operation may be the more clear, let *a* repreſent one pound of *Avoirdupois* weight; *b* one *lb.* of *Amſterdam*; *c* one *lb.* of *Bruges*, and *d.* one *lb.* of *Dantzick*; then let the queſtion be ſtated after the order in the firſt Caſe, *viz.*

Suppoſitions

$$\textit{Suppoſitions}\left\{\begin{array}{l} 10\,a = 9\,b \\ 45\,b = 49\,c \\ 98\,c = 116\,d \end{array}\right.$$

$$\textit{The queſtion}\ 112\,a = \ ?\,d$$

Theſe things premiſed, all queſtions which fall under Caſe 2. before mentioned, may be reſolved by this Rule, *viz.*

Rule II.

Multiply all the given terms which ſtand in even place (to wit in the latter column) and the laſt odd term in the firſt column according to the Rule of continual Multiplication, and reſerve the laſt product for a Dividend; again, multiply continually the reſt of the terms which ſtand in odd places (to-wit in the firſt column) for a Diviſor, ſo ſhall the quotient ariſing be the anſwer of the queſtion.

Or in this latter caſe if you place the laſt of the given terms in the ſame column with the even terms, the rule for ſolving queſtions, which fall under the latter caſe will be this which followeth, *viz.*

Multiply continually all the numbers in the latter column for a Dividend; alſo multiply continually all the numbers in the firſt column for a Diviſor, ſo ſhall the quotient ariſing be the anſwer of the queſtion. Thus the anſwer of the laſt mentioned queſtion will be found 129. 92, to wit, 192 $\frac{92}{100}$ *lb.* of *Dantzick*, as is evident by the ſubſequent operation.

$$\begin{array}{r|l} 10 & 9 \\ 45 & 49 \\ 98 & 116 \\ & 112 \\ \hline \end{array}$$

$$44100 \quad 5729472\,(129.92$$

The Reaſon of the ſaid Rule II. will be manifeſt by ſolving the queſtion propounded, by three ſingle Rules of three, thus,

$$I\; 9\,b.\; 10\,a :: 45\,b\; ..\; \frac{45\times10}{9}\,a\,(= 49\,c.$$

$$II.\; \frac{49}{1}c\; \frac{45\times10}{9}a :: \frac{98}{1}c\; \frac{45\times10\times98}{49\times9}a\,(=116\,d.$$

$$III.\; \frac{45\times10\times98}{49\times9}a.\; \frac{116}{1}d.\; \frac{112}{1}a.\; \frac{49\times9\times116\times112}{45\times10\times98}d.$$

vhich fourth proportional laſt found, to wit, $\frac{9\times9\times116\times112}{5\times10\times98}$ being well viewed and compaed with the forementioned order of placing the erms given in the queſtion diſcovers the very Rule I. before expreſt in words.

Note, When the ſame Numbers happen to be Iultiplicators in the Dividend, and alſo in the Diviſor, ſuch Multiplicators may be cancelled in oth, and thereby much labour will oftentimes be pared.

Such which have much practice in calculating *Exchanges*, and do exactly know the rate or proportion between two different weights or meaſures or coins, which they would compare together, may by the *Rule of Three* frame Tables of proportions for the more ſpeedy reducing of a given quantity of one kind of Weight, Meaſure, *&c.* into a quantity of the ſame value in another kind of weight, *&c.* In the expreſſing of which proportions it will be very convenient that the firſt Number or Antecedent of each Proportion be made 1 or unity, and the ſecond term or conſequent a Decimal, or elſe a mixt number whoſe Fractional part is a Decimal, for then the Coin, Weight, *&c.* of the one place (whoſe term is 1) may be reduced into that of the other place, by help of thoſe Tables and of Multiplication of Decimals without ſenſible error: For Example, It hath been obſerved by ſome ingenious Merchants that 100 *lb* of *Avoirdupois* weight at *London*, are equal unto 89 *lb* in *Paris* by the King's Beam, and conſequently 1 *lb* *Avoirdupois* is equal to $\frac{89}{100}$ *lb* or 89 *lb* at *Paris*; (for if 100 give 89, then 1 will give 89;) therefore any number of Pounds *Avoirdupois* being multiplied by 89 (with reſpect unto Multiplication of Decimals, explained in the 26 Chapter of the preceeding Book) will produce Pounds of *Paris*: Again, if 89 *lb* of *Paris* be equal to 100 *lb* *Avoirdupois*, then 1 *lb* of *Paris* will be near equal to 1.1235 *lb* of *Avoirdupois*; therefore any number of Pounds of *Paris* being multiplied by 1.1235 will produce Pounds *Avoirdupois* very near.

Upon this ground I have collected the proportions in the following Tables, wherein I would not have any to confide further than they ſhall know them to be agreeable to truth, for I have only derived them from thoſe delivered by Mr. *Lewis Roberts*, Merchant, in his *Map of Commerce*, and do herein only aim at the Inſtruction of ingenious *Merchants* and *Factors* in the briefeſt ways of calculating their exchanges, the rate or proportion being truly known; in which Practice, *Decimal Arithmetick* (which hath no Enemy but the Ignorant) will be very ſerviceable.

A Table for the Reduction of Avoirdupois *Weight at* London, *to the Weights of divers Foreign Cities and remarkable Places.*

		lb.
One Pound of *Avoirdupois* weight at *London*, makes at	*Antwerp*	.9615
	Amſterdam	.9
	Abeville	.91
	Acona	1 .282
	Avignon	1 .12
	Burdeaux	.91
	Burgoigne	.91
	Bollonia	1 .25
	Bruges	.98
	Callabria	1 .3698
	Callais	1 .07
	Conſtantinople	.8474 *Loder*;
	Deep	.91
	Dantzick	1 .16
	Ferrara	1 .3333
	Florence	1 .282
	Flanders in general	1 .06
	Geneva	.9345

		lb.
One Pound of *Avoirdupois* weight at *London,* makes at	*Genova*	1 .4084 *ſuttle*
		1 .4285 *groſs*
	Hamburg	.92
	Holland	.95
	Lisbon	.881
	Lyons	1 .07 *common weight.*
		.98 *ſilk weight.*
		.9 *cuſtomers weight.*
	Leghorn	1 .3333
	Millan	1 .4285
	Mirandola	1 .3333
	Norimberg	.88
	Naples	1 .4084
	Paris	.89
	Prague	.83
	Placentia	1 .3888
	Rotchel	1 .12
	Rome	1 .27
	Rovan	.875 *by vicont.*
		.9017 *com. weight.*
	Sevil	1 .08
	Tholouſa	1 .12
	Turin	1 .2195
	Venetia	1 .5625 *ſubtil*
		.9433 *groſs*
	Vienna	.813

The uſe of the precedent Table will be manifeſt by the ſubſequent *Example, viz.*

How much weight at *Dantzick* do 320 *lb. Avoirdupois* make? *Anſwer*, 371 .2 *lb.* Seek in the precedent Table for *Dantzick*, and right againſt it you ſhall find 1 ,16. which ſhew that 1 *lb Avoirdupois* is equal to 1 .16 *lb.* at *Dantzick*, therefore multiply 320 by 1 .16, ſo will the product be 371 .2 *lb.* of *Dantzick*, as by the Operation is manifeſt.

Avoir. Dantz. Avoir. Dantz.
1 : 1 , 16 :: 320 : 372.2
1.16

1920
320
320

371|20

A

A Table for the Reduction of the Weights of divers Foreign Cities and remarkable Places to Avoirdupois *Weight at* London.

			lb.
One Pound weight in	*Antwerp*	makes at *London* of *Avoirdupois* weight	1 .04
	Amſterdam		1 .1111
	Abbeville		1 .0989
	Ancona		.78
	Avignon		.8928
	Burdeaux		1 .0989
	Burgoyne		1 .0989
	Bollonia		.8
	Bruges		1 .0204
	Callabria		.73
	Callais		.9345
	Deep		1 .0989
	Dantzick		.862
	Ferrara		.75
	Florence		.78
	Flanders in general		.9433
	Geneva		1. 07
	Genoa ſubtil		.71
	Genoa groſs,		.70

One Pound weight in		makes at London of *Avoirdupois* weight lb.
Hamburg		1 0865
Holland		1 .0526
Lisbon		1 .135
Lyons	common weight.	.9345
	ſilk weight.	1 .0204
	cuſtom weight.	1 .1111
Leghorn		.75
Millan		.7
Mirandola		.75
Norimberg		1 .1363
Naples		.71
Paris		1 .1235
Prague		1 .2048
Placentia		.72
Rotchel		.8928
Rome		.7874
Ronan	by Vicont.	1 .1428
	com. weight.	1 .1089
Sevil.		.9259
Tholouſa		.8928
Turin		.82
Venetia	ſuttle,	.64
	groſs,	1 .06
Vienna		1 .23

The uſe of the laſt mentioned Table, will be manifeſt by this example, *viz.*

In 224 *lb.* weight at *Hamburg*, how many pounds *Avoirdupois*?

Anſw. 243.376 *lb.*

Seek in the Table for *Hamburg*, and right againſt it you will find 1.0865, which ſheweth that 1 *lb.* of *Hamburg* makes 1.0865 *lb. Avoirdupois*, therefore if 1.0865 be multiplied by 224 the product will be pounds *Avoirdupois.*

```
1 . . . 1.0865 . . . 224
           224
      ----------
         43460
        21730
       21730
      ----------
       243|3760
```

A Table for the Reduction of Engliſh *Ells to the Meaſures of divers Foreign Cities, and remarkable Places.*

One Ell at *London* makes at	*Amſterdam*		1 .6949	Ells
	Antwerp		1 .6666	
	Bruges		1 .64	
	Arras		1 .65	
	Norimberg		1 .74	
	Colen		2 .08	
	Liſle		1 .66	
	Maſtrich		1 .57	
	Frankfort		2 .0866	
	Dantzick		1 .3833	
	Vienna		1 .45	
	Paris		.95	
	Rouan		1 .03	Aulnes
	Lions		1. 0166	
	Callais		1 .57	
	Venice	linen	1 .8	Braces
		ſilk.	1 .96	
	Lucca		2	
	Florence		2 .04	
	Milan		2 .3	
	Leghorn		2	
	Madera Iſles		1 .0328	

One Ell at *London* makes at	*Sevil*	1 .35	Vares
	Lisbon	1.	
	Caſtilia	1 .3875	
	Andoluzia	1 .3625	
	Granado	1 .3625	
	Genoa	4 .8083	Palmes
	Saragoſa	.55	Canes
	Rome	.56	
	Barſelona	.7125	
	Valentia	1 .2125	

The uſe of the aforeſaid Table will be manifeſt by the ſubſequent Example, *viz.*

In 325 ells of *London*, how many ells at *Antwerp?* *Anſw.* 541 .645 ells: Seek in the Table for *Antwerp*, and right againſt it you ſhall find 1 .6666 which being multiplied by 325 produceth 541 .645 ells of *Antwerp*, as by the operation is manifeſt.

1 . . . 1. 6666 . . . 325
325

83330
33332
49998

541|6450

A Table for the Reduction of the Meaſures of divers Foreign Cities and remarkable Places to Engliſh *Ells.*

One Ell at	*Amſterdam*		makes at *London*	.59
	Antwerp			.6
	Bruges			.6097
	Arras			.606
	Morimberg			.5747
	Colen			.4807
	Liſle			.6024
	Maſtrich			.6369
	Franckfort			.4792
	Dantzick			.7228
	Vienna			.6896
One Auln at	*Paris*			1 .0526
	Rouan			.9708
	Lions			.9836
	Callais			.6369
One Brace at	*Venice*	linen		.5555
		ſilk,		.5102
	Lucques			.5
	Florence			.4901
	Milan			.4347
	Legborn			.5
	Madera Iſles			.9681

Sevil

One Cane at One Vare at	*Sevil*	makes at *London*	.7407	Ells
	Lisbon		1.	
	Caſtilia		.7407	
	Andoluzia		.7339	
	Granado		.7339	
	One Palm at *Genoa*		.2079	
	Saragoſa		1 .8181	
	Rome		1 .7875	
	Barſelona		1 .4035	
	Valentia		.8247	

The uſe of the ſaid Table will be manifeſt by the ſubſequent Example, *viz.*

In 730 Aulneſs at *Lions*, how many Ells at *London.*

Anſw. 718 .028. Seek in the Table for *Lions*, and right againſt it you ſhall find .9836, which being multiplied by 730 produceth 718.028 Ells of *London*, as by the operation is manifeſt.

1 . . . 9836 . . . 730
730

295080
68852

71810280

Note, That one and the ſame kind of Weight or Meaſure doth ſeldom or never alter from its peculiar quantity, in the Kingdom or Common-wealth, where ſuch Weight or Meaſure was firſt eſtabliſhed; but one and the ſame kind of Money doth often riſe and fall in its value in foreign Parts: For which cauſe I have ſpared the pains of calculating *Decimal Tables* for *Coins*, yet to give ſome light to ſuch as read modern Relations, and want experimental knowledge in this matter, I ſhall here inſert a *Table* in the ſame eſtate as I find it in the aforeſaid *Map* of *Commerce*, and refer the Reader for further ſatisfaction, to the *Table* in *Rider*'s *Dictionary*, concerning *Coins*, *Weights* and *Meaſures*, both Ancient and Modern.

Of Exchanges of London, *with divers foreign Cities.*

London doth exchange with

	Pence			
Placentia ſterl.	64	for	1	*Crown*
Lyons	64	for	1	*Crown*
Rome	66	for	1	*Ducat*
Genoa	65	for	1	*Crown*
Milan	64 ¾	for	1	*Crown*
Venice	50	for	1	*Ducat*
Lucca	53 ½	for	1	*Duccaton*
Naples	50	for	1	*Ducat*
Lecchia in *Calabria*	50 ½	for	1	*Ducat*
Barri	51	for	1	*Ducat*
Palermo	57 ½	for	1	*Ducat*
Meſina	56 ½	for	1	*Ducat*
Antwerp & *Colen* 1 *l. ſterl.*		for	34 ½	*ſhill. ſlem.*
Valentia	57 ½	for	1	*Ducat*
Saragoſa	59	for	1	*Ducat*
Barſelona	64	for	1	*Ducat*
Lisbon	53 ½	for	1	*Ducat*
Bollonia	53 ⅓	for	1	*Ducaton*
Bergamo	52	for	1	*Ducaton*
Frankfort	59 ½	for	1	*Florin*
Genoa	65	for	1	*Crown*

London exchangeth in the denomination of *Pence ſterling* with all other Countries, *Antwerp* and thoſe neighbouring Countries of *Flanders* and *Holland* excepted, with which it exchangeth by the entire *Pound* of 20 *Shillings Engliſh* (or *ſterling.*)

CHAP. IV.

Practical Queſtions about various Things: viz. *Tare, Tret, Loſs, Gain, Barter, Factorſhip, and Meaſuring of Tapeſtry.*

Of Abatements and Allowances in Traffick, viz. 1. *Of Tare.*

IN the Trade of Merchandize there are in uſe various Allowances, and Abatements, known by the names of *Tare*, *Tret*, &c. concerning which I ſhall give a few examples, whereby the practical *Arithmetician* will eaſily ſee, that there is more difficulty in the name than in the thing; for the rate, or proportion agreed upon, in any Allowance or Abatement (be it called by what name ſoever) being once known, the Arithmetical work will quickly be diſpatch'd by the *Rule of Three*, or elſe by that and ſome of the former rules mixtly uſed, as will partly appear by the following queſtions.

Groſs weight is compoſed of the Neat weight of the Commodity, and alſo the Tare, to wit, the Cheſt, Bag, But, &c. which containeth the Commodity.

Queſt. 1. A Factor buyeth 4 Cheſts of Sugar marked A. B. C. D. The groſs weight of each Cheſt in *Avoirdupois* greater weight is as followeth.

	C.	q.	lb.
A.	11	1	19
B.	10	3	20
C.	11	2	13
D.	10	1	17
The total groſs weight	44	1	13.

Now ſuppoſing the *Tare* or weight of each Cheſt, when it is empty, to be 37 *lb.* the queſtion is what neat weight of Sugar will remain, when the total *Tare* is ſubtracted? *Anſw.* 43 C. 0 q. 4 *lb.*

	C.	q.	lb.
from	44	1	13
Subtr.	1	1	08
Rem.	43	0	05 *the neat weight of Sug.*

Queſt. 2. If from 990 *C.* 3. *qu.* 21 *lb. groſs weight*, *Tare* is to be ſubtracted after the rate of 14 *lb. per C.* (or 112 *lb.*) of *groſs weight*, how many C. neat will remain? *Anſw.* 867 *C.* 0 *q.* 7 $\frac{7}{8}$ *lb.*

I. The groſs weight being converted into pounds by the ſixth Rule of the 7th Chapter of the preceding Book, will give 110985 *lb.*

II. Then by the *Rule of Three*

112 . 14 :: 110985 . 13873 $\frac{1}{8}$
or 8 . 1 :: 110985 . 13873 $\frac{1}{8}$

lb.
III. From 110985 *the groſs weight:*
Subtr. $13873\frac{1}{8}$ *the total Tare.*

	C.	qu.	lb.
Reſt neat $97111\frac{7}{8}$ =	867	0	$7\frac{7}{8}$

Note, When the number of *lb.* to be abated *per C.* for *Tare* is an *aliquot part* of 112, as in the laſt mentioned Example, where $14 = \frac{1}{8}$ of 112, the operation may be thus:

$$\begin{array}{ccccccccc} C. & C. & & C. & q. & lb. & C. & qu. & lb. \\ 1 & \frac{1}{8} & :: & 990 & : 3 & : 21 & : (123 & : 3 & : 13\frac{1}{8} \end{array}$$

$$\frac{1}{8}\text{ of}\left\{\begin{array}{ll} 990\ c. = & 123 : 3 : 00 \\ 3\ q. = & 00 : 0 : 10\frac{4}{8} \\ 21\ lb. = & 00 : 0 : 02\frac{5}{8} \end{array}\right.$$

Total Tare $123 : 3 : 13\frac{1}{8}$
Reſt neat $867 : 0 : 07\frac{7}{8}$

Of Tret.

Queſt. 3. Suppoſe at ſome City, there is a Cuſtom in ſelling of certain Merchandice by weight, to allow or caſt in as an Overplus to the Buyer, 4 *lb.* weight for every 100 *lb.* weight that is bought, and in that proportion for a greater or leſſer quantity. Now if a Merchant buy 1175 *lb.* weight of ſome commodity, and is to be allowed thereupon after the aforeſaid rate, the queſtion is, how many *lb.* weight ought he to receive in all? *Anſw.* 1222 *lb.* weight.

100. 104 : : 1175 . 1222

This kind of allowance is commonly called *Tret.*

Queſt. 4. Suppoſe a Merchant hath 1222 *lb.* weight of a certain commodity, part whereof he bought at a certain rate *per lb.* and the reſt was al- owed to him or caſt in as an Overplus, after the rate of 4 *lb.* weight for every 100 *lb* weight which he bought; the queſtion is, to know how many pounds neat weight he bought? *Anſw.* 1175 *lb.* weight.

104. 100 : : 1222. 1175.

This queſtion is the converſe of the former, and heweth how to make abatement for *Tret.*

Queſt. 5. If from 55 *C.* 1 *qu.* of groſs weight *Tare* is to be ſubtracted after the rate of 16 *lb. per C.* nd from the remainder *Tret* is to be abated after the ate of 4 *lb. per* 104 *lb.* the Queſtion is, what the eat weight is worth in money after the rate of *l.* 8 *s.* for every *C.* (or 112 *lb.*) *Anſw.* 382 ½ *l.*

I. The groſs weight in *lb.* is 6188 *l.*
II. 112 . 16 : : 6188 . 884
or 7 . 1 : : 6188 . 884
III. 6188—884=5304
IV. 104 . 100 : : 5304 . 5100
V. 112 . 8⅖ : : 5100 . 382½

Of Loſs and Gain.

Queſt. 6. A Merchant hath bought inen Cloth at 11 *s. per* Ell, which pro- ng worſe than he expected, he is wil- ng to ſell it at ſuch a price that he may loſe pre- ſely after the rate of 1⅔ *l.* for every 20 *l.* that he id out; the queſtion is to know at what price he ght to ſell the ell, that the proportion in the ſaid loſs may be obſerved? *Anſw.* 10 *s.* 1 *d.* *per* ell.

I. $20 - 1\frac{2}{3} = 18\frac{1}{3}$
II. $20 \,.\, 18\frac{1}{3} :: 11 \,.\, 10\frac{1}{12}$ *pence*

Otherwiſe,

I. $20 \,.\, 1\frac{2}{3} :: 11 \,.\, \frac{11}{12}$
II. $11 - \frac{11}{12} = 10\frac{1}{12}$

Queſt. 7. If 100 *lb.* weight of any commodity coſt 30 *s.* at what price muſt 1 *lb.* weight of that commodity be ſold to gain after the rate of 10 *l.* for every 100 laid out? *Anſw.* $3\frac{24}{25}$ *d. per lb.* weight.

I. $100 \,.\, 110 :: 30 \,.\, 33$
II. $100 \,.\, 33 :: 1 \,.\, \frac{33}{100}$ *s.* (or $3\frac{24}{25}$ *d.*

Queſt. 8. A Merchant ſelleth a parcel of Jewels which coſt him 250 *l.* ready money, for 559 *l.* payable at the end of 6 months; the queſtion is (his ſecurity being ſuppoſed to be good) what his gain was worth in ready money upon rebate of intereſt at the rate of 6 *l.* for 100 *l.* for year? *Anſw.* 300 *l.*

$559 - 250 = 309$
$103 \,.\, 100 :: 309 \,.\, 300$

Of Barter. *Queſt.* 9. How much Sugar at 8 *d. per lb.* weight may be bought for 20 C. of Tobacco at 3 *l. per* C.? *Anſw.* 1800 *lb.* weight of Sugar.

$$1 \,.\, 3 :: 20 \,.\, 60$$
$$\tfrac{1}{30} \,.\, 1 :: 60 \,.\, 1800$$

Queſt. 10. A. hath 100 pieces of Silk, which re worth but 3 *l. per* piece in ready money, yet he arters them with B. at 4 *lb. per* piece, and at that ate takes their value of B. in Wools at 7 *l.* 10 *s. per* . which are worth but 6 *l. per C.* in ready money, he queſtion is to know what quantity of Wool pays or the Silk, and which of the two A. or B. is the ainer, and how much? *Anſw.* $53\frac{1}{3}$ C. of Wool pays or the Silk, and A. gaineth 20 *l.* by the barter.

I. $7\frac{1}{2} \,.\, 1 :: 400 \,.\, 53\frac{1}{3}$

II. $\begin{cases} 1 \,.\, 6 :: 53\frac{1}{3} \,.\, 320 \\ \text{or } 7\frac{1}{2} \,.\, 6 :: 400 \,.\, 320 \end{cases}$

So it is evident that the true worth of the Wool hich B. delivered was 320 *l.* for which he receive nly of A. the worth of 300 *l.* in Silks, and therere B. loſeth 20 *l.* by the barter.

Queſt. 11. A Merchant delivered to his Factor oo *l.* upon condition that if the Fa-or add to it 250 *l.* of his own mo-ey, and beſtow his pains in mana-ng the whole ſtock, he ſhall then ave $\frac{2}{5}$ parts of the total gain. The ueſtion is to know what ſtock the actor's ſervice was eſtimated at? *nſw.* 150 *l.*

Of Factorſhip. See brief rules for computing of Factors allowances in the 19, *and* 20 *Rules of the ſecond Chap. of this Appendix.*

I. The Factor's part of the gain being $\frac{2}{5}$ the Mer-ant muſt neceſſarily have the remainder, which $\frac{3}{5}$.

II. $\frac{3}{5} \,.\, \frac{2}{5} :: 600 \,.\, 400$

III, $400 - 250 = 150$

Queſt. 12. A Merchant delivereth to his Facto 320 *l.* and permitteth him to add to it 64 *l.* of hi own money, to be employed in traffick, and b agreement between them the Factor's ſervice i eſtimated equivalent to a certain ſtock; which i ſuch, that if the total gain be divided proportionably according to thoſe three Stocks, the Factor i to receive $\frac{1}{5}$ of the total gain, in conſideration o the ſaid imaginary ſtock (being the value of hi ſervice;) the queſtion is to know the full part o the gain belonging to each, and what ſtock th Factor's ſervice was valued at? *Anſw.* The Merchant $\frac{2}{3}$ of the gain, and the Factor $\frac{1}{3}$, whoſe ſervice was valued at 96 *l.* ſtock.

I. $320 + 64 = 384$

II. $\frac{4}{5} . \frac{1}{5} :: 384 . 96$

$$\text{III.} \left\{\begin{array}{l}320\\ 64\\ 96\\ \hline 480 . 1 ::\end{array}\right. \qquad \left\{\begin{array}{l}320 . \frac{2}{3}\\ \\ 160 . \frac{1}{3}\end{array}\right.$$

Of Meaſuring of Tapeſtry.

Queſt. 13. If a piece of Arras Hangings, in th form of a long ſquare, hath for i length 6 $\frac{1}{4}$ yards *Engliſh*, and breadt 4 yards; how many ſquare ells, o ſticks *Flemiſh* are contained in that piece, when th length of a *Flemiſh* ell is equal to $\frac{3}{4}$ yard *Engliſh* *Anſwer.* 44 $\frac{4}{9}$ ſquare ells or ſticks *Flemiſh*.

Foraſmuch as by ſuppoſition, a *Flemiſh* ell i length, hath ſuch proportion to an *Engliſh* yard i length, as 3 to 4, and conſequently the ſquare o the one to the ſquare of the other, as 9 to 1

Therefor

Therefore in a direct proportion, as 9 is to 16; ſo s any given number of ſquare yards *Engliſh* to a number of ſquare ells *Flemiſh*, which will take up equal ſpace with the ſaid ſquare ells *Engliſh*. Alſo n a direct proportion, as 16 is to 9, ſo is any given number of ſquare ells *Flemiſh* to a number of ſquare yards *Engliſh*, which will take up an equal ſpace with the ſaid *Flemiſh* ells: Therefore to reſolve the aforeſaid queſtion, firſt find the number of ſquare yards *Engliſh* contained in the ſaid piece of Arras, by multiplying the length and breadth in yards mutually one by the other, then proceed according to the aforeſaid proportion; ſo the work will ſtand thus,

I. $6\frac{1}{4} \times 4 = 25$ ſquare yards *Engliſh*.
II. $9 \,.\, 16 :: 25 \,.\, 44\frac{4}{9}$ ſquare ells *Flemiſh*.

Otherwiſe.

$6\frac{1}{4}$ yards *Engliſh* in length give by the *Rule of Three* in **Flemiſh** ells——	$8\frac{1}{3}$ *length.*
Alſo 4 yards **Engliſh** give in *Flemiſh* ells—— —— —— ——	$5\frac{1}{3}$ *breadth.*
Therefore the product of the ſaid $8\frac{1}{3}$ multipyed by $5\frac{1}{3}$, gives for the ſuperficial content as before—— ——	——— ——— $44\frac{4}{9}$

Queſt. 4 If a piece of Tapeſtry in the form of a long ſquare be in length $15\frac{1}{4}$ ells *Flemiſh*, and in breadth $4\frac{1}{3}$ ells *Flemiſh*, how many ſquare yards *Engliſh* are contained in that piece, when 4 ells *Flemiſh* in length are equal to 3 yards *Engliſh*? *Anſw.* $37\frac{11}{64}$ ſquare yards *Engliſh*.

I. $15\frac{1}{4} \times 4\frac{1}{3} = 66\frac{1}{12}$.
II. $16 \,.\, 9 :: 66\frac{1}{12} \,.\, 37\frac{11}{64}$.

CHAP. V.

Concerning the Intereſt of Money, and the Conſtruction of Tables to that purpoſe.

I. IN reſolving queſtions concerning intereſt of money, four things are to be well obſerved, to wit, Firſt, the Principal, or money lent for gain or intereſt; Secondly, the time for which the ſaid Principal is lent; Thirdly, the rate or proportion which the Principal bears to the ſum of the Principal and Intereſt; and Fourthly, the Intereſt it ſelf: So if 100 *l.* be lent upon condition that 106 *l.* ſhall be repaid at the end of a year, the ſaid 100 *l.* is called Principal; the time for which the ſaid principal is lent is one year; the proportion which the principal bears to the ſum of the principal and intereſt is ſuch as 100 hath to 106; Laſtly, the intereſt it ſelf is 6 *l.*

II. Intereſt is either Simple or Compound.

III. Simple Intereſt is that which ariſeth or is computed from the principal only: So if 100 *l.* be lent for two years, the ſimple Intereſt thereof after the rate of 6 pounds for 100 pounds for 1 year will be 12 pounds, *viz.* 6 pounds due at the firſt years end, and 6 pounds due at the ſecond years end.

IV. Compound Intereſt is that which ariſeth from the Principal, and alſo from the intereſt thereof, and therefore it is called intereſt upon intereſt: So if 100 pounds be lent and forborn 3 years, and compound intereſt thereof is to be computed after the rate of 6 pounds for 100 *l.* for one year; there will ariſe beſides the ſimple intereſt of the principal for three years, the intereſt of 6 pounds (due at the firſt years end) for 2 years, and the intereſt of 6 pounds (due at the ſecond years end) for one year following.

V. Rebate or diſcompt of money is, when a ſum of money due at any time to come, is ſatisfied by the payment of ſo much preſent money, which if it were put forth at a certain rate of intereſt for the ſaid time, would become equal to the ſum firſt due: So if 100 pounds be due at the end of two years, and is to be ſatisfied by the payment of preſent money upon rebate, after the rate of 6 pounds *per centum, per annum, ſimple intereſt,* there ought to be ſo much ready money paid, which in two years after the ſaid rate of intereſt would be augmented unto 100 *l.* In like manner if the rebate or diſcompt were to be made after any rate of compound intereſt, ſo much ready money ought to be paid, which at ſuch rate of compound intereſt, for the time agreed on, would become equal to the ſum firſt due. *Examples* of the manner of computation by rebate may be ſeen in the tenth and fourteenth Rules of this Chapter.

VI. In the taking of intereſt, or uſe money, for the loan or forbearance of money lent, reſpect muſt be had to the rate limited by Act of Parliament, which now reſtraineth all perſons from taking more than 6 *l.* for the intereſt or uſe of 100 *l.* lent for a year, but what part of 6 *l.* may be taken for the intereſt of 100 *l.* lent for half a year, a quarter of

The foundation upon which the rules for computing ſimple intereſt are grounded.

of a year, a month, or any other part of a year, is not expreſt in the Act; In this caſe therefore we muſt obſerve cuſtom and daily practice, ſo we ſhall find that 3 *l.* is uſually taken for half a years Intereſt of 100 *l.* and 30 *s.* for a quarter of a year, *&c.* by which practice, this following Analogy (which is the ground or reaſon of the common Rules for computing ſimple Intereſt) ſeems to be aſſumed for a ſafe Expoſition of the Statute, *viz.* That ſuch proportion as the whole year (ſuppoſed to conſiſt of 365 days) hath to any propounded ſpace of time more or leſs than a year, ſuch proportion any Intereſt (not exceeding the rate limited by the Act) for any Principal lent for a year, ought to have to the Intereſt of the ſame Principal for the time propounded: This Analogy being granted, the manner of computing ſimple Intereſt, for any Principal lent and forborn any time propounded, will be ſuch as is expreſt in the two next Sections.

VII. The Intereſt or Gain of 100 *l.* principal Money forborn for a year being known, the Intereſt of any other principal Money for the ſame time may be found out by one ſingle *Rule of Three*; for as 100 *l.* principal is in proportion to the Intereſt thereof, ſo is any other principal to its Intereſt: So if it be demanded what 270 *l.* will gain in a year at the rate of 6 *l.* for 100 *l.* for one year, the *Anſwer* will be found to be 16 *l.* 4 *s.* For,

l. *l.* *l.* *l.* *l.* *s.* *d.*
100 : 6 : : 270. 16,2 (or 16 : 4 : 0

A ſecond Example. What is the Intereſt of 175 *l.* 18 *s.* 11 *d.* for a year, at the rate of 6 *l.* for 100 *l.*

for a year? *Anſw.* 10 *l.* 11 *s.* 1 $\frac{62}{100}$ *d.* as by the following operation (which is performed after the practical manner delivered in the nineteenth Rule of the ſecond Chapter of this *Appendix*) is evident.

```
 l.    l.     l.    s.    d.    l.   s.   d.
100 . 6 : : 175 : 18 : 11 ( 10 : 11 : 1 62/100
            multiply by . . 6
        ——————————————————————
 l. . . . . . . 10|55 : 13 : 6
                  |20
      ———————————————————
 s. . . . . . . 11|13
                  |12
     ———————————————————
 d. . . . .       1|62
```

VIII If the Intereſt of 100 *l.* principal for one whole year, or 365 days be known, the ſimple intereſt of any other principal, for any number of days more or leſs than 365, may be found out by the following Rule, *viz.*

Multiply theſe three numbers according to the Rule of continual Multiplication, to wit, the given intereſt of 100 *l.* for a year, the principal, whoſe intereſt is required, and the number of days preſcribed, reſerving the laſt product for a Dividend: Alſo multiply 365 by 100, and reſerve this product for a Diviſor; Laſtly finiſh Diviſion, ſo ſhall the quotient be the intereſt or gain ſought.

A Rule for computing ſimple intereſt for any number of days.

Note here, that the two principals, to wit 100 *l.* and the other propounded, are ſuppoſed to be of one and the ſame denomination: Alſo the intereſt required will be of the ſame denomination with the given intereſt of 100 *l.*

For an example of this Rule, let it be required to find out the intereſt of 400 *l.* for a week, or 7 days at the rate of 6 *l.* for 100 *l* for a year, or 365 days: Firſt multiplying theſe three numbers 6, 400, and 7 continually (*viz.* multiplying 6 by 400, and the product thence ariſing by 7) the laſt product will be 16800 for a Dividend; alſo multiplying 365 by 100, the product is 36500 for a Diviſor; Laſtly, dividing 16800 by 36500 (after cyphers at pleaſure are added to 16800) the quotient (according to the fourth Rule of the 27th Chapter of the preceeding Book) will be diſcovered to be this decimal .4602, which is equal to 9 *s.* 2 *d.* 1 *farth.* (as will appear by the brief way of valuing a decimal fraction in the fourth Rule of the 26th Chapter.

The reaſon of the above mentioned rule for the computing of intereſt for days, will be manifeſt by this following way of ſolving the ſame queſtion by two ſingle *Rules of Three*, viz.

$$\text{I. } 100 \;.\; 6 :: 400 \;.\; \frac{6 \times 400}{100}$$

$$\text{II. } \frac{365}{1} : \frac{6 \times 400}{100} :: \frac{7}{1} : \frac{6 \times 400 \times 7}{365 \times 100}$$

Which fourth proportional in the latter *Rule of Three*, to wit, $\frac{6 \times 400 \times 7}{365 \times 100}$ being well viewed the truth of the rule before delivered will be manifeſt.

Hence one vulgar errour in computing intereſt is diſcovered, for ſome argue thus, 6 *l.* is the intereſt of 100 *l.* for a year, therefore 10 *s.* (or $\frac{1}{12}$ of 6 *l.*) is the intereſt for a month, and conſequently 2 *s.* 6 *d.* for a week or ſeven days, and ſo the intereſt of 400 *l.* for 7 days, computed after that manner would be 10 *s.* which exceeds the anſwer found by the preceeding Rule by $9\frac{3}{4}$ *d.* very near, which fallacy hath its riſe from the taking, (or rather miſtaking) of 28 days for $\frac{1}{12}$ part of the number of days in a year, when indeed the juſt $\frac{1}{12}$ part of 365 days conſiſts of $30\frac{5}{12}$.

Moreover, by the help of this decimal fraction of a pound, to wit, .000164383, which is very near the intereſt of one pound for a day at the rate of 6 *per cent. per annum* (as will appear by the preceeding rule) the intereſt of any principal (ſuppoſed to be pounds or decimal parts of a pound) for any number of days propounded at the ſaid rate of intereſt, may be found out by multiplication only, *viz.* Firſt multiply the ſaid decimal .000164383, by the principal whoſe intereſt is required, then multiply that product by the number of days propounded, ſo ſhall this laſt product be the intereſt required; (but in theſe multiplications reſpect muſt be had to the cutting off of places in the products, according to the ſecond and third Rules of the 26th Chapter of the preceeding Book;) for example, if it be required to find the intereſt of 1000 *l.* for 131 days, at the rate of 6 *per cent. per annum*, the *Anſw.* will be found 21.534 +, or 21 *l.* 10 *s.* 8 *d.* + for according to the rule laſt given.

Another Rule for computing ſimple Intereſt for days.

.000164383

$.000164383 \times 1000 \times 131 = 21.534 +$

But at another rate of intereſt, a peculiar decimal inſtead of the ſaid .000164383 (which ſerves only for 6 *per cen. per annum*) muſt be found out by the firſt rule aforegoing before the latter rule can take place, the reaſon of which latter rule doth alſo evidently ariſe from two ſingle rules of three.

The manner of ſumming up Annuities in arrear with allowances of ſimple intereſt.

IX. When an Annuity payable yearly is an arrear for any number of years, and it is required to know what the ſame will amount unto, ſimple intereſt being computed for each particular yearly payment, from the time it became due, until the end of the term of years, the work will be as in this following example, *viz.* If an Annuity, or yearly rent of 134 *l.* 10 *s* 6 *d.* be all forborn till the end of 4 years, what will it then amount unto, ſimple intereſt being allowed at the rate of 6 *per cent. per annum* for each years rent, from the time on which it was due, until the end of the ſaid term of four years? *Anſw.* 586 *l.* 10 *s.* $6\frac{92}{100}$ *d.*

It is evident by the queſtion, that at the rate of intereſt propounded, there muſt be computed the intereſt of 134 *l.* 10 *s.* 6 *d.* (due at the third years end) for one year (to wit, the fourth year;) alſo the intereſt of the like ſum due at the ſecond years end, for two years (to wit, the third and fourth years;) likewiſe the intereſt of the ſame ſum due at the firſt years end, for three years (to wit, the ſecond, third and fourth years:) all which intereſt being added to the ſum of the four years rent, the total ſum will ſhew what the ſaid Annuity will amount

mount unto at the end of the ſaid term of 4 years.

Explication.

	years	*l.*	*s.*	*d.*
The intereſt of 134 *l.* 10 *s.* 6 *d.* at 6 *per cent. per annum*, for	1 is...	8	1	5. 16
	2 is...	16	2	10. 32
	3 is...	24	4	3. 48
The ſum of the 4 years rent (to wit, 4 times 134 *l.* 10 *s.* 6 *d.*	is...	538	2	0
All which added together give the *Anſwer* of the queſtion, to wit,	...	586	10	6.96

X. When it is required to find out how much ready Money will ſatisfie a Debt due at the end of any ſpace of time to come, by rebating or diſcompting at a given rate of ſimple intereſt, it may be effected by this rule, *viz.* Firſt, find out the intereſt of 100 *l.* at the given rate of intereſt, for the time which the ready money is to be paid beforehand, then adding the intereſt ſo found to 100 *l.* make always the ſum of that addition the firſt term in a rule of three; 100 *l.* the ſecond term; and the debt propounded to be ſatisfied the third term; laſtly, the fourth proportional found out by the ſaid *Rule of Three* ſhall be the ready money which ought to be paid in ſatisfaction of the debt propounded.

Of rebate or diſcompt of money at ſimple intereſt.

Example 1. If a debt of 100 *l.* be payable at the end of a year to come, how much ready money will diſcharge that debt by rebating or diſcomptin at the rate of *6 per cent. per annum?* *Anſw.* 94 *l.* 6 *s.*

6 *s.* 9 *d.* 2 *f.* very near; for by the *Rule of Three*,

106 . 100 :: 100. 94,3396 +

That is to ſay, if 106 *l.* (which is compos'd of 100 *l.* principal and 6 *l.* intereſt) proceeds from 100 *l.* principal forborn for a year, from what principal forborn for a year doth 100 *l.* (compos'd of principal and intereſt) proceed from? *Anſw.* 94.3396 *l.* + (or 94 *l.* 6 *s.* 9 $\frac{1}{2}$ *d.* very near) principal money: therefore 94 *l.* 6 *s.* 9 $\frac{1}{2}$ *d.* in ready money, is of equal value with 100 *l.* due at the end of the year to come; for if the ſaid 94 *l.* 6 *s.* 9 $\frac{1}{2}$ *d.* be put forth at intereſt for a year, at the rate of 6 *per cent. per ann.* it will gain 5 *l.* 13 *s.* 2 $\frac{1}{2}$ *d.* very near, which together with the ſaid 94 *l.* 6 *s.* 9 $\frac{1}{2}$ *d.* makes the 100 *l.* the debt firſt propounded to be diſcharged by rebate.

Example 2. If 150 *l.* 10 *s.* be payable at the end of 73 days to come, how much preſent money will diſcharge the ſaid debt, by rebating after the rate of 6 *per cent. per annum*? *Anſw.* 148 *l.* 14 *s.* 3 $\frac{1}{2}$ *d.* + as by the following operation is manifeſt.

	days	*l.*	*days*	*l.*
I.	365 .	6 ::	73 .	1,2

	l.	*l.*	*l.*	*l.*
II.	101,2	100 ::	150,5.	148,7154 +

That is to ſay, Firſt, I ſeek by a ſingle *Rule of Three* the intereſt of 100 *l.* for 73 days, at the rate of intereſt propounded, ſaying, if 365 days (or a year) gain 6 *l.* what will 73 days gain? *Anſw.* 1 $\frac{2}{10}$ *l.* or 1,2. *l.* Then adding the ſaid 1. 2 to 100, I ſay, by a ſecond *Rule of Three*, if 101. 2 *l.* principal and intereſt, payable at the end of 73 days to come, be equivalent to 100 *l.* ready money, what ready money is 150 *l.* 10 *s.* (or 150. 5) payable at the end of 73 days to come equivalent unto? So by multiplying and dividing (according to the rules of decimal Multiplication and Diviſion explained in Chapter 26 and 27 of the preceding Book) the quotient or anſwer of the queſtion will be found 148.7154+, that is, 148 *l.* 14 *s.* 3½ *d.*+ for the decimal .7154 being valued according to the brief way at the end of the fourth rule of the 26th Chapter, will by inſpection only be diſcovered to be 14 *s.* 3½ *d.* which rule I ſhall here once for all, adviſe the Learner to be well acquainted with.

The Proof.

Seek (by the *Rule of Three*) what the ready money found as aforeſaid will gain, in ſo much time as it is paid before hand at the rate of intereſt propounded; then having added this gain to the ſaid ready money, if the ſum be equal to the debt firſt propounded to be ſatisfied by rebate, the ready money was rightly found out. So the laſt example will be thus proved,

l. *l.* *l.* *l.*
100. 1.2 :: 148.7154 . (1.7845

Which fourth proportional 1.7845 being added o 148.7154, the ſum will be 150.4999+, which oth not want a farthing of 150 *l.* 10 *s.* the debt rſt propounded.

Of the preſent worth of Annuities by diſcompting at ſimple intereſt.

XI. When it is required to find the preſent worth of an annuity, by rebating or diſcompting at a given rate of ſimple intereſt, the operation will be as in the following example, *viz.* How much preſent money is equivalent to an annuity or rent of 100 *l.* *per annum* to continue five years, rebate being made at the rate of 6 *l.* for 100 *l.* for one year, at ſimple intereſt? *Anſw.* 425 *l.* 18 *s.* 9 ½ *d.* very near.

It is manifeſt that there muſt be computed the preſent worth of 100 *l.* due at the firſt years end; alſo the preſent worth of 100 *l.* due at the ſecond years end, and in like manner for the third, fourth and fifth years; all which particular preſent worths being added together, the aggregate or ſum will be the total preſent Worth of the Annuity, to wit, in the example above propounded, $425\frac{8286150}{8821267}$ *l.* that is, 425 *l.* 18 *s.* 9 ½ *d.* very near.

The operation by decimals (which will come near enough to the truth) will be as followeth, *viz.*

	l.	*l.*	*l.*	*l.*
1.	106 .	100 ::	100 .	94,33962 +
2.	112 .	100 ::	100 .	89,28571 +
3.	118 .	100 ::	100 .	84,74576 +
4.	124 .	100 ::	100 .	80,64516 +
5.	130 .	100 ::	100 .	76,92307 +
			Anſw.	425,93933 +

Here by the way, from the manner of reſolving the laſt mentioned queſtion, that *Rule* commonly called *Equation* of *Payments*, which is inſiſted on by divers *Arithmetical Writers*, will be found erroneous, which I thus prove.

1. Since that rule aims at the reducing of ſeveral days of payment, upon which particular ſums of money are due, unto a mean time upon which the aggregate or total of thoſe particular ſums ought to be paid, without damage to the *Debitor* or *Creditor*, there muſt be neceſſarily ſome rate of intereſt implied; for otherwiſe why may not any day at pleaſure be aſſigned for one intire payment?

2. If ſome rate of intereſt be implied, then equity requires that the preſent worth of the total ſum payable at one entire payment, rebate or diſcompt being made according to that rate of intereſt, may be equal to the ſum of the preſent worths of the particular ſums of money, rebate being made at the ſame rate of intereſt.

3. In regard the ſaid *Rule* doth mention no particular rate of intereſt, it ought to be true at any rate of intereſt whatſoever.

4 Let us therefore examine the ſaid *Rule* according to the rate of 6 *per cent. per annum*, ſimple intereſt, by taking the laſt mentioned queſtion for an example, which according to the accuſtomed manner) will be thus ſtated, *viz.* If 500 *l.* ought to be paid by five equal yearly payments, to wit, 100 *l.* at each years end, what time ought to be given for the payment of the ſaid 500 *l.* at one entire payment, without loſs either to the *Debitor* or *Creditor*.

5. By proceeding according to the ſaid *Rule* of *Equation* of *Payments* (which ſaith, if the ſum of the

products, ariſing from the multiplication of each particular ſum of money by its reſpective time, be divided by the ſum or aggregate of the ſaid particular ſums of money, the quotient will be the mean time to be aſſigned for one entire payment) there will be found three years, which time (according to the ſaid rule) ought to be given for the payment of the whole 500 *l.*

6. Now if 500 *l.* due at the end of three years to come be worth as much in preſent money, as is the preſent worth of an *Annuity* of 100 *l.* to continue five years, then the ſaid *Rule of Equation* is true; otherwiſe falſe; but the preſent worth of 500 *l.* due at the end of three years to come, rebate being made at the rate of 6 *per centum*, *per annum*, *ſimple intereſt*, will be found (by the tenth rule of this Chapter) to be 423 *l.* 14 *s.* 6 *d.* 3 *f.* very near; alſo the preſent worth of the ſaid *Annuity*, rebate being made as before, is found, (as appeareth by the reſolution of the laſt mentioned queſtion) to be 425 *l.* 18 *s.* 9 $\frac{1}{2}$ *d.* very near; wherefore it is evident that the *Creditor* loſeth 2 *l.* 4 *s.* 2 $\frac{3}{4}$ *d.* very near, by receiving the whole 500 *l.* at the three years end: Moreover at 6 *per centum*, *per annum*, *compound intereſt*, he would loſe 1 *l.* 8 *s.* 6 *d.* very near; as will be manifeſt by the *Tables of compound intereſt* hereafter expreſſed: So that the loſs will be either more or leſs according as the rate of intereſt doth differ: And therefore I conclude the ſaid Rule (as alſo all other rules or reſolutions of queſtions which have dependance thereon) to be erroneous.

Although queſtions of this nature ſeldom come into practice, yet he that will take the pains, may find out ſuch a mean time as is required by the ſaid

Rule of Equation of *payments*, at any rate of ſimple intereſt by this following rule, *viz.*

Firſt, By the preceding tenth Rule of this Chapter find out the preſent worth of every particular ſum in the queſtion payable at a time to come, by rebating at the rate of intereſt agreed on; then find in what time the ſum of thoſe preſent worths will be augmented unto the total of all the particular ſums payable at times to come, according to the firſt agreement, ſo ſhall the time found out be the mean time for the payment of the whole debt: thus the mean or equated time in the laſt example will be found to be 2.8979, *&c.* years (not three years, as the ſaid *Rule* of *Equation* of *payments* would have it) for by rebating at 6 *per cent. per annum*, ſimple intereſt, 500 *l.* payable at the end of 2.8979, *&c.* years to come (that is 2 years and 328 days very near) is worth in ready money 425 *l.* 18 *s.* 9½ *d.* very near, and the ſame ready money is alſo the preſent value of 100 *l.* Annuity for five years, at the ſame rate of intereſt, as before hath been manifeſted. But to return to the path from which I have made a digreſſion.

From the preceding tenth rule of this Chapter the following *Tables* I. and II. are deduced, whoſe conſtruction and uſe are afterwards declared.

Years	*Table* I. Which ſheweth in decimal parts of a pound, the preſent worth of one pound due at the end of any number of years to come, not exceeding 7 years, at the rate of 6 *per centum, per annum,* ſimple intereſt.	*Years*	*Table* II. Which ſheweth in pounds and decimal parts of a pound, the preſent worth of one pound Annuity, to continue any number of years not exceeding 7, at the rate of 6 *per centum, per annum,* ſimple intereſt.
1	.943396	1	. 943396
2	.892857	2	1 . 836253
3	.847457	3	2 . 683710
4	.806451	4	3 . 490162
5	.769230	5	4 . 259393
6	.735294	6	4 . 994687
7	.704225	7	5 . 698912

The Conſtruction of Table I.

The numbers in the firſt Table which are placed right againſt the numbers of Years 1, 2, 3, 4, 5, 6, and 7, are decimal fractions, one pound of Engliſh money being the Integer, and are thus found (according to the preceding tenth Rule of this Chapter) *viz.*

106 . 100 : : 1 . ,943396 +
112 . 100 : : 1 . ,892857 +
118 . 100 : : 1 . ,847457 +

whereby it appears, that 1 *l.* due at the end of a year to come, is worth in ready money .943396+, that is, 18 *s.* 10 *d.* 1 *f.* and ſomewhat more. Alſo 1*l.* due at the end of two years to come, is worth in ready money .892857+, or 17 *s.* 10¼ *d.* rebate being made at the rate of 6 *per centum*, *per annum*, ſimple intereſt, the like is to be underſtood of the reſt of the numbers in *Table* I. which may be continued to more years, and other *Tables* alſo of rebate may be framed upon the ſame ground, for months, or days, by the ingenious Artiſt.

The Uſe of Table I.

The practical uſe of the ſaid firſt Table will be manifeſt by ſolving this following queſtion; *viz.* How much ready money will diſcharge 345 *l.* 15 *s.* 6 *d.* due at the end of five years to come, by rebating ſimple intereſt at the rate of 6 *per centum per annum*? *Anſw.* 265 *l.* 19 *s.* 7¼*d.* which is thus found out; *viz.* In the preceding *Table* I. right againſt 5 years, I find the decimal .76923, which ſhews that 1 *l.* due at the end of five years to come is worth in ready money .76923 (that is, 15 *s.* 4½.) then inſtead of 15 *s.* 6 *d.* mentioned in the queſtion propounded, taking the decimal .775 which is equal to 15 *s.* 6 *d.* (the ſame being reduced according to the fifth Rule of the 23d. Chapter of the preceding Book) I ſay, by the *Rule of Three.*

1 . ,76923 :: 345.775 . (265.9805 +

That is to ſay, if 1*l.* give .76923 *l.* what will 345 .775 *l.* give? *Anſ.* 265.9805 *l.* for multiplying 345 .775 by .76923, according to the ſecond Rule of the 26 Chapter of the preceding Book, the product will be 265.9805, that is, 265*l.* 19 . 7¼*d.*

The Conſtruction of Table II.

The numbers in the ſecond *Table* are found out by the addition of thoſe in the firſt, *viz.* the firſt number in the latter *Table* is the ſame with the firſt number in the former, the ſecond in the latter is the ſum of the firſt and ſecond in the former, the third in the latter is the ſum of the firſt, ſecond and third in the former, and in that manner the reſt are found; (the reaſon of which compoſition is manifeſt from the example of the eleventh rule aforegoing;) otherwiſe the numbers in *Table* II. may be found more eaſily thus, *viz.* the firſt number in the ſaid *Table* II. is the ſame with the firſt number in *Table* I. the ſecond number in the latter *Table* is compos'd of the ſecond number in the former and the firſt in the latter, the third number in the latter *Table* is compos'd of the third number in the former and the ſecond in the latter, the fourth in the latter is compos'd of the fourth in the former and the third in the latter; the like is to be underſtood of the reſt of the numbers in *Table* II. which might be continued to more years, and fitted to other rates of intereſt, but I ſhall ſpare that labour, in regard a more equal way of finding out the preſent worth of an Annuity, agreeable to the accuſtomed and practical rates of buying and ſelling Annuities of Rents, for terms of years, is grounded upon a computation of intereſt upon intereſt as will hereafter be made manifeſt, for at ſimple intereſt an Annuity will be overvalued.

The uſe of Table II.

The uſe of *Table* II. will appear by this following

ing example; *viz.* What is the preſent worth of an Annuity of 100 *l. per annum* payable yearly during the term of five years, diſcompt or rebate being made at the rate of 6 *per centum, per annum,* ſimple intereſt? *Anſwer,* 425 *l.* 18 *s.* 9½*d.* very near, which is thus found out, *viz.* In the preceding *Table* II. right againſt five years, I find this number 4.259393, which ſhews that an Annuity of 1 *l.* payable yearly during five years, is worth in ready money 4.259393 *l.* (that is 4 *l.* 5 *s.* 2 *d.* and ſomewhat more) therefore, I ſay, by the *Rule of Three*;

l. *l.* *l.* *l.*
1 . 4.259393 :: 100 . (425.9393.

That is to ſay, if 1*l.* give 4.259393 *l.* what will 100 *l.* give? *Anſw.* 425 *l.* 18 *s.* 9½*d.* very near, for by multiplying 4.259393 by 100, the product (according to the ſecond rule of the 26th. Chapter of the preceding Book) is 425.9393, that is, 425 *l.* 18 *s.* 9½*d.* very near. Which operation being compared with the manner of ſolving the ſame queſtion before mentioned in the eleventh Rule of this Chapter, the great benefit of Tables of this kind in point of expedition will be apparent.

Of the forbearance of Money at compound intereſt.

XII. When it is required to know, unto what ſum of money any propounded principal forborn any number of years will at the end of ſuch term be augmented unto, intereſt upon intereſt being computed at a given rate, there muſt be found a rank of continual proportionals, more in number by one than is the number of years in the queſtion; of which proportionals the firſt is the principal aſſigned, the ſecond muſt increaſe or proceed from the firſt, the third from the ſecond, *&c.* in ſuch manner or rate, as 106 proceeds from 100 (or as 108 from 100, if the rate of intereſt be 8 *per centum*) then will the laſt proportional be the Anſwer of the queſtion: So if 300 pounds principal money be put forth at intereſt upon intereſt, at the rate of 6 *l.* for 100 *l.* for one year, and all forborn until the end of 4 years, there will then be due 378.743088, or 378 *l.* 14 *s.* $10\frac{1}{2}$ *d.* very near, as by the four following *Rules of Three* is manifeſt.

$$100 \;.\; 106 :: \left\{ \begin{array}{ll} 300 & .\; 318 \\ 318 & .\; 337.08 \\ 337.08 & .\; 357.3048 \\ 357.3048 & .\; 378.743088 \end{array} \right.$$

For the ſaid 300 *l.* will at the firſt years end be augmented unto 318 *l.* which 318 *l.* being put forth as a *principal* for 1 year, will (at the ſecond years end) be augmented unto 337.08, again this 337.08 being put forth as a *principal* for 1 year, will (at the third years end) be augmented unto 357.3048, in like manner 357.3048 being put forth as a *principal* for 1 year, will (at the fourth years end) be augmented unto 378.743088, which is the number required by the queſtion. And if the work be well examined, it will appear (as was before declared) that the *principal* firſt aſſigned, to wit 300 *l.* and the numbers reſulting ſucceſſively at the ends of the ſeveral years are *continual proportionals*, *viz.* theſe five numbers are ſo qualified, that if the

300 | 318 | 337.08 | 357.3048 | 378.743088

ſecond be multiplied by it ſelf, the product will be equal to the product of the firſt and third; alſo if the third be multiplied by it ſelf, the product will be equal to the product of the ſecond and fourth; in like manner, if there were more *continual proportionals* in a rank, if any one proportional which is placed between two next on each ſide of ſuch one, be multiplied by it ſelf, the product will be equal to the product of thoſe two extreams. (which is a property peculiar to continual proportionals.)

Two numbers being given to find a third, a fourth, a fifth, &c. *in continual proportion.*

Note here by the way, that if any two numbers be propounded, ſuppoſe 300 and 318, and it be required to find to them a third, a fourth, a fifth, &c. in continual proportion, multiply the ſecond proportional 318 by it ſelf, and divide the product 101.124 by the firſt proportional 300, ſo ſhall the quotient 337.08 be a third in continual proportion: In like manner if you multiply the third proportional 337.08 by it ſelf, and divide the product 113622.9264 by the ſecond proportional 318, the quotient 357.3048 ſhall be a fourth in continual proportion, and after the ſame manner a fifth, a ſixth, or as many as you pleaſe may be found out.

From what hath been ſaid by way of Explication of the preceding twelfth Rule, the following *Table* III. is deduced, the conſtruction and uſe whereof is afterwards declared.

TABLE

TABLE III.

Which ſheweth what one Pound will amount unto, being forborn unto the end of any term of years under 31, compound intereſt being computed yearly, at any of theſe rates, to wit, 4, 5, 6, 7, 8, 9, 10, 11, *and* 12 per cent. per annum.

Years	4	5	6	7	8	9	10	11	12
1	1,04000	1.05000	1,06000	1,07000	1.08000	1.09000	1.10000	1.11000	1.12000
2	1.08160	1.10250	1,12360	1.14490	1.16640	1.18810	1,21000	1.23216	1.25440
3	1.12486	1.15762	1.19101	1,22504	1,25971	1.29502	1.33100	1.36763	1.40492
4	1.16985	1.21550	1.26247	1.31079	1.36048	1.41158	1.46410	1.51807	1.57351
5	1.21665	1.27628	1.33823	1.40255	1.46932	1.53862	1.61051	1.68505	1.76234
6	1.26531	1.34009	1.41851	1,50073	1.58087	1.67710	1.77156	1.87041	1.97382
7	1.31593	1.40710	1.50363	1.60578	1.71382	1.82803	1.94871	2.07616	2.21068
8	1.36856	1.47745	1.59384	1.71818	1.85093	1.99256	2.14358	2.30453	2.47596
9	1.42331	1.55132	1.68947	1.83845	1.99900	2.17189	2.35794	2.55803	2.77307
10	1.48024	1.62889	1.79084	1.96715	2.15892	2.36736	2.59374	2.83942	3.10584
11	1.53945	1.71033	1.89829	2.10485	2.33163	2 58042	2.85311	3.15175	3.478 4
12	1.60103	1.79585	2.01219	2.25219	2.51817	2.81266	3.13842	3.49849	3.89597
13	1.66507	1.88564	2.13292	2.40984	2.71962	3.06580	3.45227	3.88328	4.36349
14	1.73167	1.97993	2.26090	2.57853	2.93719	3.34172	3.79749	3.31044	4.88711
15	1.80094	2 07892	2.39655	2.75903	3 17216	3.64248	3.17724	3.78458	4.47356

A continuation of the preceding Table III.

Years	4	5	6	7	8	9	10	11	12
16	1.87298	2.18287	2.54035	2.95216	3.42549	3.97030	4.59497	5.31089	6.13039
17	1.94790	2.29 01	2.69277	3.15881	3.70001	4.31763	5.05447	5.89509	6.86604
18	2.02581	2.40661	2.85433	3.37993	3.99601	4.71712	5.55991	6.54355	7.68996
19	2.10684	2.52695	3.02559	3.61652	4.31570	5.14166	6.11590	7.26234	8.61276
20	2.10112	2.65329	3.20713	3.86968	4.66095	5.60441	6.72749	8.06231	9.64629
21	2.27876	2.78596	3.39956	4.14056	5.03383	6.10880	7.40024	8.94916	10.80384
22	2.36991	2.92529	3.60353	4.43040	5.43654	6.65860	8.14027	9.93357	12.10031
23	2.46471	3.07152	3.81975	4.74053	5.87146	7.25787	8.95430	11.02626	13.55234
24	2.56330	3.22509	4.04893	5.07236	6.34118	7.91108	9.84973	12.23915	15.17862
25	2.66583	3.38535	4.29187	5.42743	6.84847	8.62308	10.83470	13.58546	17.00006
26	2.77246	3.55567	4.54938	5.80735	7.39635	9.39915	11.91817	15.07986	19.04007
27	2.88336	3.73345	4.82234	6.21386	7.98806	10.24508	13.10999	16.73864	21.32488
28	2.99870	3.92012	5.11168	6.64883	8.62710	11.16713	14.42099	18.57990	23.88386
29	3.11865	4.11613	5.41838	7.11425	9.31727	12.17218	15.86309	20.62369	26.74993
30	3.24330	4.32194	5.74349	7.61225	10.06265	13.26767	17.44940	22.89229	29.55993

The Conſtruction of the preceeding Table III.

The numbers 1, 2, 3, 4, *&c.* to 30, in the firſt Column on the left hand ſignifie years; the numbers 4, 5, 6, 7, 8, 9, 10, 11, and 12, placed at the head of the reſt of the Columns ſignifie rates of intereſt, for 100 *l.* lent for a year, and the numbers placed in the ſeveral columns underneath thoſe rates of intereſt, are found out by the *Rule of Three* in decimals, in manner following, *viz.*

I. | 100 : 104 :: 1 :(1.04
II. | 100 : 104 :: 1.04 :(1.0816
III. | 100 : 104 :: 1.0816 :(1.12486

That is to ſay, Firſt if 100 *l.* put forth at intereſt for a year be augmented to 104 *l.* at the years end, what will 1 *l.* be then augmented unto at the ſame rate? *Anſw.* 1.040 *l.* (that is 1 *l.* 0 *s.* 9 *d.* 2 *f.* and ſomewhat more) which 1.04 (or 1.04000, the Cyphers after the 4 being of no value in decimals) is the firſt number in the ſecond column belonging to 4 *per cent.* and is placed right againſt 1 year in the firſt column.

Secondly, ſay if 100 *l.* lent for a year be augmented to 104 *l.* at the years end, what will 1.04 *l.* be then augmented unto at the ſame rate? *Anſw.* 1,0816 *l.* (that is 1 *l.* 1 *s.* 7 *d.* 2 *f.*+) which 1.0816 is the ſecond number in the ſaid column of 4 *per cent.* and is placed right againſt 2 years in the firſt column.

Thirdly, as 100 is to 104, ſo is 1.0816 to 1.124864 (or 1 *l.* 2 *s.* 5 *d.* 2*f.*+) which 1.12486 is the third number in the column of 4 *per cent.* and is placed right againſt 3 years in the firſt column. Hence it appears, that 1 *l.* at 4 *per cent. per annum*, compound intereſt, will at the end of 3 years be augmented unto 1.124864 *l.* (that is, 1 *l.* 2 *s.* 5 *d.* 2 *f.* and ſomewhat more.)

After the ſame manner the reſt of the numbers in the ſecond column, as alſo in the other columns are found out *(mutatis mutandis.)*

The Uſe of the preceding third Table.

Queſt. 1 What will 136 *l.* 15 *s.* 6 *d.* be augmented unto, being forborn 20 years, intereſt upon intereſt being computed at the rate of 6 *per cent. per annum? Anſw.* 438 *l.* 13 *s.* 1 *d.* very near, which is thus found out.

Firſt, looking into the fourth column of the ſaid third *Table*, to wit, that column which hath the figure 6 placed at the head of it, I find right againſt 20 years the number 3.20713, which ſhews that 1 *l.* being continued 20 years at 6 *per cent. per annum*, compound intereſt, and all forborn until the end of the ſaid term will be augmented unto 3.20713 *l.* (that is 3 *l.* 4 *s.* 1 *d.* 2 *f.* and ſomewhat more) therefore after the 15 *s.* 6 *d.* in the queſtion is reduced to the decimal .775 (by the ſixteenth rule of the 23 Chapter of the preceding Book) I multiply the ſaid tabular number 3.20713 by 136.775 (the ſum propounded in the queſtion) according to the ſecond rule of the 26th. Chapter, ſo the product is found to be 438.665, &c. that is, 438 *l.* 13 *s.* 1 *d.* for the anſwer of the queſtion. View the operation here following.

1 . 3.20713 : : 136.775 . (438.665 +
136.775

1603565
2244991
2244991
1924278
962139
320713

438|65520575

Queſt. 2. If 320 *l.* be forborn 11 years, at intereſt upon intereſt at 5 *per centum, per annum,* what will be due at the end of thoſe eleven years for principal and intereſt? *Anſw.* 547 *l.* 6 *s.* 1 *d.* +. For in the third Column of the third Table, under the figure 5 at the head of the column and right againſt 11 years you will find this number 1.71033, which ſhews that 1 *l.* at the end of 11 years will at five *per centum, per annum,* compound intereſt, be augmented to 1.71033 (that is 1 *l.* 14 *s.* 2 *d.* 1 *f.* and ſomewhat more) wherefore by multiplying the ſaid 1.71033 by 320 the number of pounds propounded in the queſtion) the product will be 547.305, &c. that is 547 *l.* 6 *s.* 1 *d.* + for the anſwer of the queſtion. See the following operation:

$$\begin{array}{l} 1 : 1.71033 :: 320 \,.\, (547.305 + \\ \quad\quad\quad 320 \\ \hline \quad\quad 3420660 \\ \quad\quad 513099 \\ \hline \quad\quad 547|30560 \end{array}$$

After the ſame manner the numbers belonging to any of the other rates of intereſt mentioned in the third Table are to be uſed.

The manner of ſumming up Annuities in arrear with allowances of intereſt upon intereſt.

XIII. When an Annuity payable yearly is in arrear for any number of years, and it is required to know what the ſame will amount unto, compound intereſt being computed or each particular Annuity from he time it became due until the end of the term of years, the work will be as in the ollowing example; *viz.* Suppoſe an Annuity of 300 *l.* payable at yearly payments be forborn, and ll unpaid until the end of four years, the queſtion s, what will then be due, compound intereſt being computed at the rate of 6 *per cent. per annum,* or each yearly payment from the time it becomes due to the end of the ſaid term of four years? *Anſwer,* 1312 *l.* 7 *s.* 8 *d.* very near.

It is evident by the queſtion, that there muſt be omputed what 300 *l.* due at the third years end will be augmented unto in one year (to wit, the ourth year) at 6 *per cent.* Alſo what 300 *l.* due at the econd years end will be augmented unto in two ears (to wit, the third and fourth years; likewiſe what 300 *l.* due at the firſt years end, will be augmented unto, in the three following years (to wit the ſecond, third and fourth years) all which ſums being added to 300 *l.* (the payment due at the end of the fourth year, which is incapable of any improvement) the aggregate or ſum will be the total money in Arrear at the end of the fourth year, to wit, 1312, $\frac{3848}{10000}$ *l.* as may appear by the following operation, *viz.*

	l.
The laſt payment of the Annuity due at the end of the fourth year is	300.
Again, the 300 *l.* due at the third years end, will in one year after the rate of 6 *per centum*, be augmented unto	318.
Alſo 300 *l.* due at the ſecond years end, will in two years at the rate of 6 *per centum*, *per annum*, compound intereſt, be augmented unto (as appears by the firſt example of the twelfth Rule aforegoing.)	337.08
In like manner; 300 *l.* due at the firſt years end, will in three years be augmented unto	357.3048
The ſumm due at the four years end	1312.3848

The invention of the numbers before-mentioned being well examined, it will appear, that if an Annuity or Rent payable at yearly payments be improved

mproved to the utmoſt as intereſt upon intereſt,
nd all forborn or reſpited unto the end of certain
ears, the total then due will be the ſum of a rank
f continual proportionals as many in number as
here are yearly payments, the firſt of which proportionals is the firſt (or any one) years rent, and
he ſecond proportional proceeds from the firſt in
he ſame rate as 106 proceeds from 100, if the rate
f intereſt be 6 *per centum*, (or as 108 proceeds
rom 100, if the rate of intereſt be 8 *per centum*,
&c.) and ſo likewiſe the third from the ſecond,
he fourth from the third, *&c.* (after the manner
f the operation in the firſt example of the twelfth
Rule of this Chapter.)

Otherwiſe.

Find a principal which may have ſuch propor-
ion to 300 as 100 hath to 6, and ſay by the *Rule*
f *Three*,

6 . 100 : : 300 . 5000.

That is to ſay, as 6 *l.* intereſt hath 100 *l.* for a
rincipal, ſo 300 *l.* intereſt hath 5000 *l.* for a
rincipal; then ſeek what 5000 *l.* will be augmen-
ed unto, being forborn four years at 6 *per centum*,
er centum, compound intereſt (after the manner
f the firſt Example of the 12th. Rule aforegoing:)
o will you find 6312.3848, from which ſubtract-
ng the ſaid principal 5000 *l.* the remainder (as be-
ore) is 1312.3848 *l.* being the ſum which 300 *l.*
nnuity will be augmented unto at the end of
ur years, according to the ſaid rate of intereſt,
e Annuity being payable at yearly payments.

The Reaſon of the latter Rule.

If a principal be put forth at intereſt upon intereſt payable by yearly payments, and all be forborn untill the end of certain years, the total then due is equal to the aggregate or ſum of theſe three numbers, to wit, the ſaid principal firſt put forth; the ſum of the annual ſimple intereſt of that principal; and the utmoſt improvement of thoſe ſimple intereſts by computing intereſt upon intereſt; wherefore if from the ſaid aggregate the firſt principal be ſubtracted, the remainder muſt neceſſarily conſiſt of the ſum of the annual ſimple intereſts, (which are in the nature of an Annuity) and the utmoſt improvement of thoſe ſimple intereſts (or Annuity) by computing intereſt upon intereſt.

The Conſtruction of the following Table IV.

Upon the aforeſaid grounds, the following *Table* IV. is calculated, to ſhew what one pound Annuity, payable at yearly payments, and forborn any number of years under 31, will amount unto by computing intereſt upon intereſt at any of the rates expreſt at the head of the ſaid Table.

But the ſame *Table* may be more eaſily compoſed by the addition of the numbers in the preceding *Table* III. in this manner, *viz.* the firſt number in each of thoſe columns in the following *Table* IV. at the head whereof are placed the numbers 4, 5, 6, 7, 8, 9, 10, 11, and 12, ſignifying rates of intereſt *per centum*, is 1 or unity, the ſecond number in each of theſe columns in the latter *Table* is compos'd of 1 or unity, and the firſt number in the reſpective columns of the ſaid preceding *Table* III.

Alſo the third number in each of the ſaid columns of this *latter Table* is compos'd of 1, and the ſum of the firſt and ſecond number of the reſpective columns of the *former Table*, and in that order the reſt are found out; or more eaſily thus: the third number in the *latter Table* is compos'd of the ſecond number in the *latter*, and of the ſecond in the *former*; the fourth number in the *latter* is compos'd of the third in the *latter*, and of the third in the *former*, &c. But you are to obſerve that according to either of theſe ways of compoſing the fourth *Table* by Addition, the numbers in the preceding *Table* III. ought to be continued to more places than are there expreſt to prevent error which may happen by adding of defective decimal fractions.

TABLE IV.

Which ſheweth what one Pound Annuity, payable by yearly payments, and forborn any number of years under 31, *will amount unto, at the end of the term, compound intereſt being computed at any of theſe rates, to wit,* 4, 5, 6, 7, 8, 9, 10, 11, *and* 12 per cent. per annum.

Years	4	5	6	7	8	9	10	11	12
1	1.00000	1.00000	1.00000	1.00000	1.00000	1.00000	1.00000	1.00000	1.00000
2	2.04000	2.50000	2.06000	2.07000	2.08000	2.09000	2.10000	2.11000	2.12000
3	3.12160	3.15250	3.18360	3.21490	3.24640	3.27810	3.31000	3.34210	3.37440
4	3.24646	4.31012	4.37461	4.43994	4.50611	4.57312	4.64100	4.70973	4.77932
5	5.41632	5.52563	5.63709	5.75073	5.86660	5.08471	6.10510	6.22780	6.35284
6	6.63297	6.80190	6.97531	7.15329	7.33592	7.52333	7.71561	7.91285	8.11518
7	7.89829	8.14200	8.39383	8.65402	8.92280	9.20043	9.48717	9.78327	10.08901
8	9.21422	9.64900	9.89746	10.25980	10.63662	11.02847	11.43588	11.85943	12.29969
9	10.58279	11.02656	11.49131	11.97798	12.48755	13.02103	13.57947	14.16397	14.77565
10	12.00610	12.55789	13.18079	13.81644	14.48656	15.19292	15.93742	16.72200	17.54873
11	13.48635	14.20678	14.97164	15.78359	16.64548	17 56029	18.53116	19.56142	20.65458
12	15.02580	15.91712	16.86994	17.88845	18.97712	20.14071	21.38428	22.71318	24.13313
13	16.62683	17.71298	18.88213	20.14064	21.49529	22.95338	24.52271	26.21163	28.02910
14	18.29191	19.59863	21.01506	22.55048	24.21492	26.01918	27.97498	30.09491	32.39260
15	20.02358	21.57856	22.27596	25.12902	27.15211	29.36091	31.77248	34.40535	37.27971

A continuation of the preceding Table IV.

Years	4	5	6	7	8	9	10	11	12
16	21.82453	23.65749	25.67252	27.88805	30.32428	33.00339	35.94972	39.18993	42.75328
17	23.69751	25.84036	28.21287	30.84021	33.75022	36.97370	40.54470	44.50084	48.88367
18	25.64541	28.13238	30.90565	33 99903	37.45024	41.30133	45.59917	50.39592	55.74971
19	27.67120	30.53900	33.75999	37.37896	41.44626	46.01845	51.15009	56 93948	63.43968
20	29.77807	33.06595	36.78559	40.99549	45.76196	51.16011	57.27499	64.20283	72.05244
21	31.96920	35.71925	39.99272	44.86517	50.42292	56.76453	64.00249	72.26514	81.69873
22	34.24796	38 50521	43.39228	49.00573	55.45675	62.87333	71.40274	81.21430	92.50258
23	36.61788	41 43047	46.99582	53.43614	60.89329	69.53193	79.54302	91.14788	104.60289
24	39.08260	44.50199	50.81557	58.17667	66.76475	76.78981	88.49732	102.17415	118.15524
25	41.64590	47.72709	54.86451	63 24903	73.10599	84 70087	98.34705	114 41330	133.33387
26	44.31174	51.11345	59.15638	68.67646	79.95441	93.32397	109.18176	127.99877	150.33393
27	47.08421	54.66912	63.70576	74.48382	87.35076	102.72313	121.09994	143.07863	169.37400
28	49.96758	58.40258	68.52810	80.69769	95.33882	112.96821	134.20993	159.81728	190 69888
29	52.96628	62.32271	73.63979	87.34652	103.96593	124.13535	148.63092	178.39718	214.58275
20	56.08493	66 43884	79.05818	94 46078	112.28321	136.30753	164.49402	199.02087	241.33268

The uſe of the preceding Table IV.

The uſe of the ſaid fourth *Table* will be manifeſt by the manner of ſolving this Queſtion, *viz.* if an Annuity of 20 *l.* payable by yearly payments for 15 years; be all forborn or unpaid until the end of the ſaid term, what will it then amount unto, upon a computation of intereſt upon intereſt, at the rate of 6 *per centum, per annum? Anſw.* 465*l.* 10*s.* 4 *d.* 2 *f.* very near, as by the following operation is evident; For in the column belonging to 6 *per centum* (to wit, that column which hath the figure 6 placed at the head of it) right againſt 15 years, you will find 23.27596, which ſhews that an Annuity of 1 *l.* payable at yearly payments for 15 years, will at the end of the ſaid term (compound intereſt being computed at 6 *per cent. per annum*) amount unto 23.27596 *l.* (or 23 *l.* 5 *s.* 6 *d.*+) Therefore multiplying the ſaid tabular number 23.27596 by 20. (becauſe the Annuity propounded is 20 *l.*) the product will be 465.519 +, that is 465 *l.* 10 *s.* 4*d.* 2*f.* which is the anſwer of the queſtion; view the following operation.

$$
\begin{array}{r}
1 \;.\; 23.27596 :: 20 \;.\; (465.519+ \\
20 \\
\hline
465|51920
\end{array}
$$

In the ſame manner the numbers in the other columns are to be uſed.

Of rebate at Compound intereſt.

XIV. When a ſum of money is due at a time to come, and it is required to know what it is worth in ready money, rebate being made at a given rate of compound intereſt, the work will not be much different from the 12 Rule of this Chapter, *viz.* there muſt be found a ſeries or rank of continual proportionals more in number by one, than is the number of years in the queſtion; of which proportionals, the firſt is the money propounded to be rebated, the ſecond muſt decreaſe or leſſen from the firſt, the third from the ſecond, &c. in ſuch manner or rate as 100 decreaſeth from 106 (or as 100 from 108, if the rate of intereſt be 8 *per cent.*) then will the laſt proportional be the anſwer of the queſtion: So if $378\frac{743088}{1000000}$*l.* be due at the end of four years wholly to come, it will be found to be worth in ready money 300*l.* rebate being made at compound intereſt at 6 *per cent.* as by the four following *Rules of Three* is manifeſt, which may be proved by the preceding twelfth Rule, where it will appear that 300*l.* being forborn four years, will at the ſaid rate of compound intereſt be augmented unto 378.743088 *l.*

$$106 \,.\, 100 :: \left\{ \begin{array}{ll} \text{l.} & \text{l.} \\ 378.743088 & .\ 357.3048 \\ 357.3048 & .\ 337.08 \\ 337.08 & .\ 318. \\ 318. & .\ 300. \end{array} \right.$$

Upon this ground the following *Table* V. is calculated, to ſhew what one pound due at the end of any number of years to come, is worth in preſent money, rebate being made at the rates of compound intereſt, mentioned in the ſaid *Table*; by the help whereof and of *Multiplication*, queſtions of rebate for any ſum propounded may be performed without conſiderable error.

TABLE

TABLE V.

Which ſheweth what one pound, payable at the end of any term of years to come under 31, is worth in ready money, diſcompt or rebate being yearly computed at any of theſe rates, to wit, 4, 5, 6, 7, 8, 9, 10, 11 *and* 12 per cent. per ann. *compound intereſt.*

Years	4	5	6	7	8	9	10	11	12
1	.961538	952381	943396	934579	.925925	.917431	.909090	.900900	.892857
2	.924556	.907029	.889996	.873438	.857338	.841680	.826446	.811622	.797193
3	.888996	.863837	839619	.816297	.793832	.772183	.751314	.731191	.711780
4	.854802	.822702	792095	762895	.735029	.708425	.683013	.658731	.635518
5	.821927	.783526	747258	712986	.680583	.649931	.620921	.593451	.567426
6	.790314	.740215	.704960	.666342	.630169	.596267	.546447	.534640	.506631
7	.759917	.710681	.655057	.622749	.583490	.547034	.513158	.481658	.452349
8	.730690	.676839	.627412	582009	.540268	.501866	.466507	.433926	.403883
9	.702586	.644608	.591898	.543933	.500248	.460427	.424097	.390924	.360610
10	.675564	.613913	.558391	.508349	.463193	.422410	.385543	.352184	.321973
11	.649580	.584679	.526787	.475092	.428882	.387532	350494	317283	.287476
12	.624596	.556837	.496989	.444012	.397113	.355534	.318630	.285840	.256675
13	.600573	.530321	468839	.414964	.367697	.326178	289664	.257514	.229174
14	.577474	.505067	442300	.387817	.340461	.299246	.263331	.231994	.204619
15	.555264	.481017	.417265	.362446	.315241	.274538	239392	.209004	.182696

A continuation of the preceding Table V.

Years	4	5	6	7	8	9	10	11	12
16	.533908	.458111	.393646	.338734	.291860	.251869	.217629	.188292	.163121
17	.513373	.436296	.371364	316574	.270269	.231073	.197844	.169632	.145644
18	.493628	.415520	.350343	.295864	.250249	.211993	.179858	.152822	130039
19	.474642	.395733	.330512	.276508	.231712	.194489	.163508	.137677	.116106
20	.456386	.376889	.311804	.258419	.214548	.178430	.148643	.124034	.103666
21	.438833	.358942	.294155	.241513	.198655	.163698	.135130	.111742	.092559
22	.421955	.341849	.277505	.225713	.183940	.150181	.122840	.100668	.082642
23	.405726	.325571	.261797	.210947	.170315	137781	.111678	.090692	.073787
24	.390121	.310067	.246978	.197146	.157699	126405	.101525	.081705	.065882
25	.375116	.295302	.232998	.184249	.146018	.115967	.092296	.073608	.058823
26	.360689	.281240	.219810	.172195	.135201	.106392	083905	.066313	.052520
27	.346816	267848	.207367	.160930	.125186	.097607	.076277	.059742	.046893
28	.333477	.255093	.195630	.150402	.115913	.089548	.069343	.053821	.041869
29	.320651	.242946	.184556	.140562	.107327	.082154	063039	.048487	.037383
30	.308318	.231377	.174110	.131367	.099377	.075371	057308	.043682	.033377

The Conſtruction of the preceding Table V.

The numbers 1, 2, 3, 4, *&c.* to 30, in the firſt column on the left hand, ſignifie years; the numbers 4, 5, 6, 7, 8, 9, 10, 11 and 12, placed at the head of the reſt of the columns ſignifie rates of intereſt for 100 *l.* lent for a year, and the numbers placed in the ſeveral columns underneath thoſe rates of intereſt are found out by the *Rule of Three* in decimals, in manner following, *viz.*

I. | 104.100 : : 1 . (,9615384615, *&c.*
II. | 104.100 : : ,9615384615+ (,9245562, *&c.*
III. | 104.100 : : ,9245562, *&c.* (,888996+

That is to ſay, Firſt, if 104 decreaſe to 100, or if 104 *l.* payable at the end of a year to come be worth 100 *l.* ready money, what ready money, is 1 *l.* due at the end of a year to come worth? *Anſw.* .9615384615 + (or 19 *s.* 2 *d.* 3 *f.* very near.) So that .961538 is the firſt decimal in the ſecond column belonging to 4 *per centum*, in *Table* V. and is placed right againſt 1 year in the firſt column.

Secondly, ſay in like manner if 104 decreaſe to 100, what will .9615384615, *&c.* (the decimal found by the firſt *Rule of Three*) decreaſe unto? *Anſw.* 9245562, *&c.* the firſt 6 places whereof, to wit, 924556 are the ſecond decimal in the ſaid column of 4 *per cent.* which is placed right againſt two years.

Thirdly, as 104 is to 100; ſo is, 9245562, *&c.* (the decimal found by the ſecond *Rule of Three*) to .888996+ (or 17*s.* 9*d.* 1*f.*+) which is the third decimal in the column of 4 *per centum.* Hence it appears, that 1 *l.* due at the end of 3 years to come is worth .888996+ (or 17*s.* 9*d.* 1*f.* and ſomewhat more) in ready money, rebate being made at the rate of 4 *per centum, per annum,*) compound intereſt.

After the ſame manner the reſt of the decimal fractions in the ſaid ſecond column, as alſo in the other columns are found out (*mutatis mutantis.*)

The uſe of the preceding Table V.

To exemplifie the ſaid fifth Table, let it be required to find out how much ready money will diſcharge a debt of 356 *l.* payable at the end of ſeven years to come, by rebating at the rate of 7 *per cent. per annum*, compound intereſt? *Anſw.* 221 *l.* 13 *s.* 11 *d.* 3 *f.* very near. For in the fifth column, at the head whereof is placed 7, ſignifying 7 *per centum*, right againſt 7 years, I find .622749, which ſhews that 1 *l.* due at the end of 7 years to come is worth in preſent money .622749 decimal parts of a pound, rebate being made at the ſaid rate of compound intereſt. Therefore multiplying the ſaid tabular number .622749 by the ſaid 356 *l.* (the debt propounded) the product (according to the ſecond Rule of the 26th. Chapter) will be 221.698, *&c.* that is, 221 *l.* 13 *s.* 11 *d.* 3 *f.* which is the Anſwer of the queſtion. See the ſubſequent operation.

1 . ,622749 : : 356 . (221.698 +
356

3736494
3113745
1868247

221'698644

In the ſame manner the numbers in the other columns are to be uſed.

To find the preſent worth of Annuities by a computation of compound intereſt.

XV. The finding out the preſent worth of an Annuity, is grounded upon this foundation, to wit, if the preſent money which is paid for the purchaſe of Annuity, to continue any term of years, be put forth at any rate of compound intereſt, and all forborn until the end of the ſaid term, and that the total money then due be put into one Scale: Alſo if the total ſum of the utmoſt improvements of the annual payments of the Annuity, put forth at the ſame rate of compound intereſt, from the time thoſe annual payments become due until the end of the term, be put into the other Scale, the Scales muſt be even, *viz.* the ſaid two total ſums of money muſt be equal one to the other.

Now to find out ſuch a preſent worth of an Annuity, there are divers ways, ſome of which I ſhall here explain by examples:

Firſt therefore let it be required to find the preſent worth of an Annuity of 378.743088 *l.* to continue three years compound intereſt being computed at 6 *per cent. per ann.* *Anſw.* 1012.3848 *l.*

It is evident by the queſtion, that there muſt be computed (after the manner of the Example upon the fourteenth Rule aforegoing) the preſent worth of $378\frac{743088}{1000000}$ *l.* due at the firſt years end, alſo the preſent worth of the like ſum due at the ſecond years end, and in like manner for the third year; all which particular preſent values being added together, the aggregate or ſum will be the total preſent worth of the Annuity propounded, *viz.*

	l.
378.743088 *l.* payable at the end of one year is worth in ready money (as is evident by the fourteenth Rule aforegoing.)	357.3048
Alſo the like ſum payable at the end of 2 years to come, is worth in ready money	337.08
Again, the like ſum payable at the end of three years to come, is worth in ready money	318.
Therefore the total preſent worth of an Annuity of 378.74088 *l.* to continue 3 years is	1012 3848

Otherwiſe,

Find a Principal which may be in ſuch proportion to the propounded Annuity 378.743088 *l.* as 100 is to 6. Which will be exactly 6312.3848 *l.* for

6 . 100 :: 378,743088 . 6312, 3848

Then ſuppoſing this principal ſo found to be a ſum due at the end of three years to come, find what it will be worth in ready money, by diminiſhing it according to the fourteenth Rule of this Chapter, ſo will you find 5300 *l.* for the ready money equivalent to the ſaid 6312.3848 *l.* due at the end of three years, which ready money 5300 *l.* being ſubtracted from the ſaid 6312.3848 *l.* leaves (as before) 1012.3848 *l.* for the preſent worth of the ſaid Annuity of 378.743088 *l.* to continue three years, compound intereſt being allowed at 6 *per centum, per annum.*

The Reaſon of the latter Rule.

It will not be difficult to apprehend, that if 6312.3848 *l.* ready money be put forth as a Principal at intereſt upon intereſt, it will at three years end be augmented unto an Aggregate or ſum compos'd of theſe three numbers, to wit, the ſaid Principal 6312.3848; the ſum of the annual ſimple intereſts of that Principal, and the utmoſt improvement of thoſe ſimple intereſts by intereſt upon intereſt: And becauſe (by the operation aforegoing) 5300 *l.* ready money (part of the ſaid ready money 6312.3848.) will at three years end be augmented unto 6312.3848 *l.* part of the ſaid Aggregate, therefore 1012.3848*l.* the complement or remaining part of the ſaid ready money 6313.3848 *l.* muſt neceſſarily be augmented unto the complement or remaining part of the ſaid Aggregate, which remaining part laſt mentioned is compoſed of the ſum of the aforeſaid ſimple intereſts, and of their utmoſt improvement at intereſt upon intereſt, that is, the ſaid remainder is the utmoſt improvement of an Annuity of 378.743088 *l.* to continue three years, compound intereſt being allowed at 6 *per centum, per annum.*

The Conſtruction of the following Table VI.

Upon the aforeſaid grounds the following Table VI. is calculated to ſhew how much ready money an Annuity of one pound to continue any number of years under 31. and payable at yearly payments, is worth, upon a computation of compound intereſt at any of the rates *per centum*, mentioned at the head of the ſaid Table. But the ſaid Table VI. may more eaſily be compos'd by the help of the preceding Table V. in this manner, *viz.* the firſt number in every of the Columns (except the Column of years) in the following Table VI. is the ſame with the firſt number in the like Columns reſpectively in the preceding Table V. the ſecond number in each of the ſaid Columns of the ſixth Table is the ſum of the firſt and ſecond numbers in the reſpective Columns of the ſixth Table; the third number in the ſaid Columns of the fifth Table is the ſum of the firſt, ſecond and third numbers in the reſpective Columns of the fifth Table: Or yet more eaſily thus, the third number in the ſixth Table, is compoſed of the third in the fifth Table and of the ſecond in the ſixth; the fourth number in the ſixth Table is compoſed of the fourth in the fifth and of the third in the ſixth; the like is to be underſtood of the reſt. But you are to obſerve that according to this way of compoſing the ſixth Table by Addition, the numbers of the fifth Table muſt be continued to more places than are there expreſt, to prevent error ariſing by the addition of defective decimal Fractions.

TABLE VI.

Which ſheweth the preſent worth of one Pound Annuity, to continue any term of years under 31, and payable by yearly payments, compound intereſt being computed at any of theſe rates, to wit, 4, 5, 6, 7, 8, 9, 10, 11, *and* 12 per cent. per annum.

Years	4	5	6	7	8	9	10	11	12
1	.96153	.95238	94339	.93457	.92592	.91734	.90909	.90090	.89285
2	1.88609	1.85941	1.83339	1.80801	1.78326	1.75911	1.73553	1.71252	1.69005
3	2.77509	2.72324	2.67301	2.62431	2.57709	2.53129	2.48685	2 44371	2.40183
4	3.62989	3.54595	3.46510	3.3872	3.31212	3.23971	3.16986	3.10244	3.03734
5	4.45182	4.32947	4.21236	4.10010	3.99270	3.88965	3.79078	3.69589	3.60477
6	5.24213	5.07569	4.91732	4.76653	4.62287	4.48591	4.35526	4.23053	4.11140
7	6.00205	5.78637	5.58233	5.38926	5.20636	5.03295	4.86841	4.71219	4.56375
8	6.73274	6.46321	6.20979	5.97129	5.74663	5.53481	5.34492	5.14612	4.96763
9	7.43533	7.10782	6.80169	6.51523	6.24688	5.99524	5.57901	5.53704	5.32824
10	8.11089	7.72173	7.36008	7.02358	6.71008	6.41765	6.14456	5.88923	5.65022
11	8.76047	8.30641	7.88687	7.49867	7.13896	6.80519	6.49506	6.20651	5.93769
12	9.38507	8.86325	8.38384	7.94268	7.53607	7.16072	6.81369	6.49235	6.19437
13	9.98964	9.39357	8.85268	8.35765	7.90377	7.48690	7.10335	6.74987	6.42354
14	10.56312	9.89864	9.29498	8.74546	8.24423	7.78614	7.36668	6.98186	6.62816
15	11.11838	10.37965	9.71224	9.10791	8.55947	8.06068	7.60608	7.19087	6.81086

A continuation of the preceding Table VI.

Years	4	5	6	7	8	9	10	11	12
16	11.65229	10.83776	10.10589	9.44664	8.85136	8.31255	7.82371	7.37916	6.97398
17	12.16560	11.27406	10.47725	9.76322	9.12163	8.54363	8.02155	7.54879	7.11962
18	12.65929	11.68958	10.82760	10.05908	9.37188	8.75562	8.20141	7.70161	7.24966
19	13.13393	12.08531	11.15811	10.33559	9.60359	8.95011	8.36492	7 83929	7.46577
20	13.59032	12 46220	11.46992	10.59401	9.81814	9.12854	8.51356	7.96332	7.46944
21	14.02915	12.82115	11.76407	10.83557	10.01680	9.29224	8 64869	8.07507	7.56200
22	14.45111	13 16300	12.04158	11.06124	10.20074	9.44242	8.77154	8.17574	7.64464
23	14.85683	13.48857	12.30337	11.27218	10.37105	9.58020	8.88322	8.26643	7.71843
24	15.24696	13.79864	12.55035	11.46933	10.52875	9.70661	8.98474	8.34813	7.78431
25	15.62207	14.09394	12.78335	11 65358	10.67477	9 82257	9.07704	8.42174	7 84313
26	15.98276	14.37518	13 00316	11.82577	10.80997	9 92897	9.16094	8.48805	7 89505
27	16.32958	14.64303	13.21053	11.98671	10.93516	10.02657	9.23722	8.54780	7.94255
28	16.66305	14.89812	13.40616	12.13711	11.05107	10.11612	9.30656	8.60162	7 98442
29	16.98371	15.14107	13 59071	12.27767	11.15840	10.19828	9.36960	8.65011	8.02180
20	17.29292	15.39244	13.76482	12.40004	11.25778	10.27365	9.42691	8 69379	8.05516

The Uſe of the preceding Table VI.

The uſe of the ſaid ſixth Table will appear by the manner of ſolving theſe two ſubſequent queſtions, *viz.*

Queſt. 1. What is the preſent worth of an Annuity or rent of 56 *l.* *per annum* payable by yearly payments for 21 years, accompting intereſt upo intereſt at the rate of 6 *per centum, per annum* *Anſw.* 658 *l.* 15 *s.* 9 *d.* very near, thus found out In the fourth Column of the preceding *Table* VI under the figure 6 at the head, and right againſ 21 years, I find 11.76407, which ſhews that ar Annuity of 1 *l.* payable by yearly payment for 2 years, is worth in preſent money 11.76407 *l.* o 11 *l.* 15 *s.* 3 *d.* 1 *f.* and ſomewhat more) intereſ upon intereſt being computed on both ſides at th rate of 6 *per centum, per annum*; therefore multiplying the ſaid tabular number 11.76407 by 56 (56 becauſe the Annuity propounded is 56 pound the product (according to the ſecond rule of th 26th. Chapter of the preceding Book) will b found to be 658.787, &c. that is, 658 *l.* 15 *s.* 9 very near: Wherefore I conclude that the Anſwe of the Queſtion is 658 *l.* 15 *s.* 9 *d.* View the fo lowing Operation.

1 . 11.76407 : : 56 . (658,787 +
56

7058442
5882035

658|78792

Queſt. 2. What is the preſent worth of an An-
ual Rent of 45 *l.* payable by yearly payments for
1 years, intereſt upon intereſt being computed at
0 *per centum, per annum*? *Anſw.* 389 *l.* 3*s.* 10 *d.*
ery near; for in the Column of 10 *per centum*, in
he ſaid ſixth *Table*, right againſt 21 years, and
nder 10, at the head I find this number 8.64869;
hich ſhews that at 10 *per centum*, compound inte-
eſt, an Annuity or Rent of 1 *l.* payable by yearly
ayments for 21 years, is worth in ready money
.64869 *l.* that is 8 *l.* 12 *s.* 11 *d.* 3 *f.* therefore
ultiplying the ſaid tabular number 8.64869, by
5 (the Rent propounded) the Product will be
89.191 +, that is 389 *l.* 3 *s.* 10 *d.* very near,
hich is the Anſwer of the Queſtion.

1 . 8.64869 : : 45 . (389.191 +
45

4324345
3459476

389|19105

In the ſame manner the numbers in the other Columns of *Table* VI. are to be uſed.

To find how many years purchaſe an Annuity or a Leaſe for years is worth.

Moreover the numbers in the ſaid ſixth *Table* will at firſt ſight ſhew how many years purchaſe an Annuity to continue any number of years under 31 is worth, to be ſold for preſent money, compound intereſt being computed on both ſides, at any of the ſaid rates 4, 5, 6, 7, 8, 9, 10, 11 and 12 *per centum*: So if you deſire to know how many years purchaſe an Annuity iſſuing out of Lands for 21 years, to begin preſently, is worth, if it were to be ſold for ready money, when the current rate of intereſt is 6 *per centum*; ſeek in the firſt Column of *Table* VI. for 21 years, and carry your eye from thence equidiſtant to the head-line of the *Table* till you come under 6, which (as before hath been ſaid) ſignifies 6 *per centum*. So in the fourth Column you will find 11.76407, whereof you need only conſider 11.76, which ſhews that the ſaid Annuity is worth 11 years purchaſe, (or 11 times one years rent whatever it be) and 76 parts of one years purchaſe divided into 100 parts, or a $11\frac{1}{4}$ years purchaſe and a little more. The ſame Annuity when money was at 8 *per centum* was worth 10 years purchaſe and about $\frac{1}{100}$ part of a years purchaſe more, as the number in the Column of 10 *per centum* right againſt 21 years will diſcover.

In like manner ſuppoſing 10 *per centum* to be a fit rate to be allowed in the valuation of Leaſes of houſes, the Leaſe of a houſe for 21 years will be found by the ſaid *Table* to be worth 8 years purchaſe and $\frac{64}{100}$ parts of a years purchaſe, or 8 years purchaſe and an half, and half a quarter of a years purchaſe, and ſomewhat more : But note here, that in valuing of Leaſes, the rate *per centum* is to be ſet higher or lower according to the goodneſs of the thing leaſed, and the certainty or uncertainty of the rent.

XVI. When a ſum of money is propounded, and it is required to know what Annuity (to continue any number of years, and according to any given rate) that ſum will buy, you may preſuppoſe at pleaſure an Annuity for the term of years propounded, and find the value of that Annuity in ready money (according to the fifteenth Rule aforegoing) at the rate aſſigned ; then will the proportion be as followeth.

Of the purchaſe of Annuities at compound intereſt.

> *As the value found is in proportion to the ſuppoſed Annuity ; ſo is the ſum of Money propounded, to the Annuity required.*

So if it be required to find what Annuity to begin preſently, and to continue three years 500 *l.* in preſent money will purchaſe, compound intereſt being computed at 6 *per centum, per annum* : The *Anſwer* will be 187 *l.* 1 *s.* 1 *d.* very near.

For preſuppoſing an Annuity at pleaſure, to wit, 378.743088 *l.* payable yearly for 3 years, the value thereof in preſent money will (by the 15th. Rule of this Chapter) be found to be 1012.3848 *l.* Therefore by the *Rule of proportion,* ſay,

1012,3848 : 378,743088 : : 500 . 187,054.

That is to ſay, if 1012.3848 *l.* in ready money will buy an Annuity of 378.743088 *l.* (to continue three years, then 500 *l.* in preſent money will purchaſe an Annuity (to continue the ſame term of years, and at the ſame rate of intereſt) of 187.054, *&c.* that is, 187 *l.* 1 *s.* 1 *d.* very near.

The Conſtruction of the following Table VII.

Upon this ground the following *Table* VII. is calculated to ſhew what Annuity (to continue any term of years under 31, and at any rate of intereſt mentioned at the head of that *Table*) one pound will purchaſe, by which *Table*, and by the help of *Multiplication*, queſtions concerning the *Purchaſe of Annuities*, *Rents* or *Penſions*, by any ſum of ready money propounded, may be reſolved without conſiderable Error. But a more ready way to make the ſaid *Table* VII. may be this following, *viz.*

Foraſmuch as it is evident by the conſtruction of the third *Table* aforegoing, that one pound ready money is equivalent unto 1.06 *l.* payable at the end of a year to come, at the rate of 6 *per centum, per annum*; therefore this 1.06 is to be the firſt number in the Column intitled 6 *per centum* in the ſubſequent *Table* VII. Again, the preſent value of one Pound Annuity to continue two years at the ſaid rate will be found by the preceding *Table* VI. to be near 1.83339 *l.* Therefore by the Rule of Proportion, ſay,

1.83339

1.83339 : 1 :: 1. 54543, &c.

That is, if 1.83339 *l.* ready money will purchaſe an Annuity of 1 *l.* (to continue two years; what Annuity to continue the ſame time will 1 *l.* in preſent money purchaſe? *Anſwer*, an Annuity of .54543 *l.* that is 10 *s.* 11 *d.* very near, to continue two years; therefore the ſaid Decimal .54543 *l.* is to be placed as the ſecond number in the fourth Column of the ſubſequent *Table* VII. Hence it follows, that if 1 or unity be divided by every one of the numbers in all the Columns of *Table* VI. except the firſt Column of years, the quotients will give the reſpective numbers to be placed in the like Columns of the following *Table* VII. in which operation it will be requiſite, that the numbers in the preceding *Table* VI. be continued to more places than are there expreſt, to prevent error that will ariſe by adding of defective Decimals.

TABLE

TABLE VII.

Which ſheweth what Annuity, payable by yearly payments, to continue any term of years under 31, *one pound will purchaſe, compound intereſt, being computed at any of theſe rates, to wit,* 4, 5, 6, 7, 8, 9, 10, 11, *and* 12 per cent. per ann.

Years	4	5	6	7	8	9	10	11	12
1	1.04000	1.05000	1.06000	1.07000	1.08000	1.09000	1.10000	1.11000	1.12000
2	.53019	.53780	.54543	.55309	.56076	.56846	.57619	.58393	.59169
3	.36034	.36720	.37411	.38105	.38803	.39505	.40211	.40921	.41634
4	.27549	.28209	.28859	.29519	.30192	.30866	.31547	.32232	.32923
5	.22462	.23097	.23739	.24389	.25045	.25709	.26379	.27057	.27740
6	.19076	.19701	.20336	.20979	.21631	.22291	.22960	.23637	.24322
7	.16660	.17282	.17913	.18555	.19207	.19869	.20545	.21221	.21911
8	.14852	.15472	.16103	.16746	.17401	.18067	.18744	.19432	.20130
9	.13449	.14069	.14702	.15348	.16007	.16679	.17364	.18060	.18767
10	.12329	.12950	.13586	.14237	.14903	.15582	.16274	.16980	.17698
11	.11414	.12038	.12079	.13335	.14007	.14694	.15396	.16112	.16841
12	.10655	.11282	.11927	.12590	.13269	.13965	.14676	.15402	.16143
13	.10010	.10645	.10296	.11965	.12652	.13356	.14077	.14815	.15567
14	.09466	.10102	.10758	.11434	.12129	.12843	.13574	.14322	.15087
15	.08994	.09634	.10296	.10979	.11682	.12405	.13147	.13906	.14682

A continuation of the preceding Table VII.

Years	4	5	6	7	8	9	10	11	12
16	.08581	.09226	.09895	.10585	.11298	.10209	.12781	.13551	.14339
17	.08219	08869	.09544	.10242	.10962	.11704	.12466	.13247	.14056
18	.07899	.08554	.09235	.09941	.10670	.11421	.12192	.12984	.13793
19	.07613	.08274	.08962	.09675	.10412	.11173	.11954	.12756	.13576
20	.07358	.08024	.08718	.09439	.10184	.10954	.11745	.12557	.13387
21	.07128	.07799	.08500	.09228	.09983	.10761	.11562	.12383	.13224
22	.06919	.07597	.08304	.09040	.09803	.10590	.11400	.12231	.13081
23	.06739	.07413	.08127	.08871	.09642	.10438	.11257	.12097	.12955
24	.06655	.07247	.07967	.08718	.09497	.10302	.11126	.11978	.12846
25	.06401	.07095	.07822	.08581	.09367	.10180	.11016	.11874	.12749
26	.06256	.06956	.07690	.08456	.09250	.10071	.10915	.11781	.12665
27	.06123	.06829	.07569	.08342	.09144	.09973	.10825	.11698	.12590
28	.06001	.06712	.07459	.08239	.09048	.09885	.10745	.11625	.12524
29	.05887	.06604	.07357	.08144	.08961	.09805	.10672	.11565	.12465
30	.05783	.06496	.07264	.08058	.08882	.09733	.10607	.11502	.12414

The uſe of the preceding Table VII.

Queſt. 1. What Annuity or yearly rent iſſuing out of Lands, to begin preſently, and to continue 14 years, will 320 *l.* purchaſe, compound intereſt being reckoned on both ſides, at the rate of 6 *per centum, per annum?* *Anſw.* 34 *l.* 8 *s.* 6 *d.* very near, which is thus found out, *viz.* In the 4th. Column of the preceding *Table* VII. under 6 at the head of that Column, and right againſt 14 years, you will find this Decimal .10758, which ſhews that 1 *l.* ready money will purchaſe an Annuity of .10758 *l.* (that is 2 *s.* 1 *d.* 2 *f.* +) therefore multiplying the ſaid Decimal .10758 by the ſaid 320; the product (according to the ſecond Rule of the 26th. Chapter of the preceding Book) will be found to be 34.425, *&c.* that is 34 *l.* 8 *s.* 6 *d.* very near, which is the *Anſwer* of the *Queſtion.*

1 . ,10758 : : 320 . (34.425 +
320
215160
32274
34|42560

In like manner, if 10 *per centum* be thought a fit rate of intereſt to be allowed in purchaſing Leaſes of Houſes, 500 *l.* will buy a preſent yearly rent of 63 *l.* 18 *s.* 1 *d.* payable for 16 years out of a houſe. For underneath 10 at the head of the 8th. Column, and right againſt 16 years (in the preceding *Table* VII.) you will find this decimal ,12781, which being multiplied by 500, (the number of pounds propounded to purchaſe the Leaſe (the product will be found to be 63.90500, that is, 63 *l.* 18 *s.* 1 *d.* ¼, as by the ſubſequent operation is manifeſt.

1 : ,12781 :: 500 : (63.905
500
—————
63,90500

The making of Tables for half yearly and quarterly payments.

XVII. Upon the ſame foundations which hath been laid in the 12, 13, 14, 15 and 16 Rules of this Chapter, for the making of Tables which reſpect yearly payment, Tables may be made for half yearly and quarterly payments, the intereſt of 100 *l.* for ½ year, and likewiſe for ¼ year being firſt agreed upon: For if we ſuppoſe that at the rate of 6 *l.* for 100 *l.* for a year, the intereſt of 100 *l.* for ½ year is 3 *l.* the numbers 100 and 103 are to be uſed in the ſame manner to calculate *Tables* for half yearly payments, as the numbers 100 and 106 have been before uſed to form *Tables* for yearly payments. But if at the rate of 6 *per centum, per annum,* the intereſt of 100 *l.* for ½ year ought to be ſuch, that being added to the ſaid principal 100 *l.* and the whole put forth at intereſt for the next half year, at the ſaid rate, the ſum then due (to wit, at the years end) muſt exactly amount unto 106 *l.* In this caſe a Geometrical mean proportional number between the extreams 100 and 106 muſt be ſought, which mean will (by the following 18th. Rule) be found to be near 102.956301 +, and then the numbers 100 and 102.956301, *&c.* are to be uſed inſtead of the numbers 100 and 106 in manner aforeſaid. In like manner, if it be ſuppoſed that at the rate of 6 *per centum*, *per annum*, the intereſt of 100*l.* for ¼ year is 1*l.* 10*s.* or 1.5*l.* the numbers 100 and 101,5 are to be uſed for the calculating of Tables for quarterly payments, in the ſame manner as the numbers 100 and 106 for yearly payments. But if at the rate of 6 *per cent. per ann.* the intereſt of 100 *l.* for ¼ year ought to be ſuch, that being added to the ſaid 100*l.* and the whole put forth at the ſame rate of intereſt for the next ¼ of a year, and in that manner for the third and fourth quarters, and that the ſum due at the years end muſt exactly amount unto 106 *l.* In this caſe a ſeries or rank of five numbers in Geometrical proportion continued muſt be conſidered, *viz.* the principal 100 *l.* (which is the leſſer of the two extream proportionals;) the three ſums (compoſed of principal and intereſts) due at the end of the firſt, ſecond and third quarters of the year, (which are the three mean proportionals) and 106 *l.* due at the years end (which is the greater of the two extream proportionals;) now between the ſaid extreams 100 and 106, the firſt (to wit the leaſt) of the ſaid three mean proportionals is to be ſought, which (by the following 20th. Rule of this Chapter) will be found to be near 101.4673+. And then the numbers 100 and 101.4673, *&c.* are to be uſed inſtead of the numbers 100 and 106 in manner aforeſaid.

To find a Geometrical mean proportional number between two numbers given

XVIII. Two numbers being given to find a Geometrical mean proportional between them; multiply the two given numbers one by the other, and extract the ſquare root of the product, ſo is ſuch ſquare root the mean proportional ſought : For Example, if 8 and 18 are two numbers given, and it is required to find a mean number Geometrically proportional between them, multiply 18 by 8, ſo is the product 144, whoſe ſquare root is 12 for the mean proportional ſought, ſo that 8, 12 and 18, are three numbers in Geometrical proportion continued, *viz.* As 8 is in proportion to 12, ſo is 12 to 18. In like manner a Geometrical mean proportional between the extreams 100 and 106 will be found near 102.956301 +.

XIX. Two numbers being given, to find the firſt of two Geometrical mean proportional numbers between the extreams given, multiply the ſquare of the leſſer extream by the greater, and extract the cube root of the product, ſo is ſuch cube root the leſſer of the two mean proportionals required : for example, if 8 and 27 are aſſigned for two extreams, the leſſer mean will be found 12 ; for according to the rule, the ſquare of 8 the leſſer extream is 64, which being multiplied by 27 (the greater extream) produceth 1728, whoſe cube root is 12 the leſſer mean ſought, then may the greater mean be found more eaſily by the[*] *Rule of Three*, for 8 . 12 : : 12 . 18, ſo that 12 and 18 are two means Geometrically proportional between the extreams 8 and 27, *viz.* theſe four numbers are in Geometrical proportion continued, to wit, 8. 12. 18 and 27.

To find the firſt of the Geometrical mean proportional numbers between two extream numbers given.

To find the firſt of three Geometrical mean proportionals between two extream numbers given.

XX. Two numbers being given to find the firſt of three Geometrical mean proportionals between the extreams given, multiply the cube of the leſſer extream by the greater, and extract the Biquadrate root of the product, ſo is ſuch Biquadrate root the firſt (to wit, the leaſt) of the three mean proportionals required: For Example, if 2 and 32 are two extreams given, the firſt and leaſt of three Geometrical mean proportionals will be found to be 4, for (according to *Rule*) the cube of 2 (the leſſer extream given) is 8, which being multiplied by 32 (the greater extream) produceth 256, the *Biquadrate root* whereof being extracted (according to the 29th. *Rule* of the 33th. *Chapter* of the preceding *Treatiſe*) gives 4 for the firſt and leaſt of the three means ſought, the other means may be eaſily found by the *Rule of Three*; for,

2 . 4 : : 4 . 8 : : 8 . 16 : : 16 . 32.

So that theſe five numbers will appear to be in Geometrical proportion continued, to wit,

2 . 4 . 8 . 16 . 32.

In like manner the firſt and leaſt of three *Geometrical mean proportionals* between the extreams 100 and 106, will be found to be near 101.4673, *&c.* Thus have I ſhewed the moſt eaſie ways (raiſed from clear grounds) to make *Tables* for the reſolution of the uſual queſtions, which depend upon the computation of intereſt, by the help of *Multiplication* only.

Queſt.

Queſtions to exerciſe the precedent Tables, with their uſe in ſolving Queſtions of the ſame nature, when the number of years exceeds 30.

Queſt. 1. If the Leaſe of an Houſe be worth 153 *l.* *Fine,* and 16 *l.* *yearly rent,* payable yearly for 21 years, and the *Leſſee* be deſirous to bring down the *Fine* to 50 *l.* and ſo to pay the more *Rent,* the queſtion is, what rent the Tenant ſhall pay accompting compound intereſt at the rate of 10 *per centum,* per *annum?* *Anſw.* 27 *l.* 18 *s.* $1\frac{3}{4}$ *d.* near.

Firſt, find the difference between the *Fines,* which is 103 *l.* Then after the manner of the examples of the uſe of the preceding *Table* VII. ſeek what Annuity or Rent to continue 21 years, 103 *l.* ready money will purchaſe at 10 *per cent.* ſo will you find 11 *l.* 18 *s.* $1\frac{3}{4}$ *d.* which being added to the old rent 16 *l.* gives 27 *l.* 18 *s.* $1\frac{3}{4}$ *d.* which the Tenant muſt pay to the end that the *Fine* may be diminiſhed unto 50 *l.*

Queſt. 2. There is a Leaſe of certain Lands to be let for 14 years for 250 *l.* *Fine,* and 44 *l.* *Rent per annum,* payable yearly, but the Tenant is deſirous to pay leſs *Rent,* *viz.* 20 pounds *per annum,* and to give a greater *Fine;* the queſtion is what *Fine* ought to be paid to bring down the *Rent* to 20 *l.* *per annum,* accompting compound intereſt, at the rate of 6 *per centum, per annum?* *Anſw.* 473 *l.* 1 *s.* 7 *d.*

Firſt find the difference between the *Rents,* which will be 24 pounds *per annum.* Then by the help of the preceding *Table* VI. ſeek what Annuity or Rent of 24 *l.* *per ann.* to continue 14 years, is worth in ready money at 6 *per centum, per annum,* ſo will you find 223 *l.* 1 *s.* 7 *d.* which being added to the firſt *Fine* 250 pounds, gives 473 *l.* 1 *s.* 7 *d.* which the *Tenant* muſt pay, to the end the *Rent* may be brought down to 20 *l. per annum.*

Queſt. 3. There is a Leaſe of certain Lands worth 32 *l. per annum*, more than the Rent paid to the Lord for it, of which Leaſe ſeven years are yet in being, and the Leſſee is deſirous to take a Leaſe in *reverſion* for 21 years, to begin when his old Leaſe is expired, the *Queſtion* is what ſum of money is to be paid for this Leaſe in *reverſion*, accompting compound intereſt at the rate of 6 *per cent. per ann.* *Anſw.* 250 *l.* 7 *s.* 2 *d.*+

Firſt, by adding the 7 years of the Leaſe in being to the 21 years you would have in *reverſion* after thoſe 7 are expired, the ſum is 28. Then by the preceding *Table* VI.

The preſent worth of 1 *l.* Annuity for 28 years at 6 *per centum* compound intereſt, is———	13.40616
Likewiſe the preſent worth of 1 *l.* Annuity for 7 years is———	5.58233
Therefore the difference of thoſe preſent worths, ſhall be the preſent value of 1 *l.* Annuity for 21 years in reverſion after 7 years———	7.82383
Which multiplied by 32 (the yearly rent propounded) gives the *Anſwer* of the queſtion———	250.36256

Otherwiſe thus.

Firſt, By the help of the ſaid *Table* VI. find out how much 32 *l.* yearly rent for 21 years is worth in ready money, as if the 21 years were to begin preſently, at the rate of 6 *per centum*, which ready money will be found 376.45024 *l.* Then by *Table* V. find what 376 45024 *l.* due at the end of 7 years to come, is worth in ready money, ſo will it be 250 *l.* 7 *s.* 2 *d.* which agrees with the *Anſwer* before found.

Queſt. 4. One would beſtow 630 *l.* to purchaſe a preſent yearly rent or Annuity of 60 *l.* to be paid by yearly payments, the queſtion is to know how many years the ſaid Annuity muſt continue, compound intereſt at 6 *per centum*, *per annum*, being allow'd on both ſides. *Anſw.* 17 years, and 23 days, very near.

Firſt, I divide 630 by 60, the quotient is 10.5, which ſhews that 10 years purchaſe and an half are given for the Annuity; then ſearching for 10.5 in *Table* VI. in the Column of 6 *per cent.* I find it not exactly, but the neareſt leſs than it, is 10.47725, ſtanding right againſt 17 years, and the next greater than 10.5 is 10.82760 which is placed againſt 18 years. Whence I infer that the Annuity muſt continue 17 years and more, yet leſs than 18 years. Now the proportional part of a year to be added to 17 years, may be found out near enough for uſe, thus, *viz.* ſubtract the ſaid leſſer tabular number 10.47725 from the greater 10.82760, ſo the remainder will be found .35035: Alſo ſubtracting the ſaid 10.47725 from 10.5 (the quotient

tient firſt found) the remainder will be .02275; then ſay by the rule of three in decimals, as .35035 the greater remainder is to .02275 the leſſer; ſo is 1 year (the difference between 17 and 18 years) to .0649 parts of a year, or 23 days + (as will appear by the fourth *Rule* of the 26th. *Chapter* of the preceding *Book*;) therefore the number of years ſought by the queſtion is 17 years, 23 days.

Queſt. 5. If an Annuity of 96 *l.* payable by yearly payments for 14 years be ſold for 826 *l.* what rate of intereſt *per centum*, is implied in that Bargain? *Anſw.* 7 *l.* 5 *s.* 7½ *d.* near.

Firſt, dividing 826 by 96, the quotient is 8.60146, which ſhews how many years purchaſe was given for the Annuity; then ſearching for 8.60416 in *Table* VI. in a right line paſſing from 14 years, equi-diſtant to the head line of the *Table*, I find it not exactly, but the neareſt leſs than it is 8.24423 (which ſtands in the Column of 8 *per cent.*) and the neareſt greater is 8.74546 (which ſtands in the Column of 7 *per cent.*) whence I infer, that the rate of intereſt required is betwen 7 and 8 *per cent.* and the proportional part of 1 *l.* to be added to 7 *l.* may be found out near enough for practice thus, *viz.* ſubtract the ſaid leſſer tabular number 8.24423 from the greater 8.74546, the remainder will be .50123. Alſo ſubtract 8.60416 (the quotient firſt found, which falls between the ſaid tabular numbers from the ſaid greater tabular number 8.74546,) the remainder will be .14130; then ſay by the *Rule of three* in decimals, as .50123 the great remainder (or difference between the two tabular numbers) is to .14130 the leſſer remainder, ſo is 1 *l.* (the difference between 7 *per cent.* and 8 *per cent.*) to .2819, &c.

or 5*s.* 7*d.* 2*f.* which added to 7*l.* gives 7*l.* 5*s.* 7*d.* 2*f.* which is near the rate of intereſt *per. cent.* required.

Queſt. 6. If a years rent, or one years purchaſe, be paid as a *Fine*, for renewing or adding 7 years to 14 years yet to come of an old Leaſe for 21 years, and accordingly a new Leaſe to be taken for 21 years, to begin preſently (which proportion is ordinarily obſerved by *Biſhops*, *Deans* and *Chapters*, *Heads* and *Fellows* of *Colleges* in letting Leaſes of their Lands) what rate of intereſt *per centum* is implied in that Agreement? *Anſw.* 11*l.* 11*s.* 8*d.* 1*f.* and ſomewhat more.

To ſolve this Queſtion, firſt I ſearch in the preceding *Table* VI. to find out two numbers ſo ſeated in ſome one Column of intereſt, that one of them may ſtand right againſt 14 years, and the other againſt 21 years; and ſo qualified that the difference between them may be exactly 1 or unity; but not finding any two numbers preciſely anſwering thoſe Conditions, I take thoſe numbers that come neareſt, which will be found in the Columns of 11 and 12 *per cent.* for the difference between the numbers 6.98186 and 8.07507, which ſtand in the Column of 11 *per cent.* right againſt 14 years and 21 years, is 1.09321, which exceed 1 (that is 1 years purchaſe) by .09321; alſo the difference between 6.62816 and 7.56200, which ſtand in the Column of 12 *per cent.* right againſt 14 years and 21 years, is .93384, which wants .06616 of 1: therefore I divide 1 *l.* (the difference between 11 *l.* and 12 *l.* *per cent.*) into two parts, in ſuch proportion one to the other, as the ſaid decimals .09321 and .06616 are one to the other; ſo I find the ſaid part of 1 *l.* to be near .5848 and .4151; or 11 *s.* 8 *d.* 1*f.*+ and 8*s.*

5 *d.* 2 *f.* +; the former of which being added to 11 *per centum*, or the latter being ſubtracted from 12 *l. per cent.* gives 11.5848 *l.* or 11 *l.* 11 *s.* 8 *d.* 1 *f.* +, which is very near the rate of intereſt required by the queſtion.

Queſt. 7. What is the preſent worth of 1 *l. per ann.* payable yearly for 10 years, compound intereſt being computed at the rate of 11.5848 *l. per cent. per an.* 5 *l.* 15 *s.* 0 *d* very near, which is found out by the help of the preceding *Table* VI. in this manner, *viz.*

The tabular number for 10 years at 11 *l. per centum* is ——— ——— 5.88923
The tabular number for 10 years at 12 *per centum* is —— —— —— —— 5.65022
Their difference is ——— ——— ——— 0.23901

Then ſay by the *Rule of Three* in decimals, as 1 *l.* (the difference between 11 and 12 *per cent.*) is to .5848 *l.* (to wit, the decimal by which the given rate in the queſtion exceeds 11 *per cent.*) ſo is .23901 (the difference found out as above) to .13977 +, which being ſubtracted from 5.88923 (the greater of the two tabular numbers above-mentioned) there will remain 5.74946, or 5 *l.* 15 *s.* 0 *d.* which is near the preſent worth of one pound yearly rent to continue 10 years, at the propoſed rate of 11.5848 *l. per centum.*

After the ſame manner the preſent worth of 1 *l.* yearly rent payable for 21 years, at the ſame rate of intereſt, will be found to be 7.77503 *l.* or 7 *l.* 15 *s.* 6 *d.* very near, from which if you ſubtract 5.74946, (being the aforementioned preſent worth of 1 *l.* yearly rent for 10 years) there will remain 2.02557 or 2*l.* 0*s.* 6*d.* which is near the preſent worth of a Leaſe of 1*l.* rent *per annum*, for 11 years in reverſion, to begin after 10 years yet to come in a Leaſe are expir'd: Hence it is evident, that if a Tenant to a College hath 10 years yet to come in a Leaſe, at 1*l.* rent *per annum*, and deſires to have 11 years renewed, or added to thoſe 10, and ſo take a new Leaſe for 21 years, to begin preſently at the ſame rent, he muſt give 2*l.* 0*s.* 6*d.* or two years purchaſe and $\frac{1}{40}$ part of a years purchaſe, very near (according to the fundamental proportion before aſſumed in the ſixth queſtion.) The like may be done for any other term of years under 30, by the help of the ſaid *Table* VI.

Concerning the renewing of a College Leaſe of Lands.

But yet by a *Table* calculated purpoſely for the ſaid rate of 11.5848 *l.* *per centum*, (according to the fifteenth *Rule* of this *Chapter*) queſtions of the ſame kind with the two laſt, may be more eaſily anſwered, and therefore (for that they come often in practice) I ſhall here inſert ſuch a *Table*, as I find it ready calculated to my hand by Doctor *Newton*, in his Scale of Intereſt lately publiſh'd, which *Table* is to be uſed in every reſpect like to the preceding *Table* VI. and will be very ready and uſeful, for the proportioning of Fines, in the renewing of Leaſes held from Cathedral Churches and Colleges, as will be manifeſt by the manner of ſolving the two following Queſtions.

Queſt. 8. If a College-Tenant hath 7 years yet to come or unſpent in a Leaſe of Lands for 21 years, at 1 *l.* yearly rent, and deſires to have 14 years renewed or added to thoſe 7 years, and ſo to take a new Leaſe for 21 years to begin preſently, what muſt he pay for a Fine? *Anſ.* 3*l.* 3*s.* 0*d.*

The rule for finding out the Anſwer of the Queſtion propoſed, and ſuch like, is this, *viz.*

From 7.77507 (being the number which anſwers to 21 years in this *Table* VIII.) ſubtract always the tabular number which belongs to the number of years to come or unſpent in the old Leaſe, ſo the remainder will ſhew what Fine muſt be paid for the years to be renewed or added, to make thoſe unſpent years in the old Leaſe to be 21 years compleat again, at 1 *l.* yearly rent.

So to ſolve the queſtion propoſed.

TABLE VIII.

Shewing the preſent worth of one pound Annuity for any number of years under 22, at the rate of 11 *l.* 11 *s.* 8 *d.* 1 $\frac{3}{12}$ *f.* per cent. *compound intereſt.*

Years	*preſent worth*
1	0.90034
2	1.69938
3	2.41922
4	3.06438
5	3.64262
6	4.16088
7	4.62540
8	5.04176
9	5.41496
10	5.74948
11	6.04934
12	6.31819
13	6.55907
14	6.77507
15	6.96868
16	7.14226
17	7.29786
18	7.43737
19	7.56243
20	7.67455
21	7.77507

From

From the preſent worth of 1 *l.* yearly rent for 21 years, which is—	7.77507
Subtract the preſent worth of the ſame rent for 7 years, that were unſpent in the old Leaſe.———	4.62540
And there will remain the Fine ſought, to wit,———	3.14967

That is to ſay, 3.14967 *l.* or 3 *l.* 3 *s.* 0 *d.* (very near) muſt be paid as a Fine, for renewing or adding 14 years to 7 years, that were unſpent in the old Leaſe, the yearly rent being 1 *l.* Alſo the ſaid 3.14967 ſhews, that ſuch a renewal is worth 3 years purchaſe and near $\frac{15}{100}$ parts of a years purchaſe (whatever the rent be.)

Queſt. 9. If a Tenant that hath 17 years yet to come in a Leaſe of Lands held of a College for 21 years, at 50 *l.* yearly rent, be deſirous to renew 4 years, and ſo make thoſe 17 years to be 21 years compleat again at the ſame rent, what muſt he give for a Fine? *Anſw.* 23 *l.* 17 *s.* 2 *d.* 1 *f.* For according to the Rule before given,

From the preſent worth of 1 *l.* yearly rent for 21 years--———	7.77507
Subtract the preſent worth of the ſame rent for 17 years (that were unſpent in the old Leaſe.)———	7.29786
And there will remain,———	0.47721
Which multiplied by the rent———	50
The product will be the Fine ſought, to wit, 23 *l.* 17 *s.* 2 *d.* 1 *f.*	23|86050

Queſtions of this nature may be readily ſolved without the loſs of one ſixteenth part of a years Purchaſe by the help of the following *Table* IX, which I have drawn from the forgoing *Table* VIII. for the benefit of ſuch as underſtand not Decimal fractions: For example, if a College-Tenant deſireth to have 10 years added to 11 years that are to come or unſpent in a Leaſe of Lands that he may have a new Leaſe for the term of 21 years to begin preſently, the following *Table* IX, ſhews that he muſt give for a Fine 1 years Purchaſe, and 2 quarters of a years Purchaſe, and 3 quarters of a quarter of a years Purchaſe, *viz.* one years rent, and half a years rent, and three quarters of a quarter of a years rent: Suppoſing then the rent to be 48*l.* *per annum*, the Fine may be computed thus.

	l.	*s.*	*d.*
One years rent is ————	48	00	00
Half a years rent is ————	24	00	00
Three quarters of a quarter of a years is ————	9	00	00
The ſum is the Fine required	81	00	00

Whence it appears that the Tenant muſt give 81 *l.* as a Fine, for adding of 10 years to 11 years that were unexpired in his old Leaſe, to the end he may have a new Leaſe for 21 years in being.

In like manner the following *Table* IX. ſhews that the Fine for renewing or adding 7 years to 14 years that are unſpent in a Leaſe of Lands, to the end there may be a new Leaſe for 21 years in being, is valued at 1 years Purchaſe preciſely, which is the fundamental proportion aſſumed in calculating the foregoing *Table* VIII, as before was ſaid.

TABLE IX.

TABLE IX.

	Years		*Years*		*Years purchaſe.*	*Quarters of a Year.*	*Quarters of a quarter.*
The Fine for renewing or adding	1	to	20	is valued at	0	0	1
	2	to	19		0	0	3
	3	to	18		0	1	1
	4	to	17		0	1	3
	5	to	16		0	2	2
	6	to	15	is valued at	0	3	0
	7	to	14		1	0	0
	8	to	13		1	0	3
	9	to	12		1	1	3
	10	to	11		1	2	3
	11	to	10	is valued at	2	0	0
	12	to	9		2	1	1
	13	to	8		2	2	3
	14	to	7		3	0	2
	15	to	6		3	2	1
	16	to	5	is valued at	4	0	2
	17	to	4		4	2	3
	18	to	3		5	1	1
	19	to	2		6	0	1
	20	to	1		6	3	2

The like may be done for renewing any other term of years under 21, at any rent propoſed.

Of finding out tabular numbers for any term of years above 30.

But becauſe it may ſometimes happen, that the number of years in queſtions belonging to the preceding 3, 4, 5, 6. and 7 *Tables* may exceed 30, I ſhall by the five following Queſtions ſhew, how by the help of thoſe *Tables* the anſwer to any queſtion of that nature may be found out near the truth, when the term of years is above 30.

Queſt. 10. If 340 *l.* be put forth at 4 *per centum*, compound intereſt, and both principal and intereſt be forborn until the end of 45 years, what will then be due? *Anſw.* 1986 *l.* very near.

To reſolve this queſtion and the like, obſerve this rule, *viz.* Firſt make choice of ſuch numbers of years in *Table* III. that if they be added together will make the number of years propoſed in the queſtion, as 17 and 28. or 15 and 30, each of which pairs make 45, then looking into *Table* III. in the Column belonging to 4 *per centum*, you will find right againſt 17 and 28 years theſe numbers, 1.94790 and 2.99870, which being multiplied one by the other will produce 5.84116+, or 5 *l.* 16 *s.* 10 *d.* which ſhall be the increaſe of 1 *l.* forborn 45 years at 4 *per centum*, compound intereſt; therefore multiplying the ſaid 5.84116 by 340, the product will give 1985.994, *&c.* or 1986 *l.* very near for the Anſwer of the Queſtion.

The reaſon of the ſaid Rule will be manifeſt by this Theorem, *viz.* If there be a rank of numbers in Geometrical proportion continued, beginning with 1 or unity, as 1, 2, 4, 8, 16, 32, 64, 128, *&c.* Alſo if the firſt term 1 be caſt away, and over or under all the reſt of the terms there be placed another rank of numbers, beginning at 1 and proceeding according to the natural order of numbers, as 1, 2, 3, 4, 5, 6, 7, *&c.* which may be called the *Indices* of thoſe in the firſt rank, after the firſt term 1 is caſt away; I ſay if any two of thoſe remaining Geometrical proportionals be multiplied one by the other, the product ſhall be a proportional correſpondent to that *Index*, which is equal to the ſum of the *Indices* anſwering to the two proportionals that were multiplied one by the other.

Proport.	2	4	8	16	32	64	128
Indices.	1	2	3	4	5	6	7

So if 4 and 32, which are the ſecond and fifth proportionals in the upper rank, be multiplied one by the other, the product is 128, which ſhall be the ſeventh proportional, becauſe the ſum of the *Indices* 2 and 5, which Anſwer to the ſaid 4 and 32, is 7. In like manner, becauſe the ſum of the *Indices* 3 and 4 is 7, therefore if the third and fourth proportionals, to wit, 8 and 16, be multiplied one by the other, the product ſhall alſo give the ſeventh proportional 128. Now foraſmuch as the numbers in every one of the Columns, except the firſt Column of years in the preceding *Table* III. are continual proportionals whoſe firſt term is 1, but 'tis excluded out of the ſaid Columns, as appears by the Conſtruction of that *Table*, and for that the numbers of years 1, 2, 3, 4, 5, *&c.* are placed as *Indices* ſhewing the order or ſeat of thoſe proportionals inſerted in the Columns, therefore the rule before given for continuing that Table to any number of years is manifeſt.

Queſt. 11. If one pound be due or payable 50 years hence, what is it worth in ready money, by rebating at 5 *per centum*, *per annum*, compound intereſt? *Anſw.* .08720, *&c.* or 1 *s.* 9 *d.* + which is found out by the help of *Table* V. in the ſame manner as the Anſwer to the laſt Queſtion; (reſpect being had to the ſecond and third rules of the 26th. Chapter of the preceding Book concerning the multiplication of decimal fractions.)

Queſt. 12. If an Annuity of one pound payable yearly for 40 years, be all forborn until the end of that term, what will it then amount unto, compound intereſt being computed at 5 *per centum*, *per annum*? *Anſw.* 120*l.* 16*s.* 0*d.* thus found out: Firſt, according to the ſecond way of calculating the fourth *Table* in the thirteenth *Section* of this *Chapter*, find out a principal, which may have ſuch proportion to the propoſed Annuity 1 *l.* as 100 *l.* hath to 5, ſaying, if 5 *l.* intereſt hath 100 *l.* for a principal, what principal muſt 1 *l.* intereſt have? *Anſwer*, 20 *l.* Secondly, ſeek (after the manner of the preceding tenth queſtion) what 20 *l.* will be augmented unto being forborn 40 years, at the rate of 5 *per centum*, *per annum*, compound intereſt, ſo you will find 140.798+, from which ſubtracting the ſaid principal 20 *l.* the remainder will be 120.798+, or 120 *l.* 16 *s.* which is the anſwer of the queſtion.

Queſt. 13. If an Annuity of one pound payable yearly for 37 years, be to be ſold for preſent money, what is it worth, compound intereſt being computed on both ſides at 6 *per centum*, *per annum?* *Anſwer*, 14 *l.* 14 *s.* 9 *d.* which is found out thus: Firſt, according to the ſecond way of calculating the ſixth *Table* in the fifteenth *Section* of this *Chapter*, find out a principal in ſuch proportion to one pound (the propoſed Annuity) as 100 is to 6, ſo will ſuch principal be found 16.66666+, then after the manner of the preceding eleventh queſtion find out the ready money which is equivalent to 16.66666, due 37 years hence, ſo will ſuch ready money be found to be 1.92988 + (or 1*l.* 18 *s.* 7*d.*) which being ſubtracted from the ſaid principal 16.66666, the remainder will be 14.73678+, or 14 *l.* 14 *s.* 9 *d.* which is the Anſwer of the Queſtion propounded.

Queſt. 14. What Annuity payable by yearly payments to continue 37 years will one pound Purchaſe at 6 *per centum*, *per annum*, compound intereſt? *Anſwer*, 1 *s.* 4 *d.* near, which is found out thus; Firſt, find out the preſent worth of one pound Annuity to continue 37 years, which preſent worth (by the laſt Queſtion) will be found 14.73678 *l.* Then ſay by the *Rule of Three*, if 14.73678 *l.* will purchaſe an Annuity of 1 pound, (to continue 37 years) what Annuity to continue the ſame term will 1*l.* purchaſe? *Anſwer*, .06785 +, or 1*s.* 4 *d.* which is the Anſwer of the Queſtion propounded.

CHAP.

CHAP. VI.

A Demonſtration of the Rule of Three, or Rule of Proportion.

I. FOur numbers are ſaid to be proportionals, when the firſt containeth the ſecond ſo often as the third containeth the fourth; likewiſe when the firſt is ſuch part of the ſecond, as the third is of the fourth; ſo theſe numbers following are called proportionals, *viz.*

$$
\begin{array}{l}
4 \times 6 \,.\, 6 :: 4 \times 9 \,.\, 9 \\
\frac{2}{3} \times 12 \,.\, 12 :: \frac{2}{3} \times 15 \,.\, 15
\end{array}
$$

That is to ſay, 4 times 6 (or 24) is ſaid to have ſuch proportion to 6, as 4 times 9 (or 36) hath to 9. In like manner, $\frac{2}{3}$ of 12 (or 8) hath ſuch proportion to 12; as $\frac{2}{3}$ of 15 (or 10) hath to 15.

II. When four numbers are proportionals, the product ariſing from the multiplication of the two extreams is equal to the product of the two means.

Demonſtration.

By the preceding Definition 1 theſe four numbers are proportionals, *viz.*

$$
\left\{
\begin{array}{l}
4 \times 6 \,.\, 6 :: 4 \times 9 \,.\, 9 \\
B \times C \,.\, C :: B \times D \,.\, D
\end{array}
\right.
$$

The product of the two extreams is —— $\left.\begin{matrix}4 \times 6 \times 9\\ B \times C \times D\end{matrix}\right\}$

The product of the two means is —— $\left.\begin{matrix}6 \times 4 \times 9\\ C \times B \times D\end{matrix}\right\}$

$\text{But}\left\{\begin{matrix}4 \times 6 \times 9\\ B \times C \times D\end{matrix}\right\} = \left\{\begin{matrix}6 \times 4 \times 9\\ C \times B \times D\end{matrix}\right.$

Therefore the Prop. is manifeſt.

Likewiſe.

By the preceding definition theſe four numbers are proportionals, *viz.*

$\frac{2}{3} \times 12 . 12 :: \frac{2}{3} \times 15 : 15$

The product of the two extreams is —— $\left.\vphantom{\frac{2}{3}}\right\} \frac{2}{3} \times 12 \times 15$

The product of the two means is —— $\left.\vphantom{\frac{2}{3}}\right\} 12 \times \frac{2}{3} \times 15$

But $\frac{2}{3} \times 12 \times 15 = 12 \times \frac{2}{3} \times 15$

Wherefore the Propoſition is every way proved.

III. From the laſt Propoſition ariſeth the *Rule of Proportion* commonly called the *Rule of Three,* or *Golden Rule,* which teacheth by three numbers given to find a fourth proportional number in this manner, *viz. Multiply the ſecond and third numbers mutually one by the other, and divide the product by the firſt number; ſo the quotient ſhall be the fourth proportional number ſought, in a direct proportion.* This Rule hath been fully exemplified in the 8th. Chapter of the preceding Book, and the truth of the ſaid Rule may be thus demonſtrated, *viz.* Let there be three numbers given to find a fourth in direct proportion, *viz.* if 24 gives 6, what ſhall 36 give? Or as 24 is in proportion to 6, ſo is 36 to a fourth proportional number ſought, which fourth proportional (whatſoever it be) we may ſuppoſe to be Q, and then theſe four numbers will be proportionals, *viz.*

$$24 . 6 :: 36 . Q$$

Therefore by the ſecond Propoſition of this Chapter.

$$24 \times Q = 6 \times 36$$

And becauſe if equal plane numbers be ſeverally divided by one and the ſame number, the quotients will neceſſarily be equal between themſelves, therefore

$$Q = \frac{6 \times 36}{24}$$

Whereby it is manifeſt that the fourth proportional number is equal to the quotient that ariſeth by dividing the product of the multiplication of the ſecond and third proportionals by the firſt, which was to be proved.

Note, That every *Rule of three inverſe* may be made a *Rule of three direct*, by making the third term the firſt, and by proceeding forward to the other two terms; therefore one and the ſame demonſtration ſerveth for both rules.

CHAP.

CHAP. VII.

A Demonſtration of the Double Rule of Fellowſhip.

THE *Double Rule of Fellowſhip* (commonly called the *Rule of Fellowſhip with time)* preſuppoſeth two things, *viz.* 1. That the particular Stocks of Merchants in Company, have continued unequal ſpaces of time in the common Stock. 2. That at the end of their Partnerſhip, the total gain or loſs is to be divided amongſt them, in ſuch manner, that their ſhares ſhall have ſuch proportion between themſelves, as thoſe ſums of intereſt money have one to another, which at any rate *per centum*, ſimple intereſt only being computed, might be gained by the particular Stocks, within the reſpective times of their continuance in the common Stock: Now for the effecting of ſuch a proportional partition, the ſaid *Double Rule of Fellowſhip* gives this Direction, *viz.* Divide the total gain or loſs into ſuch parts, which ſhall have the ſame proportion one to the other, as is between the products ariſing out of the Multiplication of each particular Stock by its correſpondent time.

For *Example,* Suppoſe Two Merchants *A* and *B* to be Partners in Traffick, for a certain time firſt agreed on between them, and that A doth permit his Stock of 100 *l.* to be implied in their joint Traffick three months, and that B forbears his Stock of 50 *l.* eight months; I ſay (according to the ſaid *Rule of Fellowſhip* with time) what ever the total gain or loſs be, that part thereof which belongs to A muſt have ſuch proportion to the gain or loſs of B, as 100 x 3 (or 300) hath to 50 x 8 or 400.) This Rule hath been fully exemplified in the 13th. Chapter of the preceding Book, and the truth thereof, taking the two premiſed Suppoſitions for granted may be thus demonſtrated.

1. Suppoſing 100 *l.* (the Stock of A) to gain in 3 months any certain ſum of money, as two pounds; I ſeek how much 50 *l.* (the Stock of B) will gain in the ſame time, and at the ſaid rate: ſo I find $\frac{2 \times 50}{100}$ *l.* for,

$$100 : 2 :: 50 : \frac{2 \times 50}{100}$$

2. Having found what 50 *l.* will gain in three months, I ſeek how much the ſaid 50 *l.* will gain in 8 months, at the ſame rate, and ſo I find $\frac{2 \times 50 \times 8}{100 \times 3}$ *l.* for,

$$\frac{3}{1} : \frac{2 \times 50}{100} :: \frac{8}{1} : \frac{2 \times 50 \times 8}{100 \times 3}$$

3. Thus it appears, that if 100 *l.* in 3 months doth gain 2 *l.* then 50 *l.* in 8 months will gain at

the ſame rate $\frac{2 \times 50 \times 8}{100 \times 3}$; ſo that the proportion of the gain of A to the gain of B is,

$$\textit{As 2 is to } \frac{2 \times 50 \times 8}{100 \times 3}$$

4. If both the terms (to wit, the *Antecedent* and *Conſequent*) of the ſaid proportion be ſeverally multiplied by the ſaid *Denominator* 100×3, the products will be in the ſame proportion with the numbers of terms multiplied, (by 17 è 7. *Euclid.*) *viz.* the gain of A will be to the gain of B,

$$\text{As } 2 \times 100 \times 3 \text{ is to } 2 \times 50 \times 8$$

5. Laſtly, Becauſe 2 (the ſuppoſitious gain firſt aſſumed) is a Multiplicator as well in the *Antecedent* as in the *Conſequent* of the laſt mentioned proportion, it may be expung'd out of both, and ſo the gain of A will be to the gain of B in this proportion (which was to be proved) to wit,

$$\text{As } 100 \times 3 \text{ is to } 50 \times 8$$

CHAP VIII.

A Demonſtration of the Rule of Alligation alternate, and the uſe of the ſaid Rule in the Compoſition of Medicines.

I. IN order to the Demonſtration of the ſaid Rule, I ſhall premiſe this *Lemma*, *viz.* if the difference of any two numbers given, be multiplied by a number aſſigned, the product will be equal to the difference between the products which ariſe from the multiplication of thoſe two numbers ſeverally by the number aſſigned.

Suppoſitions.

Two lines or numbers given. $\left\{\begin{array}{l} AC = 10 \\ BC = 4 \end{array}\right.$

Their difference. $AB = 10 - 4$

A multiplicator aſſigned. $\Big\} AD = 5$



Which Suppoſitions, and the *Diagram* being well viewed, the truth of the ſaid *Lemma* will be evident, *viz.*

$$AB \times AD = \overline{AC \times AD} - \overline{BC \times BE}\ (AD)$$

$$\overline{10 - 4} \times 5 = \overline{10 \times 5} - \overline{4 \times 5}$$

II. To add the more light to the following *Demonſtration* of the rule of *Alligation alternate*, I ſhall propound a queſtion which properly belongs to the ſaid rule, *viz.* Suppoſe a *Vintner* having *French Wines* at 5 *d.* the *quart*, and at 10 *d.* the *quart*, would make a Mixture of them in ſuch manner, that he might ſell the mixt Quantity at 7 *d.* the *quart*, and ſo make as much money of the Mixture, as if he ſhould ſell each Quantity of *Wine* at its own price; the queſtion is to know what proportion the quantities of both ſorts of *Wine* in the Mixture muſt bear one to the other. Here, according to the *Rule of Alligation alternate*, I take the difference between the mean price aſſigned for the Mixture, and the two other given prices, and place thoſe differences alternately, *viz.* the difference between 7 and 10 being 3, I write 3 againſt 5; likewiſe 2 being the difference between 7 and 5, I write 2 againſt 10; ſo I conclude, that the quantity to be taken of that ſort of *Wine* of 10 *d.* the *quart*, muſt have ſuch proportion to the quantity of 5 *d.* the *quart*, as 2 to 3. That is to ſay, if 2 *quarts* at 10 *d.* the *quart* be mixed with 3 *quarts* at 5 *d.* the *quart*, the total mixture 5 *quarts* being ſold at 7 *d.* the *quart*, will yield as much money as the ſaid 3 *quarts* at 5 *d.* the *quart*, together with the ſaid 2 *quarts* at 10 *d.* the *quart*; as is evident by the ſubſequent work.

$$7\left\{\begin{array}{c|c}10 & 2\\ 5 & 3\\ \hline & 5\end{array}\right.$$

	quarts	*pence*		*quarts*	*pence*
I.	1.	5	::	3 .	15
II.	1.	10	::	2 .	20
III.	15+20		=	7×5=35	

From the premiſes it appears, that when two things are given to be mixt in ſuch manner as the *Rule of Alligation alternate* requires, the propoſition to be demonſtrated will be this, namely,

Three numbers A. B. C. being given in ſuch ſort that A. is leſs than B. but greater than C. if the difference between A. and B. be multipled by C. and the difference between A. and B. be multiplied by B. the ſum of thoſe products will be equal to the product ariſing from the multiplication of A. by the ſum of the ſaid differences.

Demonſtration.

mean price	*extream prices*	*differences alternate*	
A	B	A—C	BA—BC
	C	B—A	CB—CA
		B—C	BA—CA=B—C x A

The difference between B. and A. is B—A. which multiplied by C. produceth (as is evident by the

Lemma aforegoing in the firſt Section of this Chapter) CB—CA. Alſo the difference between A and C is A—C. which multiplied by B produceth BA—BC. Then the ſum of thoſe two products is BA—CA. (for+CB and —CB expunge one the other) which ſum is manifeſtly the ſame with the product ariſing from the multiplication of A the mean price, by B—C the ſum of the aforeſaid differences (to wit, the ſum of A—C and B—A) for + A and — A expunge one another.

When more than two things of different prices are given to be mixt as aforeſaid, the *Demonſtration* will not be otherwiſe: For if the ſum of every two products ariſing from the multiplication of two alternate differences by their reſpective prices, be equal to the product of the mean price multiplied by the ſum of the ſaid differences; the ſum of all the ſaid products will alſo be equal to the products of the mean price multiplied by the ſum of all the differences; as will clearly appear by view of the ſubſequent work.

$$\begin{array}{rl} \text{If} & \overbrace{D + E}^{\textit{Products of alternate differences multiplied by their reſpective prices.}} = \overbrace{F \; x \; G}^{\textit{Products of the mean price multiplied by the ſeveral ſums of alternate differences.}} \\ \text{and} & H + K = F \; x \; M \\ \text{Then } D + E + & H + K = F \; x \; \overline{G + M} \end{array}$$

More

Moreover, becauſe if equal numbers be ſeverally divided by one and the ſame number, the quotients will be equal between themſelves, therefore from the premiſes this *Corollary* will ariſe.

COROLLARY.

In the *Rule of Alligation alternate*, if the aggregate of the products ariſing from the multiplication of the ſeveral alternate differences by their reſpective prices, be divided by the ſum of the ſaid differences, the quotient will be equal to the main Price. This may be a Proof of any Example of the ſaid *Rule of Alligation.*

OF THE COMPOSITION OF MEDICINES.

See more of this in Mr. J. Dee's *Mathematical Preface, alſo* Tom. 2. *of* P. Herigon *and Maſter* More's *Arithmetick.*

I. Medicines and Simples in reſpect of their Qualities are conſidered in ſome of theſe five ways, *viz,* either as they are hot or cold, moiſt or dry, or as they are temperate; ſo that ſuch Simples or Medicines which work heat in our Bodies are ſaid to be hot, ſuch cold which are cauſe of coldneſs.

II. The mean or middle between the extream Qualities of *Heat* and *Coldneſs*, alſo between *Dryneſs* and *Moiſture*, is called *Temperate* or the *Temperature*;

perature; from which each of the ſaid Qualities *hot*, *cold*, *moiſt* and *dry*, doth differ in four *degrees*, ſo that a Medicine or Simple is ſaid to be either *temperate*, or elſe *hot*, *cold*, *moiſt* or *dry*, in the firſt, ſecond, third, or fourth degree.

III. If the nnmbers 1, 2, 3, 4, 5, 6, 7, 8, 9, be placed as you ſee from A to B, the differences between 5 (the middle number) and the ſuperiour numbers 6, 7, 8, 9, will be 1, 2, 3, 4, which may repreſent the 4 Degrees of the Qualities hot and dry; likewiſe the differences between 5 and the inferiour numbers 4, 3, 2, 1, will be 1, 2, 3, 4, which may repreſent the 4 degrees of the qualities cold and moiſt, the temperature repreſented by 0. being the mean or middle from whence the ſaid degrees do ſwerve.

	Ind.	*Degr.*	
B	9	4	*Qualities hot and dry*
	8	3	
	7	2	
	6	1	
	5	0	*Temperature.*
	4	1	*Qualities cold and moiſt*
	3	2	
	2	3	
A	1	4	
	Ind.	*Degr.*	

IV. Since the *Rule of Alligation alternate* requires that of two things miſcible, the one muſt exceed the mean propounded and the other be leſs, therefore the queſtions of *Alligation* in this kind are to be wrought with the numbers in the aforeſaid Column AB, for by them the degrees and qualities are diſcovered, being placed as you ſee in the Column adjacent to AB, and for diſtinction ſake, thoſe numbers in the ſaid Column AB, may be called the *Indices* or *Exponents* of the *degrees*, which *Indices* are to be uſed in the ſame manner as the prices of Merchandizes in the queſtions of *Alligation alternate* in Chapter 14 of the preceding Book, and therefore thoſe Examples may be compared with theſe.

Prop. I.

Having divers Simples whoſe qualities are known, to make a compoſition of mixture of them, in ſuch manner that the quality of the Medicine may be ſome mean amongſt the qualities of the Simples, and the quantity thereof any quantity aſſigned.

Example 1. An *Apothecary* hath four ſorts of Simples, A, B, C, D, whoſe qualities are as followeth, *viz.* A is hot in the fourth degree, B is hot in the ſecond, C is temperate, and D is cold in the third degree; the queſtion is to know what quantities of each ought to be taken, to make a Medicine, whoſe quantity may be 12 ounces, and the quality in the firſt degree of heat? Seek in the aforeſaid column AB, for the *Indices* or *Exponents* of the qualities of the Simples given, *viz.* for A which is hot in the fourth degree, take 9; for B which is hot in the ſecond, take 7; for C which is temperate, take 5; and for D which is cold in the third degree, take 2; that done, rank thoſe numbers in the ſame manner as the prices of Merchandices in the queſtions of the 14th. *Chapter*, *viz.* deſcend from the higheſt degree of heat unto the temperature, and ſo proceed downwards to the degrees of cold, ſetting 6 the *Index* or *Exponent* of the mean quality propounded, which is 1 degree of heat, as common to them all; then by crocked lines or otherwiſe connect two ſuch *Indices*, whereof one may be greater than the mean, and the other leſs, and proceeding according to the *Rule* of the fourteenth *Chapter* you will find that to make a Medicine of 9 ounces, and the quality reſulting to be in the firſt degree of heat, you muſt take 1 ounce of A (being that Simple which was hot in 4) 4 ounces of B, 3 ounces of C, and 1 ounce of D, as will be manifeſt by the proof,



ounc.	ſimp.	*The Proof.*
1	A	9 x 1 = 9
4	B	7 x 4 = 28
3	C	5 x 3 = 15
1	D	2 x 1 = 2
9		9) 54 (6

Laſtly, by the *rule of Proportion* you may increaſe he Medicine to the quantity of 12 ounces, and yet the quality to continue in the firſt degree of heat, according to the following operation.

Oun.

Oun.		Oun.		Oun.		Oun.	
9	.	1	::	12	.	$1\frac{1}{3}$	of A
9	.	4	::	12	.	$5\frac{1}{3}$	of B
9	.	3	::	12	.	4	of C
9	.	1	::	12	.	$1\frac{1}{3}$	of D

The Quantity aſſigned 12 *Ounces.*

By other connexions of the qualities, other quantities of each *Simple* would ariſe, but that hath been ſufficiently manifeſted in the queſtions of the fourteenth Chapter.

Example 2. Suppoſe there are five *Simples*, A, B, C, D, E, whoſe Qualities are as followeth, *viz.* A is hot in 3°. B is hot in 2°. C is hot in 1°. D is cold in 1°. E is cold in 3°. and it is required to mix 4 Ounces of B, with ſuch Quantities of the reſt that the Quality of the *Medicine* may be temperate?



	Oun.	Simp.	The Proof.
1	1	A	8 x 1 = 8
3	3	B	7 x 3 = 21
1	1	C	6 x 1 = 6
3+1	4	D	4 x 4 = 16
2	2	E	2 x 2 = 4
	11		11)55(5

Proceed as before, ſo will you find that to make a *Medicine* of 11 ounces, and the quality of the *Form* reſulting to be temperate, you muſt take 1 ounce of A, 3 ounces of B, 1 ounce of C, 4 ounces of D, and 2 ounces of E; then ſince the quantity of B, in the compoſition propounded is limited, *viz.* 4 ounces, find numbers which may be in ſuch proportion to 4 (the quantity of B aſſigned) as the numbers 1, 1, 4, 2 (the quantities of A, C, D, E, in the aforeſaid Compoſition of 11 ounces) are unto 3 (the quantity of B in the ſaid Compoſition) in manner following:

Oun. *Oun.* *Oun.* *Oun.*
3 . 1 :: 4 . $1\frac{1}{3}$ of A.
3 . 1 :: 4 . $1\frac{1}{3}$ of C.
3 . 4 :: 4 . $5\frac{1}{3}$ of D.
3 . 2 :: 4 . $2\frac{2}{3}$ of E.

to be mixed with 4 ounces of B.

Prop. II.

A *Medicine* being compounded of divers *Simples* whoſe qualities and quantities are known, to find the degree of the *Form reſulting, viz.* the exact temperament of the *Medicine.*

Example 1. Suppoſe a *Medicine* to be compounded of two *Simples, viz.* 6 ounces of B hot in 4°. and three ounces of C hot in 3°. and it is required, to find the temperament of the *Medicine, viz.* the degree and quality reſulting from ſuch mixture? Seek in the aforeſaid Column A B for the *Indices* of the reſpective degrees and qualities of the *Simples* given, and diſpoſe them orderly in ranks right againſt their reſpective quantities; then multiply each *Index* by its reſpective quantity, and divide the ſum of the products by the ſum of the quantities: So will the quotient be the *Index* of the degree and quality of the Medicine.

Ind. Qun. Prod.

$9 \times 6 = 54$
$8 \times 3 = 24$
— —
$9)\ 78\ (8\frac{2}{3}$

So in the ſaid Example the Quotient will be found $8\frac{2}{3}$, which is the *Index* of $3\frac{2}{3}$ degrees of heat, and therefore the ſaid *Medicine* is hot in $3\frac{2}{3}$ degrees.

Foraſmuch as any two quantities miſcible according to the *Rule of Alligation alternate*, are in ſuch proportion one to the other, as the reſpective alternate differences between the mean quality of the mixture and the qualities correſpondent unto the ſaid quantities, the Demonſtration of the aforeſaid *Rule* will be manifeſt by the *Corollary* aforegoing in this Chapter.

Example 2. Suppoſe a *Medicine* to be compounded of 4 *Simples*, whoſe qualities and quantities are known, *viz.* 2 ounces of A hot in 3°. 3 ounces of B hot in 2°. 4 ounces of C temperate, and 5 ounces of D cold 4°. and let it be required to find the mean quality reſulting from ſuch mixture. According to the aforeſaid *rule*, I multiply each *Index* by its reſpective quantity, and divide the ſum of the products by the ſum of the quantities, ſo the quotient is $4\frac{3}{7}$ which is the *Index* of $\frac{4}{7}$ degrees of cold (for the difference between 5 the *Index* of the temperature, and $4\frac{3}{7}$ the *Index* found, is $\frac{4}{7}$ degrees of cold) which is the quality of the ſaid *Medicine.*

Ind.		*Oun.*		*Prod.*
8	x	2	=	16
7	x	3	=	21
5	x	4	=	20
1	x	5	=	5

14) 62 ($4\frac{3}{7}$

Example 3. Suppoſe a Medicine to be compounded of ſeveral *Simples*, whoſe qualities and quantities are as followeth, *viz.* 4 ounces of a *Simple* which is cold in 2°. and moiſt in 1°. 5 ounces hot in 3°. and (in reſpect of dryneſs and moiſture) temperate; 3 ounces hot in 2°. and dry in 2°. 6 ounces hot in 1°. and moiſt in 4°. 4 ounces cold in 3°. and moiſt in 2°. the queſtion is to know the temper reſulting?

In the reſolution of this queſtion there muſt be two diſtinct operations, each of them like to that in the laſt Example, *viz.*

1. Find in the ſame manner as before, the degree and quality reſulting from the commixture of the qualities hot and cold; ſo will you find $5\frac{7}{22}$ which is the Index of $\frac{7}{22}$ degrees of heat (for the difference between 5 the Index of the temperature and $5\frac{7}{22}$ the Index found, is $\frac{7}{22}$ degrees of heat.)

Ind.		*Own.*		*Prod.*
3	x	4	=	12
8	x	5	=	40
7	x	3	=	21
6	x	6	=	36
2	x	4	=	8

22) 117 ($5\frac{7}{22}$

Ind.		*Own.*		*Prod.*
4	x	4	=	16
5	x	5	=	25
7	x	3	=	21
1	x	6	=	6
3	x	4	=	12

22) 80 ($3\frac{7}{11}$

2. Find in the ſame manner, the temper reſulting from the mixture of the qualities dry and moiſt; ſo will you find $3\frac{7}{11}$ which is the Index of $1\frac{4}{11}$ degree of moiſture, ſo the quality of the ſaid Medicine is $\frac{7}{22}$ degree of heat, and $1\frac{4}{11}$ degree of moiſture, as by the operation is manifeſt.

Prop. III.

To augment or diminiſh a Medicine in quality according to any degree aſſigned.

Suppoſe a Medicine to be compounded as followeth, *viz.* 1 dram of a Simple hot in 4°. 2 drams hot in 3°. 2 drams hot in 2°. 1 dram hot in 1°. 1 dram cold in 1°. and 1 dram cold in 2°. Then will the quality of the ſaid Medicine be in $1\frac{1}{2}$ degree of heat as will be manifeſt by the ſecond Propoſition.) Now let it be required to augment the ſaid Medicine in quality, *viz.* to add ſuch a quantity of ſome one of the Ingredients (or ſome other ſimple) which may raiſe the quality of the Medicine ½ degree; ſo that the temperament of the Medicine after it is increaſed in quantity, may be in 2°. of heat. Make choice of ſuch a ſimple, the Index of whoſe quality may exceed the Index of the quality aſſigned, *viz.* make choice of that ſimple which is hot in 3°. whoſe Index is 8, then proceed according to the 1 Example of the firſt Propoſition; ſo will you find that if 1 dram of the aforeſaid Medicine be mixed with ½ dram of that ſimple which is hot in 3°. the temper reſulting from ſuch mixture will be in 2°. of heat:

Laſtly, by the *Rule of Three*, ſay, if 1 dram require ½ dram, what ſhall 8 drams (the quantity of the Medicine firſt given) require?

Anſw. 4 drams: So that if 4 drams of a ſimple which is hot in 3°. be mixed with 8 drams of a Medicine which is hot in 1 ½ degree, the temper reſulting will be in 2°. of heat, as by the operation is manifeſt.

Ind. Drams.

$$7\begin{cases}6\frac{1}{2}\\ 8\end{cases}\left|\begin{array}{l}1\\ \frac{1}{2}\end{array}\right.$$

$$1 . \frac{1}{2} :: 8 . 4$$

The Proof.

Ind. Dra. Prod.

$$6\frac{1}{2} \times 8 = 52$$
$$8 \times 4 = 32$$
$$\overline{12.)\ 84\ (7}$$

If it be required to diminiſh a Medicine in quality, you are to make choice of ſuch a Simple, the Index of whoſe quality may be leſs than the Index of the quality aſſigned, and then to proceed as before.

Here obſerve, that if in queſtions of this nature, the quantities of the Simples be expreſt by weights of divers denominations, they are to be reduced to that weight which is of the loweſt denomination in the queſtion, according to the ſixth *Rule* of the ſeventh *Chapter* of the preceding *Book*.

The augmenting or diminiſhing of a Medicine in reſpect of quantity; Alſo the finding of the value of any quantity of a Medicine, the prizes of the Ingredients being known, will be familiar to ſuch as underſtand the Rule of Proportion, and therefore I ſhall not inſiſt upon them.

CHAP. IX.

A Demonſtration of the common Rule of Falſe by two Poſitions.

I. WHat the ordinary *double Rule of Falſe* is, and how to be uſed in reſolving ſuch queſtions which cannot readily be applied to any of the other rules of *Arithmetick*, hath been fully declared in the 15th. and 31th. Chapters of the preceding Book; it remaineth to ſhew what kind of operation is preſuppoſed before the ſaid Rule can be applied to the reſolution of a queſtion, and then to demonſtrate the Truth of the Rule it ſelf.

II. In the ſaid *Rule of Falſe*, look what operation the queſtion requires to be performed with the number ſought and ſome given number or numbers, the ſame kind of operation in every reſpect is to be made with each of the two feigned numbers commonly called Poſitions) and the ſaid given number or numbers; which threefold proceſs being finiſht (whether it be by any one, or all of theſe *rules*, to wit, *Addition*, *Subtraction*, *Multiplication*, and *Diviſion)* there will ariſe three remarkable numbers or reſults, to wit, one reſulting from the true number ſought, and two others reſulting from the two feigned numbers; then from theſe three Reſults, the Errors are collected, which are nothing elſe but the differences between the true Reſult, and each of the two falſe Reſults.

III. After the ſaid errors or differences are diſcovered, the *Rule of Falſe* will be of no force, unleſs this Analogy or proportionality doth ariſe, namely the firſt error muſt have the ſame proportion to the ſecond, as the difference between the number ſought and the firſt feigned number hath to the difference between the ſaid number ſought and the ſecond feigned number; here therefore it may be demanded, what kind of operation will produce the ſaid Analogy? To this I anſwer, when the queſtion requires the number ſought to be increaſed, leſſened, multiplied or divided by ſome given number, or the number ariſing from ſuch operation to be increaſed, leſſened, multiplied or divided by ſome given number; in any of thoſe caſes, the aforeſaid Analogy will neceſſarily ariſe, as I ſhall here manifeſt in all the ſaid caſes. Firſt, therefore I ſay when unto each of three numbers (namely the number ſought by the *Rule of Falſe* and the two feigned numbers) one and the ſame number is added, the ſaid Analogy will enſue, for in this caſe the difference between the firſt ſum and the ſecond will be equal to the difference between the firſt and ſecond of the ſaid three numbers; likewiſe the difference between the firſt ſum and the third will be equal to the difference between the firſt number and the third which may be proved in manner following.

Suppoſitions.

Let there be three numbers, to wit,

$$\begin{array}{ccccc} A & . & B & . & C \\ 12 & . & 7 & . & 5 \end{array}$$

Suppoſe alſo that the firſt number A is greater than either of the numbers B and C.

Suppoſe alſo, ſome number as D (3) to be added to each of the ſaid three numbers, ſo will the three ſums be,

$$\begin{array}{l|l} A+D & 15 \\ B+D & 10 \\ C+D & 8 \end{array}$$

The Propoſition to be demonſtrated is, that the difference between the firſt ſum and the ſecond is equal to the difference between the firſt number and the ſecond; alſo that the difference between the firſt ſum and the third is equal to the difference between the firſt number and the third.

Demonſtration.

The difference between the firſt number and the ſecond is,

$$A - B$$

The difference between the firſt ſum and the ſecond is,

$$A+D-B-D$$

But the latter difference is manifeſtly equal to the former (for $+D$ and $-D$ expunge one the other;) to wit,

$$A+D-B-D=A-B$$

Therefore the firſt part of the propoſition is proved.

Again, the difference between the firſt number and the third is,

$$A-C$$

The difference between the firſt ſum and the third is,

$$A+D-C-D$$

But the latter difference is manifeſtly equal to the former, for $+D$ and $-D$ expunge one the other, *viz.*

$$A+D-C-D=A-C$$

Wherefore the propoſition is fully proved.

The like property might be proved after the ſame manner, when one and the ſame number is ſubtracted from three numbers ſeverally.

Secondly, when three numbers (namely the numbers ſought by the *rule of Falſe* and the two feigned numbers) are ſeverally multiplied by one and the ſame number; the afore-mentioned Analogy will likewiſe enſue, as may be thus proved.

Suppoſitions.

Let there be three numbers, to wit,

$$A \,.\, B \,.\, C.$$
$$3 \,.\, 5 \,.\, 8.$$

Suppoſe alſo that the firſt number *A* is leſs than either of the numbers *B* and *C*.

Suppoſe alſo, each of thoſe three numbers to be multiplied by one and the ſame number as *D* (4) and the three products to be theſe,

DA	12
DB	20
DC	32

The Propoſition to be demonſtrated is, that the difference between the firſt product and the ſecond hath ſuch proportion to the difference between the firſt product and the third, as the difference between the firſt number and the ſecond hath to the difference between the firſt number and the third, *viz.*

$$DB-DA \,.\, DC-DA :: B-A \,.\, C-A$$
$$8 \,.\, 20 :: 2 \,.\, 5$$

Demonſtration.

Foraſmuch as (by the 17th. Prop. of the ſeventh Book of *Euclid*'s *Elem.*) if a number (*D*) multiplying two numbers (*B*—*A* and *C*—*A*) produceth other numbers (*DB*—*DA* and *DC*—*DA*) the numbers produced by the multiplication ſhall be in the ſame proportion as the numbers multiplied are, therefore

$$DB-DA \,.\, DC-DA :: B-A \,.\, C-A$$

which was to be demonſtrated.

Likewiſe when 3 numbers are divided by one and the ſame number, the *demonſtration* will not be otherwiſe; and becauſe by the ſecond *Section* of this *Chapter*, the errors in the *Rule of Falſe* are the differences between the true Reſult and the two falſe Reſults, therefore from the precedent *demonſtrations* it is evident, that the aforementioned Analogy or proportionality (namely, when the firſt error hath ſuch proportion to the ſecond, as the difference between the number ſought and the firſt feigned number hath the difference between the ſaid number ſought and the ſecond feigned number) will ſucceed from ſuch operation, as is before declared in the beginning of the third *Section* of this Chapter.

To know whether a queſtion be reſolvable by the Rule of Falſe or not.

IV. Now to diſcern what kind of operation will not produce the ſaid Analogy, obſerve this Note, *viz.* when a queſtion requires ſome given number to be divided by the number ſought or any part thereof, alſo when the number ſought or ſome part thereof is to be ſquared, cubed, *&c.* likewiſe when ſome parts of the number ſought are to be multiplied one by the other; I ſay from ſuch operations the aforementioned Analogy will not ariſe, and in thoſe caſes, the *ordinary Rule of Falſe* will be uſeleſs; as may partly appear by the two following examples, *viz. What number is that, by which if* 360 *be divided the quotient will be* 24? Here if two poſitions or feigned numbers be taken, and 360 be divided by each of them, the errors will not be in the ſame proportion with the differences between the true number ſought and the 2 feigned numbers, and therefore the *rule of Falſe* will be uſed in vain: Yet if it be asked what number is that, which being multiplied by 24, the product will be 360, the *Anſwer* to this latter queſtion is the ſame with the anſwer to the former, and may be found by the *Rule of Falſe*; but ſuch kind of interpretations and inferences are not always obvious, and therefore ſince the preparative work of the *Rule of Falſe* (after the number is taken by gueſs for the number ſought) proceeds gradually from one condition in the queſtion to another, it will for the moſt part be eaſie to determine whether the ordinary *Rule of Falſe* will take place or not, by comparing the conditions of a queſtion with the Note before given.

Another Example; a certain perſon being demanded what number of years he had lived, anſwered, if $\frac{1}{12}$ of that number were multiplied by $\frac{1}{4}$ of the ſame number, the product would ſhew the number, or his age: Here it will be in vain to ſearch the number ſought (which is 48) by the *rule of Falſe*; for the aforementioned Analogy or proportionality will not ſucceed, and the queſtion cannot eaſily be reſolved without *Algebra*.

Now from this ſuppoſition, that after the preparative work of the *rule of Falſe* is finiſh'd, the errors will be in ſuch proportion as aforeſaid, I ſhall make it manifeſt that the *rule of Falſe* will diſcover the number ſought.

V. In the Rule of two falſe Poſitions there are 3 caſes, *viz.* the errors are either both exceſſes and noted with +, or elſe both defects and noted with —, or laſtly, one of the errors is noted with + and the other with —.

In the two firſt caſes the Rule is this, multiply the Poſitions or feigned numbers by the altern errors, *viz.* the firſt Poſition by the ſecond error, the ſecond Poſition by the firſt error, and reſerve thoſe products; then dividing the difference of the ſaid products by the difference of the ſaid errors, the quotient ſhall be the number ſought by the queſtion.

The demonſtration of the ſaid Rule here followeth.

Caſe I. When the errors are both exceſſes, and noted with +.

Suppoſitions.

1. Let ſome number unknown and ſought by the *rule of Falſe* be repreſented by . . . . } A
2. Let the firſt poſition (or feigned number) be . . . . } B
3. And the ſecond feigned number C
4. Suppoſe alſo that B is greater than C, and each of them greater than A.
5. Moreover, ſuppoſe the error of the firſt Poſition to be . . . . } F
6. And the error of the ſecond Poſition to be } G
7. Suppoſe alſo that this Analogy will be found in the ſaid numbers, *viz.*

$$B - A \,.\, C - A :: F \,.\, G$$

8. The Propoſition to be demonſtrated.

$$A = \frac{FC - GB}{F - G}$$

Demon-

Demonſtration.

9. Foraſmuch as by ſuppoſition in 7°.

$$B-A \,.\, C-A :: F \,.\, G$$

10. Therefore by comparing the rectangle of the extreams to the rectangle of the means.

$$GB-GA=FC-FA$$

11. And by equal addition of FA.

$$FA+GB-GA=FC$$

12. Again, foraſmuch as by ſuppoſition in 4°.

$$B \overline{>} C$$

13. And conſequently out of 4°. and 12°.

$$B-A \overline{>} C-A$$

14. Therefore out of 9°. and 13°.

$$F \overline{>} G$$

15. Therefore

$$FA \overline{>} GA$$

16. Therefore

$$FA-GA \overline{>} 0$$

17. Therefore by equal ſubtraction of GB from the equation in 11°.

$$FA-GA=FC-GB$$

18. Wherefore by dividing both parts of the laſt equation by F—G, equal quotients will ariſe, *viz.*

$$A=\frac{FC-GB}{F-G}$$

which was to be demonſtrated.

Caſe II. *When the errors are both defects, and noted with—*

Suppoſitions.

1. Let ſome number unknown and ſought by the *rule of Falſe* be repreſented by .... } A
2. Let the firſt poſition (or feigned number) be .. } B
3. And the ſecond feigned number ... C
4. Suppoſe alſo that B is leſs than C, and each of them leſs than A.
5. Moreover, ſuppoſe the error of the firſt Poſition to be ..... } F
6. And the error of the ſecond Poſition .. G
7. Suppoſe alſo that this Analogy will be found in the ſaid numbers, *viz.*

$$A-B \,.\, A-C :: F \,.\, G$$

8. The Propoſition to be demonſtrated.

$$A = \frac{FC - GB}{F - G}$$

Demonſtration.

9. Foraſmuch as by ſuppoſition in 7°.

$$A - B \,.\, A - C :: F \,.\, G$$

10. Therefore by comparing the rectangle of the means to the rectangle of the extreams.

$$FA - FC - GA - GB$$

11. And by equal addition of FC.

$$FA = FC + GA - GB$$

12. Again, foraſmuch as by ſuppoſition in 4°.

$$B \leq C$$

13. And conſequently out of 4°. and 12°.

$$A - B \leq A - C$$

14. Therefore out of 9°. and 13°.

$$F \leq G$$

15. Therefore

$$FA \leq GA$$

16. Therefore

$$FA-GA < 0$$

17. Therefore by equal ſubtraction of GA from the equation in 11°.

$$FA-GA=FC-GB$$

18. Wherefore by dividing both parts of the laſt equation by F-G, equal quotients will ariſe, *viz.*

$$A=\frac{FC-GB}{F-G}$$

which was to be demonſtrated.

Caſe III. *When one of the errors is an exceſs (to wit, noted by +) and the other a defect (noted by—)*

In this third Caſe the *Rule of Falſe* is this, *viz.* Multiply the Poſitions by the altern errors, to wit, the firſt Poſition by the ſecond error, alſo the ſecond Poſition by the firſt error, and reſerve thoſe products; then dividing the ſum of the ſaid products by the ſum of the ſaid errors, the quotients ſhall be the number ſought by the queſtion.

The Demonſtration of this latter Rule here followeth.

Suppoſitions.

1. Let ſome number unknown and ſought by the *Rule of Falſe* be repreſented by . . . . . A
2. Let the firſt Poſition be . . . . B

3. And the ſecond Poſition C

4 Suppoſe alſo that B is greater than C, and alſo greater than A, and that C is leſs than A.

5. Moreover, ſuppoſe the error of the firſt Poſition to be .... F

6. And the error of the ſecond Poſition to be. G

7. Suppoſe alſo that this Analogy will be found in the ſaid numbers, *viz.*

$$B - A \,.\, A - C :: F - G$$

8. The Propoſition to be demonſtrated.

$$A = \frac{GB + FC}{F + G}$$

Demonſtration.

9. Foraſmuch as by ſuppoſition in 7°.

$$B - A \,.\, A - C :: F \,.\, G$$

10. Therefore by comparing the rectangle of the means to the rectangle of the extreams.

$$FA - FC = GB - GA$$

11. And by equal addition of FC and GA to the laſt equation, this will ariſe,

$$FA + GA = GB + FC$$

12. Wherefore by dividing both parts of the laſt

equation by F + G, equal quotients will riſe, *viz.*

$$A = \frac{GB + FC}{F + G}$$

which was to be demonſtrated.

The learned *Herigonius* (in *cap.* 13. *Tom.* 2. of his *Curſus Mathematicus*) hath delivered another way of reſolving the *rule of Falſe*, namely by the two following rules, *viz.*

When the ſigns of the errors are unlike.

Rule I. As the ſum of the errors is to the firſt error, ſo is the difference of the ſuppoſed numbers to a fourth proportional, which being added to the firſt ſuppoſed number, when the ſaid firſt ſuppoſition is leſs than the ſecond, or ſubtracted from it when it exceeds the ſecond; the ſum or remainder will be the true number ſought.

When the ſigns of the errors are unlike.

Rule II. As the difference of the errors is to the firſt error, ſo is the difference of the ſuppoſed numbers to a fourth proportional, which being added to the firſt ſuppoſed number when the ſigns are — or ſubtracted from it when the ſigns are +; the ſum or remainder will be the number ſought.

Both which rules the ſaid *Herigonius* demonſtrateth geometrically by lines, upon a ſuppoſition of the *Analogy* or *Proportionality* before mentioned in the third Section of this Chapter, and the ſame may likewiſe be eaſily demonſtrated according to the precedent Method by Letters.

CHAP.

CHAP. X.

A Collection of pleaſant and ſubtil Queſtions, to exerciſe all the parts of Vulgar Arithmetick. To which are alſo added various practical Queſtions about the menſuration of Superficial Figures and Solids.

Example of the Rule of Three mixtly uſed with other Rules.

QUeſt. 1. If a wedge of Gold weighing $17\frac{3}{7}$ *lb.* of *Troy Weight* be worth $679\frac{5}{7}$ *lb. ſterling*, what is the value of $1\frac{3}{13}$ grain of that Gold?

Anſw. 2 pence.

I. $1\frac{3}{13}$ (or $\frac{16}{13}$) of $\frac{1}{24}$ of $\frac{1}{20}$ of $\frac{1}{12} = \frac{1}{4680}$
II. $\frac{122}{7} . \frac{4758}{7} :: \frac{1}{4680} . \frac{1}{120}$

Queſt. 2. A Man dying gave to his eldeſt Son $\frac{2}{3}$ of $\frac{1}{4}$ of his eſtate, to his ſecond Son $\frac{1}{5}$ of $\frac{1}{2}$ of his eſtate, and when they had counted their Portions, the one had 40 *l.* more than the other; the remainder of the eſtate was given to the Wife and younger Children. The queſtion is, what was the Portion of the eldeſt Son, alſo of the ſecond, and how much did belong to the Wife and younger Children?

Anſw. The eldeſt Sons Portion 100 *l.* the ſecond Sons Portion 60 *l.* and 440 *l.* for the Wife and younger Children.

The fractions being reduced, it will be manifeſt that the eldeſt Son had $\frac{1}{6}$, *and the ſecond* $\frac{1}{10}$; *alſo the* *difference of the ſaid fraction is* $\frac{1}{15}$, *then ſay,*

$$\frac{1}{15} \cdot \frac{40}{1} :: \frac{1}{10} \cdot \frac{60}{1}$$

	l.
The ſecond Son's portion	60
The difference of their portions	40
The eldeſt Son's portion	100

$$\frac{1}{15} \cdot \frac{40}{1} :: \frac{1}{1} \cdot \frac{600}{1}$$

Laſtly, $600-160=440$ *for the Wife and younger Children.*

Queſt. 3. A young Man received $66\frac{2}{3}$ *l.* which was $\frac{2}{3}$ of $\frac{1}{2}$ of his eldeſt Brother's portion, and $3\frac{1}{2}$ times of his eldeſt Brother's portion, was $1\frac{1}{4}$ times of his Father's Eſtate, the queſtion is, what was the Father's Eſtate? *Anſw.* 560 *l.*

$$\frac{1}{3} \cdot 66\frac{2}{3} :: 1 \cdot 200$$
$$200 \times 3\frac{1}{2} = 700$$
$$1\frac{1}{4} \cdot 700 :: 1 \cdot 560$$

Queſt. 4. If A can finiſh a Work in 20 days, and B in 30 days; in what time will the work be finiſhed by A and B working together? *Anſwer* 12 days.

Firſt find what quantity of the work will be done by each workman in one and the ſame time; then it will be, as the ſum of thoſe quantities is in proportion to the ſaid time, ſo is 1 or the whole work to the time wherein ſuch work will be finiſhed by both Workmen working together.

days *work* *days* *work*
30 . 1 :: 20 . $\frac{2}{3}$
add 1
sum $1\frac{2}{3}$

$20 : 1 :: a : \frac{a}{20}$
$30 : 1 :: a : \frac{a}{30}$
$\frac{a}{20} + \frac{a}{30} = 1$
$3a + 2a = 60$
$5a = 60$
$a = \frac{60}{5} = 12$

Hence it appears that A and B working together 20 days will finiſh that Work once, together with $\frac{2}{3}$ of the ſame Work; therefore ſay again by the *Rule of Three.*

work *days* *work* *days*
$1\frac{2}{3}$. 20 :: 1 . 12

Queſt. 5.

Æreus adſto leo, tubuli mihi lumina bina,
Oſque etiam, dextri ſic quoque planta pedis.
Binis dextro oculo, ternis lacus iſte diebus
Impletur lævo, ſed pede bis geminis.
Ori ſufficiunt ſex horæ. Dic ſimul ergo,
Quo ſpatio os, oculi, peſque replere valent?

The ſenſe is this. A brazen Lion being placed in an artificial Fountain, conveyeth water into a Ciſtern by two ſtreams iſſuing from his Eyes, alſo by one from his mouth, and by another at the bottom of his right foot. Now the Pipes through which theſe ſtreams paſs, are of different Capacities, in ſuch ſort, that by the right eye ſet open alone the reſt of the ſtreams being ſtopt, the Ciſtern will be filled in two days (the length of a day being ſuppoſed to be 12 hours;) by the left eye alone in three days; by the foot alone in four days; and by the mouth alone in ſix hours. The queſtion is, to find in what time the Ciſtern will be filled, if all thoſe ſtreams be ſet open at once?

Anſwer, $\frac{12}{37}$ days,

Days		*Ciſt.*		*Days*		*Ciſt.*
2	.	1	: :	3	.	$1\frac{1}{2}$
4	.	1	: :	3	.	$0\frac{3}{4}$
$\frac{1}{2}$	.	1	: :	3	.	$\frac{6}{1}$
					add	1

The ſum is $9\frac{1}{4}$ *Ciſterns that will be filled in Three Days by all the Four Streams running together: Then ſay by the Rule of Three.*

Ciſt.		*Days*		*Ciſt.*		*Days*
$9\frac{1}{4}$	:	3	: :	1	.	$\frac{12}{37}$

Queſt. 6. A Ciſtern in a certain Conduit is ſupplied with Water by one Pipe of ſuch bigneſs, that if the Cock *A* at the end of the Pipe be ſet open the Ciſtern will be filled in $\frac{1}{2}$ hour; moreover at the bottom of the Ciſtern two other Cocks *B* and *C* are placed, whoſe capacities are ſuch, that by the Cock *B* ſet open alone (all the reſt being ſtopt, the Ciſtern ſuppoſed to be full) will be emptied in $1\frac{3}{7}$ hour; alſo by the Cock *C* ſet open alone the Ciſtern will be emptied in $2\frac{1}{3}$ hour: Now becauſe more Water will be infuſed by the Cocks *A* than can be expelled by both the Cocks *B* and *C* in one and the ſame time; the queſtion is to find in what time the Ciſtern will be filled if all the ſaid Three Cocks be ſet open at once? *Anſw.* $1\frac{9}{61}$ hour.

After the manner of the fourth queſtion of this

Chapter, find how many times the Ciſtern will be emptied in one and the ſame ſpace of time, by the Cocks of *B* and C running together; alſo how much of the Ciſtern will be filled by *A* in the ſame time; then will the difference ſhew how much of the Ciſtern is gained by the filling Cock in the ſaid time: Laſtly, as the Ciſterns or parts gained are in proportion to the correſpondent time; ſo is the whole Ciſtern, to the time wherein it will be gained or filled.

a

$\frac{1}{2}:1::a:\frac{2a}{1}$

$\frac{10}{7}:1::a:\frac{7a}{10}$

$\frac{7}{3}:1::a:\frac{3a}{7}$

$\frac{2a}{1}-\frac{7a}{10}-\frac{3a}{7}=1$

bou. ciſt. bou. ciſt.

I. $2\frac{1}{3}$. 1 : : $1\frac{3}{7}$. ($\frac{30}{49}$ *emptied by* C

add 1 B

ſum $1\frac{30}{49}$ *B* & C in $1\frac{3}{7}$ *bou.*

$140a-49a-30a=7$

bou. ciſt. bou.

II. $\frac{1}{2}$: 1 : : $1\frac{3}{7}$: ($2\frac{6}{7}$ *filled by A*

$61a=70$

$1\frac{12}{49}$ *gained by A*

$a=\frac{70}{61}=1\frac{9}{61}$

ciſt. bou. ciſt. bou.

III. $1\frac{12}{49}$ $1\frac{3}{7}$: : 1 : : $1\frac{9}{61}$.

Queſt. 7. Suppoſe a Dog, a Wolf and a Lion, were to devour a Sheep, and that the Dog could eat up the Sheep in an hour, the Wolf in $\frac{3}{4}$ hour, and the Lion in $\frac{1}{2}$ hour: Now if the Lion begin to eat $\frac{1}{8}$ hour before the other two, and afterwards all Three eat together, the queſtion is, in what time the Sheep would be devoured? *Anſwer*, $\frac{31}{104}$ hour.

bou. ſh. bou. ſh.

I. If $\frac{1}{2}$. 1 : : $\frac{1}{8}$: $\frac{1}{4}$

Thus it appears that $\frac{1}{4}$ of the Sheep would be eaten by the Lion, before the Dog and Wolf began to eat.

II. Proceed according to the fourth queſtion, ſo will you find the remaining $\frac{3}{4}$ to be eaten by them all in $\frac{9}{52}$ hour, which added to $\frac{1}{8}$ gives $\frac{31}{104}$ hour, in which time the Sheep would be dovoured.

Queſt. 8. If $120\frac{1}{3}$ *l.* be to be diſtributed amongſt three Perſons, A. B, C, in ſuch ſort, that as often as A takes 5, B ſhall take 4, and as often as B takes 3, C ſhall take 2; what ſhall be the ſhare of each?

Anſw. A $51\frac{4}{7}$*l.* B $41\frac{9}{35}$*l.* C $27\frac{53}{105}$*l.*

Find three Numbers which may expreſs the proportions of their ſhares, by the *Rule of Three*, or (to avoid fractions) thus,

This is the seventh question

$$\frac{1}{2} : 1 :: \overset{a}{\frac{1}{8}} : \frac{2}{8} = \frac{1}{4}$$

$$1 : 1 :: a : \frac{a}{1}$$

$$\frac{3}{4} : 1 :: a : \frac{4a}{3}$$

$$\frac{1}{2} : 1 :: a : \frac{2a}{1}$$

$$\frac{a}{1} + \frac{4a}{3} + \frac{2a}{1} = 1 - \frac{1}{4} = \frac{3}{4}$$

$$3a + 4a + 6a = \frac{9}{4}$$

$$13a = \frac{9}{4}$$

$$52a = 9$$

$$a = \frac{9}{52} + \frac{1}{8} = \frac{124}{416} = \frac{31}{104}$$

5 4

3 2

15 . 12 . 8

thus found

$5 \times 3 = 15$

$3 \times 4 = 12$

$4 \times 2 = 8$

$$35 \cdot 120\tfrac{1}{3} :: \left\{ \begin{array}{l} 15 \cdot 51\frac{4}{7} \\ 12 \cdot 41\frac{9}{35} \\ 8 \cdot 27\frac{53}{105} \end{array} \right.$$

Queſt. 9. A Governour of a certain Garriſon, being deſirous to know how much money the Port or Paſſage of the Garriſon did amount unto in certain months, made choice of a loyal ſervant, giving him order to receive of every Coachman paſſing with a Coach 4 *d.* of every Horſeman 2 *d.* and of every Footman ½*d.* Now at the years end, the ſervant making his Account to the Governour, giveth him 94 *l.* 15 *s.* 10 *d.* and lets him know that as often as 5 paſſed with Coaches, 9 paſſed on Horſeback; and as often as 6 paſſed on Horſeback, 10 paſſed on foot; the queſtion is how many Coaches, Horſemen, and Footmen paſſed? *Anſw.* 2500 Coaches, 4500 Horſemen, 7500 Footmen.

Find three proportional numbers after the manner of the 8 Queſtion, which will be 5, 9, 15, then proceed as followeth,

d.
5 *Coaches* . . 20
9 *Horſemen* 18
15 *Footmen* . 7½

If 45½ · 22750 : : 5 · 2500
9 · 4500
15 · 7500

Queſt. 10. A Factor would exchange 780 *l. Sterling* for double Ducats Dollars, and French Crowns, the Ducats at 7 *s.* 6 *d.* the piece, the Dollars at 4 *s.* 4*d.* and the French Crowns at 6*s.* the piece, to be in ſuch proportion, that ½ of the number of Ducats may be equal to ⅓ of the number of Dollars, and ¼ of the Dollars equal to 3/16 of the Crowns, the queſtion is, how many pieces of each coin he ſhall receive for his 780 pounds.

Anſw. 600 Ducats, 900 Dollars, 1200 Crowns.

Find Three proportional Numbers (after the manner of the eighth queſtion) which will be 6, 4, 3,

$\frac{1}{2}$ . . . . $\frac{1}{3}$
$\frac{1}{4}$ . . . . $\frac{1}{16}$

$\frac{1}{8}$. $\frac{1}{12}$. $\frac{1}{16}$
6 . 4 . 3

Thus it appears that ſix times the number of Ducats muſt be equal to four times the number of Dollars, alſo equal unto three times the number of Crowns. Then make choice of three numbers to anſwer thoſe proportions, ſuch as are theſe, 2, 3, 4, (for $6 \times 2 = 4 \times 3 = 3 \times 4$) with which numbers proceed as followeth,

l.
2 *ducats* . . . $\frac{3}{4}$
3 *dollars* . $\frac{13}{20}$
4 *crowns* . . $1\frac{1}{5}$

$$\text{ſay if} \ldots 2\tfrac{3}{5} \cdot \overset{l.}{780} :: \left\{ \begin{array}{ll} \overset{l.}{\frac{3}{4}} & \cdot \overset{l.}{225} \\ \frac{13}{20} & \cdot 195 \\ 1\frac{1}{5} & \cdot 360 \end{array} \right.$$

l. *ducat* *l.*
$\frac{3}{8}$. 1 :: 225 . 600 *ducats.*
doll.
$\frac{13}{60}$. 1 :: 195 . 900 *dollars.*
crown
$\frac{3}{10}$. 1 :: 360 . 1200 *crowns.*

Queſt. 11. Twenty Knights, 30 Merchants, 24 Lawyers, and 24 Citizens, ſpent at a Dinner 64 pound, which was divided amongſt them in ſuch manner, that 4 Knights paid as much as 5 Merchants, 10 Merchants as much as 16 Lawyers; and

8 Lawyers as much as 12 Citizens; the Queſtion is, to know the ſum of money paid by all the Knights, alſo by the Merchants, Lawyers and Citizens.

Anſwer, The 20 Knights paid 20 pounds, the 30 Merchants 24 pounds, the 24 Lawyers 12 pounds, and the 24 Citizens 8 pounds.

Find four numbers to expreſs the proportions of their payments by the *Rule of Three*, or (to avoid fractions) in manner following, ſo will the proportional numbers be 4, 5, 8, 12, *viz.* 4 Knights paid as much as 5 Merchants, or 8 Lawyers, or 12 Citizens.

4 5
10 16
8 12

320 . 400 . 640 . 960

$\underbrace{4 \qquad 5}\ \underbrace{8 \qquad 12}$

thus found,

$4 \times 10 \times 8 = 320$
$10 \times 8 \times 5 = 400$
$8 \times 5 \times 16 = 640$
$5 \times 16 \times 12 = 960$

e, y, u, 64
a+30e+24y+24u=
4a = 5e
10e = 16y
8y = 12u
8a = 10e
8a = 16y
16y = 24u
8a = 24u
a = 3u
4a = 5e
a = 2y
6a = 3u+5e+12y

36a + 30e − 12y − 18u =
56a + 12y + 6u = 64
6a − 12y = 0
62a + 6u = 64
2a − 6u = 0
64a = 64
a = 1 = 1
e = $\frac{4}{5}$ = $\frac{4}{5}$
y = $\frac{1}{2}$
u = $\frac{1}{3}$
20 Knights = 20
30 M ---- = 24
24 L ---- = 12
24 C ---- = 8

Then preſuppoſing that a **Knight** is to pay 4 proceed as followeth, *viz.*

Answer

Answer 20 Knights pd 1 -- each ----- 20: 0: 0
30 Merchants pd 0: 16. 0 each ---- 24: 0: 0
24 Lawyers — pd 0: 10: 0 each ——— 12: 0: 0
24 Citizens pd 0: 6. 8 each ——— 8: 0 0
64

20 Knights

	l.
20 *Knights* . . .	4
30 *Merchants* . .	$4\frac{4}{5}$
24 *Lawyers* . . .	$2\frac{2}{5}$
24 *Citizens* . . .	$1\frac{3}{5}$

$$\text{ſay, if } 12\tfrac{4}{5} \;.\; 64 :: \begin{cases} 4 & .\ 20 \\ 4\frac{4}{5} & .\ 24 \\ 2\frac{2}{5} & .\ 12 \\ 1\frac{3}{5} & .\ 8 \end{cases}$$

l. — total 64

Queſt. 12. A certain Man with his Wife did uſually drink out a Veſſel of Beer in 12 days, and the Husband found by often experience, that his Wife being abſent, he drank it out in 20 days; the queſtion is, in how many days the Wife alone could drink it out? *Anſw.* 30 days.

Note, It is to be ſuppoſed that the husband in 12 of the 20 days where he drank alone, did drink as much as in the 12 days wherein he drank with his Wife; hence it followeth, that in the remaining 8 of the ſaid 20 days, he drank as much as his Wife did in 12 days. Therefore by the *Rule of Three* ſay, If 8 give 12, what 20? *Anſw.* 30. View the following form of the Work.

From 20
Subtract 12

Then if 8 . 12 :: 20 . 30

$20 : 1 :: 12 : \frac{12}{20}$
$a : 1 :: 12 : \frac{12}{a}$
$\frac{12}{20} + \frac{12}{a} = 1$
$12a + 240 = 20a$
$8a = 240$
$a = \frac{240}{8} = 30$

Queſt. 13. If a Houſe be to be built by Three Carpenters, A, B, C, working in ſuch ſort, that A, alone will finiſh it in 30 days, B in 40 days and A, B, C, together in 15 days, in what time could C alone build the Houſe? *Anſw.* 120 days.

I. After the manner of the fourth queſtion, (find in what time A and B working together will finiſh the houſe;) *Anſw.* $17\frac{1}{7}$ days.

days *work* *days* *work*
40 . 1 . 30 . $\frac{3}{4}$
add 1

ſum $1\frac{3}{4}$

work *days* *work* *days*
$1\frac{3}{4}$. 30 : : 1 . . . $17\frac{1}{7}$

$30 : 1 :: 15 : \frac{15}{30}$
$40 : 1 :: 15 \quad \frac{15}{40}$
$a : 1 :: 15 \quad \frac{15}{a}$
$\frac{15}{30} + \frac{15}{40} + \frac{15}{a} = 1$
$\frac{60+45}{120} + \frac{15}{a} = 1$
$105a + 1800 = 120a$
$15a = 1800$
$a = \frac{1800}{15} = 120$

II. Suppoſing the work of A and B to be performed by one Perſon, as D, the Houſe will be built by D in $17\frac{1}{7}$ days, but by D and C together in 15 days; Then find (according to the 12th. Queſtion) in what time C will build the ſame. *Anſw.* 120 *days.*

From $17\frac{1}{7}$
Subtract 15.

Then if $2\frac{1}{7}$. 15 : : $17\frac{1}{7}$. 120

The Proof may be wrought according to the Fourth or Fifth Queſtions.

Queſt. 14. Two Travellers A and B perform a Journey to one and the ſame place in this manner, *viz.* A travels 14 miles every day, and had travelled 8 days before B began; upon the ninth day B ſets forward, and travels 22 miles every day;

the queſtion is, to find in what times B ſhall overtake A? *Anſw.* at the end of 14 days.

I. Find how many miles A had travelled before B ſet forward? *Anſw.* 112 miles; For

day		*miles*		*days*		*miles*
1	.	14	::	8	.	112

II. Find how many miles B gains of A in a day; *Anſw.* 8 miles; For,

$$22-14=8$$

	miles		*day*		*miles*		*days*
III. If	8	.	1	::	112	.	14

Queſt. 15. There is an Iſland which is 36 miles in compaſs. Now if at the ſame time, and from the ſame place, two Footmen A and B ſet forward to travel round about the ſaid Iſland, and follow one another in ſuch manner that A travelleth every day 9 miles, and B 7 miles; the queſtion is to find in what ſpace of time they will again meet, alſo how many miles, and how many times about the Iſland each Footman will then have travelled?

Anſw. They will meet at the end of 18 days from their firſt parting; and then A will have travelled 162 miles (or $4\frac{1}{2}$ times the compaſs of the Iſland) and B will have travelled 126 miles (or $3\frac{1}{2}$ the compaſs of the Iſland.)

miles
From ... 9
Subtract 7
——— *day* *miles* *days*
2 . 1 :: 36 . 18

mult 18	*mult.* 18.
by 9	*by* 7
36) 162 (4½	36) 126 (3½

Queſt. 16. Two Footmen A and B depart at the ſame time from *London* towards *York*, travelling at this rate, *viz.* A goeth 8 miles every day, B goeth 1 mile the firſt day, 2 miles the ſecond day, 3 miles the third day, and in that progreſſion he goeth forward, travelling in every following day one mile more than in the preceding day; the queſtion is to know in how many days B will overtake A?

$1 : 8 :: a : 8a$
$a+1 \times \frac{a}{2} = 8a$
$aa+a = 16a$
$a+1 = 16$ so $a = 15$

Anſw. 15 *days.*

To reſolve this and ſuch like queſtions, double 8 (the number of miles which A travelleth daily) which make 16, from which ſubtract 1, the remainder is 15 the number of days ſought.

Queſt. 17. If *Exeter* be diſtant from *London* 140 miles, and that at the ſame time one Footman A departed from *London* towards *Exeter*, travelling every day 8 miles, and another B from *Exeter* towards *London*, travelling every day 6 miles, the queſtion is in how many days they will meet one another, and how many miles each Footman will have then travelled?

Anſwer, They will meet at the end of 10 days, and then A will have travelled 80 miles, and B 60 miles.

a
$1 : 8 :: a : 8a$
$1 : 6 :: a : 6a$
$8a + 6a = 140$
$14a = 140$
$a = 10$ days

add { 8 miles travelled daily by A.
6 miles travelled daily by B.

ſum 14 miles which A and B together did travel daily.

m. da. miles da.
14. 1 :: 140. 10 in which time A and B will meet each other.
$10 \times 8 = 80$ miles travelled by A.
$10 \times 6 = 60$ miles travelled by B.

Queſt. 18. A certain Footman A departed from *London* towards *Lincoln*, and at the ſame time another Footman B departed from *Lincoln* towards *London*; alſo A travelleth every day $2\frac{1}{2}$ miles more than B. Now ſuppoſing thoſe two Cities to be 100 miles diſtant one from the other, and that thoſe two Footmen do meet one another at the end of 8 days after the beginning of their Journies; the queſtion is, how many miles each will have then travelled, as alſo how many miles each travelled daily?

Anſw. A 60 miles, B 40 miles. Alſo A travelled $7\frac{1}{2}$ miles every day, and B 5 miles.

$a, a + \frac{5}{2}$
$1 : a :: 8 : 8a$
$1 : a + \frac{5}{2} :: 8 : 8a+2$
$8a + 8a + 20 = 100$
$16a = 80$
$a = 5$ & $a + 2\frac{1}{2} = 7$

day	*miles*	*days*	*miles*
1 .	$2\frac{1}{2}$ ::	8 .	20

Hence it appears that at the time of their meeting A had travelled 20 miles more than B, which

20 Miles being ſubtracted from 100 miles leave 80 miles, whereof the half is 40 miles which B had travelled, therefore A had travelled 60 miles.

Now to find how many miles each travelled daily, ſay,

day *miles* *days* *miles*
8 . 40 :: 1 . 5

Therefore $\left\{\begin{matrix}A\\B\end{matrix}\right\}$ travelled $\left\{\begin{matrix}7\frac{1}{2}\\5\end{matrix}\right\}$ *miles* daily.

Queſt. 19. There is an Iſland which is 134 miles in compaſs; now at the ſame time, and from the ſame place, two footmen A and B begin a journey round about the ſaid Iſland, but they travel towards contrary parts, at this rate, *viz.* A travelleth 11 miles in every 2 days, and B 17 miles in 3 days, the Queſtion is to find in what ſpace of time A and B will meet one another; and how many miles each will then have travelled?

Anſwer. They will meet at the end of 12 days, and then A will have travelled 66 miles, and B 68 miles.

After the manner of the fourth Queſtion of this Chapter, the time ſought will be found 12 days.

days *miles* *days* *miles*
2 . 11 :: 3 . $16\frac{1}{2}$
add 17
———— *days* *miles* *days*
$33\frac{1}{2}$. 3 :: 134 . 12

The miles travelled by each will be found in this manner.

days *miles* *days*
2 . 11 :: 12 . 66 miles travelled by *A.*
3 . 17 :: 12 . 68 miles travelled by *B.*

Queſt. 20. If a Clock hath two Indices (or Hands) one of which (to wit *A*) is carried twice round the whole Circumference of the Dial in one day; and the other (*B*) once in 30 days, and that both at once ſhewing the ſame point begin to be moved; the Queſtion is, in what time they will be again conjoined?

Anſwer, $\frac{30}{59}$ day or $\frac{12}{59}$ hours.

day *circum.* *days* *circum.*
1 . 2 :: 30 . 60
ſubtract 1

59

Hence it appears, that in 30 days *A* will have run through 60 Circumferences, and *B*, one Circumference only in the ſame time; therefore *A* gains of *B* 59 Circumferences in 30 days, therefore ſay,

circum. *days* *circum.* *day*
59 . 30 :: 1. . $\frac{30}{59}$.

Queſt. 21. If 6 *lb.* of Sugar be equal in value to 7 *lb.* of Raiſins; 5 *lb.* of Raiſins to 2 *lb.* of Almonds; 3 *lb.* of Almonds to 5 *lb.* of Currants; 2 *lb.* of Currants to 18 *d*: how many pence are the value of 3 *lb.* of Sugar? *Anſw.* 21*d.*

$$\text{mult. these contin.}\left\{\begin{array}{lcl} 6S. & = & 7\,R. \\ 5R. & = & 2\,A. \\ 3A. & = & 5\,C. \\ 2C. & = & 18\,d. \\ ?\,d. & = & 3\,S. \end{array}\right\}\text{mult. these contin.}$$

$$180. \quad 3780\ (21$$

Queſt. 22. If 3 dozen pair of Gloves be equal in value to 2 pieces of Ribbon; 3 pieces of Ribbon to 7 dozen of Points; 6 dozen of Points to 2 yards of Flanders-lace; and 3 yards of Flanders lace to 81 ſhilling; how many dozen pair of Gloves may be bought for 28 ſhillings?

Anſw. 2 dozen pair of Gloves.

$$\text{mult. these contin.}\left\{\begin{array}{rcl} 3\,G. & = & 2\,R. \\ 3\,R. & = & 7\,P. \\ 6\,P. & = & 2\,L. \\ 3\,L. & = & 81\,S. \\ 28\,S. & = & ?\,G. \end{array}\right\}\text{mult. these contin.}\ 2268)\ 4536\ (2$$

$$4536 \quad 2268$$

Queſt. 23. Suppoſe a *Greyhound* to be courſing a *Hare*, in ſuch ſort that the *Hare* takes five leaps for every four leaps of the *Greyhound*, and that the *Hare* is one hundred of her own leaps diſtant from the *Greyhound*; now if three of the *Greyhounds* leaps be equal to four leaps of the *Hares*; the queſtion is to know how many leaps the *Greyhound* muſt take before he obtain his prey?

Anſw. 1200 leaps.

I. If $3 \,.\, 4 :: 4 \,.\, 5\frac{1}{3}$.

Thus it appears, that 4 of the *Greyhounds* leaps are equal to $5\frac{1}{3}$ of the *Hares* leaps; and becauſe by the queſtion the *Greyhound* takes 4 leaps for every 5 of the *Hares*, therefore the *Greyhound* in every four of his leaps gains $\frac{1}{3}$ of one of the *Hare's* leaps; therefore ſay by the *Rule of Three*,

II. If $\frac{1}{3} \,.\, 4 :: 100 \,.\, 1200$

Queſt. 24. There is a certain Room whoſe Baſis is a long ſquare, which is in circuit $50\frac{1}{2}$ feet, and the height of the walls or ſides of the Room is $8\frac{1}{4}$ feet; all which walls of the Room except a ſpace taken, out for a Window in the form of a long ſquare whoſe height is five feet, and breadth four feet, are to be furniſhed with Hangings of ell broad ſtuff at 3 *s.* 4 *d.* the yard, the queſtion is to know how much money the ſtuff will coſt.

Anſw. 5 *l.* 17 *s.* $6\frac{2}{9}$*d.*

$50\frac{1}{2} \times 8\frac{1}{4} = 416\frac{5}{8}$ *ſquare feet.*
$5 \times 4 = 20$ *ſubtract.*

$396\frac{5}{8}$

$3\frac{3}{4} \times 3 = 11\frac{1}{4}$ *ſquare feet in one yard of ſtuff.*

feet *d* *feet* *d*
If $11\frac{1}{4} \,.\, 40 :: 396\frac{5}{8} \,.\, 1410\frac{2}{9}$

Queſt. 25. There is a certain Walk which is a long ſquare, whoſe length is 40 yards, and breadth 7 yards, to be paved with Stones, each of which being in form of a long ſquare is 28 inches in length, and 24 inches in breadth; the Queſtion is to know how many ſuch Stones will be requiſite to pave the ſaid Walk?

Anſw. 540.

Inches *Inches*
1440 x 252 = 362880 *ſquare Inches.*
28 x 24 = 672 *ſquare Inches.*
672 . 1 :: 362880 . 540 *Stones.*

Queſt. 26. Suppoſe a piece of *Tapeſtry* to be $5\frac{3}{8}$ yards *Engliſh* in length, and $3\frac{7}{8}$ yards in breadth, the Queſtion is, how many ſquare ells *Flemiſh* are contained in that piece of *Tapeſtry*, when the length of 1 ell *Flemiſh* is equal $\frac{3}{4}$ of a yard *Engliſh*?

Anſw. $37\frac{1}{36}$ ſquare ells *Flemiſh.*

$5\frac{3}{8}$ x $3\frac{7}{8} = \frac{1333}{64}$ *ſquare yards.*

Then becauſe $\frac{9}{16}$ of a ſquare yard is equal to ell ſquare of *Flemiſh* meaſure (for $\frac{3}{4}$ x $\frac{3}{4} = \frac{9}{16}$) ſay,

If $\frac{9}{16}$. 1 :: $\frac{1333}{64}$. $37\frac{1}{36}$.

Queſt. 27. A Workman hath performed a piece of Tiling bearing the form of a long ſquare, whoſe length is 273 feet, 7 inches; and breadth 21 feet 5 inches; now when Tiles are ſold at the rate of 11 *s.* $10\frac{3}{4}$*d.* a 1000, and every ſquare of Tiling conſiſting of 10 feet as well in length as in breadth doth take up 1000 Tiles, what doth the ſaid piece of Tiling amount unto?

Anſw. 34 *l.* 17 *s.* $0\frac{4001}{57600}$ *d.*

I. $273\frac{7}{12} x 21\frac{5}{12} = \frac{843731}{01440}$. *ſquare feet.*

II. $100 . \overset{d.}{142\frac{3}{4}} :: \overset{d.}{\frac{843731}{144}} . 8364\frac{4001}{57600}$

Queſt. 28. A Merchant would beſtow 220 *l.* in Clove, Mace, and Nutmegs, the Cloves being at 5 *s.* the pound; the Mace at 11 *s.* the pound, and the Nutmegs at 6 *s.* the pound; now he would have of each ſort an equal quantity, the queſtion is how many pounds he may have of each ſort?

Anſw. 200 *lb.*

s.
5
11
6

22 . 1 :: 4400 : 200

The Proof.

lb.		*s.*		*l.*
200	at	5	*amounts unto*	50
200	at	11	*amounts unto* ...	110
200	at	6	*amounts unto* ...	60
				220

Queſt. 29. A Factor is to receive a ſum of money, and is offered Dollars at 4 *s.* 4 *d.* which are worth but 4 *s.* 3 *d.* or French Crowns at 6 *s.* $1\frac{1}{2}$ *d.* which are worth but 6 *s.* the queſtion is by which Coin he ſhall ſuſtain the leaſt loſs?

Anſw. the Dollars..

d. *d.* *d.* *d.*
52 : 1 : : $73\frac{1}{2}$. $1\frac{43}{104}$

That is, in receiving the Dollars every 6 *s.* $1\frac{1}{2}$ *d.* loſeth $1\frac{43}{104}$*d.* but in receiving the Crowns 6*s.* $1\frac{1}{2}$*d.* loſeth $1\frac{1}{2}$*d.* which is a greater loſs than $1\frac{43}{104}$*d.*

Queſt. 30. A Butcher agrees with a Graſier, for the feeding of 20 Oxen, during the ſpace of 12 equal months, but at 2 months end, the Butcher adds 5 Oxen more, and $6\frac{3}{5}$ months after that, he added 10 Oxen more, and then it is agreed between them that the Graſier ſhall feed them all, ſo long time as will be equivalent to the keeping of the firſt twenty during 12 months; the queſtion is how long time he ſhall feed them all, after the putting in of the laſt 10?

Anſw. 1 month.

Conſider, that as he receives more Oxen to feed he ought to keep them all the leſs time; therefore work as the queſtion imports by the *Rule of Three Inverſe.*

Mon. *Oxen*
12 20
Oxen 2 5 *Mon.* *Oxen*
If 20 : 10 : : 25 . (8 25
$6\frac{1}{5}$ 10
If 25 . . . . $1\frac{3}{5}$ . . 35 (1 *Mon.*

Examples of the Rule of Fellowſhip.

Queſt. 31. Two Merchants, *viz.* A and B, have entred Company; A puts in 500 *l.* and at 4 months end takes out a certain ſum, leaving the remainder to continue 8 months longer, B puts in 250 *l.* and at five months end puts in three hundred pounds more, and then his whole ſum continues ſeven months longer. Now at the making of their Accompt A findeth that he hath gained 106$\frac{2}{3}$ pounds, and B gained 133$\frac{1}{3}$ pounds; the queſtion is to know how much A took out of the Bank at 4 months end?

Anſw. 240 *l.*

250 x 5 = 1250
add 300

550 x 7 = 3850

5100
133$\frac{1}{3}$. 5100 :: 106$\frac{2}{3}$. 4080
500 x 4 = 2000 (*ſubtract*

8) 2080 (260

Laſtly, 500—260=240 *taken out by* A.

The Proof.

l. *mon.*
500 x 4 = 2000
Subtract 240

260 x 8 = 2080

4080

Queſt. 32. Five Merchants, *viz.* *A*, *B*, C, *D*, and *E*, have gained 2025 *l.* which they divide in ſuch ſort that $\frac{1}{2}$ of the ſhare of *A* is equal ſeverally to $\frac{1}{4}$ of the ſhare of *B*, $\frac{1}{5}$ of C, $\frac{1}{6}$ of *D*, $\frac{1}{8}$ of *E*. The queſtion is, what was the ſhare of each Merchant?

Anſw. *A* 162*l.* *B* 324*l.* C 405*l.* *D* 486*l.* *E* 648 *l.*

Divide a number at pleaſure into ſuch parts which may be in ſuch proportion as the ſhares required, and proceed according to the ſubſequent operation.

A 2
B 4
C 5
D 6
E 8

$$
\text{If } 25 \;.\; 2025 :: \begin{cases} 2 \ (162 \textit{ for } A, \textit{ whereof } \frac{1}{2} \textit{ is } 81 \\ 4 \ (324 \textit{ for } B, \textit{ whereof } \frac{1}{4} \textit{ is } 81 \\ 5 \ (405 \textit{ for } C, \textit{ whereof } \frac{1}{5} \textit{ is } 81 \\ 6 \ (486 \textit{ for } D, \textit{ whereof } \frac{1}{6} \textit{ is } 81 \\ 8 \ (648 \textit{ for } E, \textit{ whereof } \frac{1}{8} \textit{ is } 81 \end{cases}
$$

l. 2025

Queſt. 33. Two Merchants A and B are in Company, the ſum of their Stocks is 300 *l.* the money of A continuing in Company 9 months, the money of B 11 months, they gain 200 *l.* which they divide equally, the queſtion is to know how much each Merchant did put in?

Anſw. A 165 *l.* B 135 *l.*

Divide 300 into two ſuch parts which may be in proportion as 11 to 9, ſo will the greater part be the Stock of A, and the leſſer the Stock of B, which Stocks being multiplied by their reſpective times, the products will be equal.

$$\begin{array}{l} 11 \\ \underline{\;\;9\;\;} \\ 20 : 300 :: \left\{ \begin{array}{l} 11 \,.\, 165 \text{ for A} \\ \\ 9 : 135 \text{ for B} \end{array} \right. \end{array}$$

Queſt. 34. Two Merchants, *viz.* *A* and *B*, are in Company, *A* did put in 325 *l.* more than *B*, and the ſtock of *A* continued in Company $7\frac{1}{2}$ months, *B* put in a certain ſum which is unknown, and it continued in Company $10\frac{3}{4}$ months: After a certain time they divided the Gain equally; the queſtion is, what each Merchant did put in?

Anſw. *B* 750 *l.* and *A* 1075 *l.*

Divide the product of the difference of their Stocks multiplied by the time of *A*, by the difference of their times, ſo will the quotient be the Stock of *B*. which added to 325*l.* gives the Stock of *A*.

$$\begin{array}{l} 325 \times 7\frac{1}{2} = 2437\frac{1}{2} \\ 3\frac{1}{4}) \; 2437\frac{1}{2} \;\; (750 \textit{ ſtock of B} \\ \qquad\qquad\quad \underline{\textit{add } 325} \\ \qquad\qquad\quad 1075 \textit{ ſtock of A} \end{array}$$

Queſt. 35. A Goldſmith hath ſome Gold of 24 Carects, others of 22 Carects, and another ſort of 18 Carects fine; he would ſo mix theſe together that the maſs mixed might be 60 *lb.* and that the whole Mixture might bear 20 Carects fine. How much of each ſort muſt he take?

Examples of the Rule of Alligation, How the fineneſs of gold and ſilver is eſtimated, v. p. 111.

lb.

Anſw. 12 *of* 24 *Carects.*
12 *of* 22 *Carects.*
36 *of* 18 *Carects.*

$$20\left\{\begin{array}{l}24\\22\\18\end{array}\right.\quad\begin{array}{c|c}2. & 2\\2 & 2\\4+2 & 6\end{array}$$

$$10 \,.\, 60 :: \left\{\begin{array}{l}2 \,.\, 12\\2 \,.\, 12\\6 \,.\, 36\end{array}\right.$$

Note, Some may think that Queſtions of *Alligation* are capable only of ſo many ſeveral Anſwers as there are different ways to connect the mean Rate or Price with the extream Rates or Prices; yet it is moſt certain, that any ordinary Queſtion of *Alligation*, where three or more things are propounded to be mixt in ſuch manner as that Rule requires, is capable of infinite Anſwers, if Fractions be admitted, and ſometimes of many Anſwers in whole numbers, which are not diſcoverable by the common Rule of *Alligation*: So albeit to the laſt mentioned queſtion, the ſaid Rule of *Alligation* can find but one Anſwer onely, which is before given; yet there are eight other Anſwers in whole Numbers, which are theſe that follow (the Invention whereof I have ſhewn in the 19th. Queſtion of the thirteenth Chapter of my ſecond Book of *The Elements of Algebra.*)

Of 24 *Carects*	18	16	14	10
Of 22 *Carects*	3	6	9	15
Of 18 *Carects*	39	38	37	35

Of 24 *Carects*	8	6	4	2
Of 22 *Carects*	18	21	24	27
Of 18 *Carects*	34	33	32	31

See Chap. 8. *of this Appendix.*

Queſt. 36. An Apothecary hath ſeveral Simples, *viz.* A hot in 3°. B hot 2°. C temperate, D cold in 2°. and E cold in 4°. Now he deſires to make a Medicine of thoſe Simples in ſuch ſort that the temper thereof in reſpect of quality may be in 1°. of heat, and the quantity 8½ Drams; the Demand is what quantity of each Simple he muſt take?

Anſw. 4½ Drams of A, ½ Dram of B, ½ Dram of C, 1 Dram of D, and 1 Dram of E.



		Drams	
8	1, 3, 5	9	A.
7	1	1	B.
5	2, 1	3	C.
3	2	2	D.
1	2	2	E.
		17	

Drams.

17 . 8½ ::	9 .	4½	A.
	1 .	0½	B.
	3 .	1½	C.
	2 .	1	D.
	2 .	1	E.
		8½	

Examples of the Rule of Falſe Poſition.

Queſt. 37. A Merchant buyeth 2 ſorts of Cloaths, *viz*. of blacks and of whites for 68 *l.* 2 *s.* after the rate of 21 *s.* the yard for the black, and 12 *s.* the yard for the white, and he taketh ſo much of each ſort, that $\frac{5}{6}$ of the number of yards of the black, are equal to $\frac{7}{8}$ of the white; the demand is how many yards be bought of each ſort?

Anſw. 42 yards of black, and 40 yards of white.

Queſt. 38. A certain Perſon A payeth unto the uſe of B for ever 2500 *l.* in preſent money, upon this Condition, that B ſhall pay unto A an Annuity of yearly rent to be continued four years, the equality of their agreement being thus grounded, *viz*. the ſaid 2500 *l.* is ſuppoſed to be put forth at intereſt for a year (to commence from the time of their agreement) at the rate of 8 *per centum, per annum*. Then from the ſum of that principal and intereſt (ariſing due at the years end) the firſt payment of the Annuity being ſubtracted, the remiander is likewiſe ſuppoſed to be put forth at the ſame rate of intereſt for the ſecond year; then from the compoſed of this principal and intereſt (due at the ſecond years end) the ſecond payment of the Annuity being ſubtracted, the remainder is likewiſe ſuppoſed to be put forth at the ſame rate of intereſt for the third year; then from this principal and intereſt the third payment of the Annuity being ſubtracted, the remainder is in like manner ſuppoſed to be put forth at the ſame rate of intereſt for the fourth Year: Laſtly, from this principal and intereſt the fourth and laſt payment of the Annuity being ſubtracted, there muſt be nothing left: The Queſtion is, what ſum of money muſt be yearly paid to ſatisfie thoſe Conditions?

Anſw. $754\frac{14117}{17602}$*l.* as will be manifeſt by the ſubſequent proof.

I: $100 \,.\, 108 :: 2500 \,.\, 2700$
Subtract the firſt payment $754\frac{14117}{17602}$

$1945\frac{3485}{17602}$
II. $100 \,.\, 108 :: 1945\frac{3485}{17602} \,.\, 2100\frac{14325}{17602}$
Subtract the ſecond payment $754\frac{14117}{17602}$

$1346\frac{208}{17602}$
III. $100 : 108 :: 1346\frac{208}{17602} \,.\, 1453\frac{12194}{17602}$
Subtract the third payment $754\frac{14117}{17602}$

$698\frac{15679}{17602}$
IV. $100 \,.\, 108 :: 697\frac{15679}{17602} \,.\, 754\frac{14117}{17602}$
Subtract the laſt payment . . . . $754\frac{14117}{17602}$

000

Queſt. 39.

Mulæ, Aſinæque duos imponit ſervulus utres
Impletos vino; ſegnemque ut vidit Aſellam
Pondere defeſſam veſtigia figere tarda,
Mula rogat; quid chara parens cunctare, gemiſque?
Unam ex utre tuo menſuram ſi mihi reddas,
Duplum oneris tunc ipſa feram; ſed ſi tibi tradam
Unam menſuram, fient æqualia utrique
Pondera, menſuras dic docte Geometer iſtas?

The ſence is this. A *Mule* and an *Aſs* carried two unequal quantities of Wine, each conſiſting of a certain number of meaſures, in ſuch ſort, that if the *Aſs* imparted one of her meaſures to the *Mule*, then the *Mules* number of meaſures ſo increaſed would be the double of thoſe which the *Aſs* had remaining; but if the *Mule* gave one meaſure to the *Aſs*, then the *Aſſes* meaſures with that increaſe would be equal to the *Mules* remaining meaſures. The Queſtion is, how many Meaſures each carried?

Anſw. The *Mule* 7, and the *Aſs* 5,

Queſt. 40.

Æs, ferrum, ſtannum miſcens, aurique metallum,
Sexaginta minas penſantem finge coronam.
Æs aurumque duos ſimul efficiunto trientes.
Ternos quadrantes ſtanno mixtum impleat aurum:
At totidem quintas auri vis addita ferro.
Ergo age dic fulvi quantum tibi conjicis auri
Miſcendum: dic quantum æris ſtannique requiras:
Dic quoque ſufficiant duri quot pondera ferri:
Præſcriptam ut valeas rite efformare coronam.

The ſenſe is this, Suppoſe a Crown that ſhall weigh 60 *l.* is to be made of Gold, Braſs, Iron, and Tin, mixed together in ſuch proportion, that the weight of the Gold and of the Braſs together may be 40 *l.* the joint weight of the Gold and of the Tin 45 *lb.* and the joint weight of the Gold and of the Iron 36 *lb.* The Queſtion is how much of every one of thoſe four Metals muſt be taken?

$a+e+y+u=60$
$a+e=40$
$a+u=45$
$a+y=36$
$3a+e+u+y=121$
$e+y+u=60-a$
$e+u+y=121-3a$
$60-a=121-3a$

Anſwer, *l.*
30½ *of Gold,* $27-a$
9½ *of Braſs,* $4-a$
5½ *of Iron,* $3-a$
14½ *of Tin.*

$a=30\frac{1}{2}$
$e=9\frac{1}{2}$
$y=5\frac{1}{2}$
$u=14\frac{1}{2}$
60

Queſt. 41. One being demanded what was the preſent hour of the day, anſwered, That the time then paſt from noon was equal to $\frac{1}{5}$ of $\frac{3}{8}$ of the time remaining until midnight. The queſtion is, what a Clock it was? (ſuppoſing the time between noon and midnight to be divided into twelve equal parts or hours)

Anſw. $\frac{36}{43}$ hour after Noon.

Queſt. 42. A Factor delivers 6 French Crowns and 2 Dollars for 45 ſhillings ſterling; alſo at another time he delivers 9 French Crowns and 5 Dollars (at the ſame rate with the former) for 76 ſhillings. The queſtion is to know the value of a French Crown, alſo of a Dollar?

Anſw. A Crown was valued at 6 *s.* 1 *d.* and a Dollar at 4 *s.* 3 *d.*

Queſt. 43. A certain Uſurer received 36 Dollars for the ſimple intereſt of 186 *l.* lent for a certain time unknown; alſo he received 90 Dollars for the gain of 360 *l.* at the ſame rate of intereſt for a certain time unknown; now the ſum of the months wherein both the ſaid numbers of Dollars were gained was twenty months. The queſtion is to know in what time as well the 36 Dollars as the 90 Dollars were gained?

Anſw. The 36 Dollars were gained in $8\frac{8}{11}$ months, and the 90 Dollars in $11\frac{3}{11}$ months, as may be proved by the *Double Rule of Three.*

Which anſwer may be diſcovered by the following Canon found out by the *Algebraick Art.*

Multiply the Dollars firſt gained, the latter Principal, and the given time, according to the rule of continual Multiplication, for a dividend; then multiply the firſt principal by the Dollars laſt gained;

alſo multiply the latter principal by the Dollars firſt gained, and reſerve the ſum of theſe two laſt products for a Diviſor: Laſtly, divide the Dividend firſt found by the ſaid Diviſor, ſo ſhall the quotient be the time wherein the firſt number of Dollars was gained, which ſubtracted from the time given in the queſtion diſcovers the time wherein the latter number of Dollars was gained.

$$\frac{36 \times 360 \times 20 = 259200}{186 \times 90, + 300 \times 36, = 29700} = 8\tfrac{8}{11}$$

And conſequently .... $20 - 8\tfrac{8}{11} = 11\tfrac{3}{11}$

Examples of the Extraction of Roots.

Queſt. 44. 3481 Soldiers are to be placed in a ſquare Battel, how many are to be ſet in Rank or in File?

Anſw. 59 (for the ſquare root of 3481 is 59.)

Queſt. 45. If 4050 Soldiers are to be ſet in Battel in a figure, which beareth the form of a long ſquare in ſuch manner, that the number in File may be to the number in Rank, as 1 to 2, how many Soldiers are to be placed in Rank, and how many in File?

Anſw. 90 in Rank and 45 in File (found by this Canon or general Rule) *viz.*

As the greater term of the proportion given is to the leſſer, ſo is the number of Men to be placed in Battle to a fourth proportional, whoſe ſquare root is the leſſer number ſought (whether it be for the Rank or File:) Alſo as the leſſer term of the given proportion is to the greater; ſo is the number of Men to be ſet in Battle to a fourth proportional, whoſe ſquare root is the greater number ſought (whether it be for the Rank or File.)

I. | 2. 1 : : 4050 . 2025
II. | √q. 2025 = 45 (*Men in File.*
III. | 1. 2 : : 4050 . 8100
IV. | √q. 8100=90 (*Men in Rank.*

The Proof.

45 x 98=4050
Alſo 45 . 90 : : 1 . 2

Or when one of the numbers ſought (whether it be for the Rank or File) is found, the other may be diſcovered by *Diviſion, viz.*

45) 4050 (90
90) 4050 (45

Queſt. 46. Suppoſe the Wall of a Garriſon to be in height 21 feet, and the breadth of the Moat ſurrounding the ſaid Wall to be 28 feet; the Queſtion is, what length muſt a ſcaling Ladder have to reach from the outermoſt ſide of the Moat to the top of the Wall?

Anſw. 35. (to wit, the ſquare root of the ſum of the ſquares of 21 and 28.)

21 x 21=441
28 x 28=784

√q. 1225 35

Queſt. 47. If 100 *l.* being put forth for intereſt at a certain rate, will at the end of two years be augmented unto $112\frac{36}{100}$*l.* (compound intereſt, or intereſt upon intereſt being computed) what principal and intereſt will be due at the firſt years end?

Anſw. 106 *l.* compoſed of 100 *l.* principal and 6 *l.* intereſt (which 106 is a mean Geometrically proportional between 100 and 112. 36 (and may be found by the eighteenth rule of the fifth Chapter of this Appendix.)

$$100 \times 112.36 = \dot{1}1\dot{2}3\dot{6}\ (106$$

Queſt. 48. If 100 *l.* being put forth for intereſt at a certain rate, will at the end of 3 years be augmented unto 115.7625 *l.* (compound intereſt being computed) what principal and intereſt will be due at the firſt years end?

Anſw. 105 *l.* compoſed of 100 *l.* Principal, and 5 *l.* intereſt) which 105 is the firſt of two mean proportional numbers between 100 and 115,7625 *l.* (See the nineteenth rule of the fifth Chapter of this *Appendix.*)

Various Practical Queſtions to exerciſe Decimal Arithmetick, in the menſuration of Superficial Figures and Solids.

Queſt. 49. If the ſide of a ſquare Superficies be 3 feet, what is the Area or Content of that Superficies? Or (which is the ſame thing) how many Squares, each of which is a Foot ſquare, are contained in that Superficies?

See the ſecond Section of the 23d. Chapter of the preceding Book.

Anſw. 9 ſquare feet, which content is found out by multiplying the given ſide 3 by it ſelf, *viz.* 3 multiplied by 3 produceth 9.

In like manner, if the ſide of a ſquare Pavement of ſtone be 15.7 feet, the ſuperficial content of that Pavement will be 246.49 feet, that is 246 feet and an half very near, (for 15.7 multiplied by it ſelf produceth 246. 49.)

Likewiſe, a ſquare piece of Wainſcot whoſe ſide is 3. 24 yards, will be found to contain 10. 49 + yards, or 10 yards and an half almoſt; for 3. 24 multiplied by it ſelf, to wit, by 3.24 will produce 10. 49 +

Alſo if the ſide of a ſquare piece of Land be 37.25 perches, the content in ſquare perches (neglecting the fraction in the product) will be found 1387, which being reduced (according to the ſeventh *Table* in *Rule* 4, *Chapter* 7 of the preceding *Book)* will give 8 Acres, 2 Roods, and 27 Perches for the content of that ſquare piece of Land.

Queſt. 50. If a long ſquare be 8 feet in length and 5 feet in breadth, what is the ſuperficial content?

Anſw. 40 feet; which content is found out by multiplying the length by the breadth, *viz.* 8 multiplied by 5 produceth 40. So if one of the Lights of a Glaſs window ſuppoſed to be in the form of a long ſquare, hath for its length 3.06 feet, and breadth 1.47 feet, the content of that Glaſs will be 4.4982 feet, or 4 feet, and an half almoſt, (for 3.06 multiplied by 1. 47 produceth 4.4982.)

In like manner if there be a piece of Wainſcot, Plaſtring, or any other ſuperfices in the form of a long ſquare, which is in length 6.325 yards and in breadth 3.214 yards; the ſuperficial content will be found 20.32+ yards, that is 20 yards, one quarter of a yard, and ſomewhat more, for, 6.325 multiplied by 3.214 produceth 20 32+.

Likewiſe a piece of Tiling in the form of a long ſquare whoſe length is 18.5 feet, and breadth 11.7 feet will be found to contain 216.45 ſquare feet, which will be reduced to 2.1645 ſquares of Tiling by allowing (according to Cuſtom) 100 ſquare feet to one ſquare of Tiling.

Alſo if a piece of Land in the form of a long ſquare be 48.75 perches in length, and 36.25 in breadth, the Area or Content in Perches, will be found 1767.18+, which 1767 perches being reduced will give 11 acres and 7 perches for the content of that piece of Ground.

Queſt. 51. If it be required to ſet forth in a Meadow one Acre of Graſs to lie in the faſhion of a long ſquare, and that the length thereof be limited or agreed to be 20 perches, what muſt the breadth be?

Anſw. 8 perches, which breadth is found out by dividing 160 (the number of ſquare perches contained in an acre) by the given length 20. If two acres were required, then 320 (to wit, twice 160) muſt be divided by the given ſide, whether it be the length or breadth; ſo if 7.25 perches be preſcribed for the breadth of two acres, the length muſt be 44. 13 + perches.

In like manner, if the breadth of a Board be 1. 32 foot, and it be demanded how far one ought to meaſure along the ſide thereof to have a ſuperficial foot, or a foot ſquare of that Board; divide

1 by the given breadth, ſo you will find in the quotient this decimal fraction .757+, which repreſents three quarters of a foot or nine inches and ſomewhat more, and ſo much in length ought to be meaſured along the ſide of that Board to make a ſuperficial Foot. Likewiſe if the breadth of a bord be given in inches, then 144 (the number of ſquare inches contained in a ſuperficial foot ſquare) being divided by the given breadth, the quotient will ſhew how many inches ought to be meaſured along the ſide of that board to make a ſuperficial foot; ſo the breadth of a board being 9 inches, the length forward to make a ſuperficial foot will be found 16 inches.

Queſt. 52. If the three ſides of a piece of Land that lies in the form of a triangle be 15 perches, 14 perches, and 13 perches, what is the Area or number of ſquare perches contained in that triangle?

Anſw. 84 Perches, or half an Acre and four Perches, which Content is found out by this Rule, *viz.*

From half the ſum of the three ſides of any plain triangle ſubtract each of the three ſides ſeverally, and note the three remainders; then multiply the ſaid half ſum and thoſe three remainders one into the other (according to the rule of continual Multiplication;) that done, extract the ſquare root of the laſt product, ſo ſhall ſuch ſquare root be the Area or Content of the triangle.

	Perches
The 3 ſides of a triangle	15 14 13
The ſum of the 3 ſides	42
The half of that ſum	21
The 3 remainders found out by ſubtracting each ſide from the half ſum	6 7 8
The product ariſing from the continual multiplication of the four laſt numbers	7056
The ſquare root of which product is the content required, to wit,	84

Another Example.

	Perches
The 3 ſides of a triangle	120 . 5 112 . 6 90 . 3
The ſume of the 3 ſides	323 . 4
The half of that ſum	161 . 7
The 3 remainders found by ſubtracting each ſide from the half ſum	41 . 2 49 . 1 71 . 4
The product ariſing from the continual multiplication of the four laſt numbers	23355380 . 1096
The ſquare root of that product	4832 . 7+

Wherefore I conclude that the content of a plain triangle, whoſe three ſides are 120.5 perches 112.6 perches, and 90.3 perches, is 4832.7 + perches, which reduced give 30 acres and 32 perches (the fraction of a perch being neglected.)

Now foraſmuch as every irregular piece of Ground may be divided into triangles, for a four-ſided Field will be divided into two triangles by one imaginary ſtreight Line leading overthwart from corner to corner called a *Diagonal Line* ; a five-ſided Field into three triangles by two *Diagonals* ; a ſix-ſided Ground into four trangles by three *Diagonals*, &c. the Rule before given will be of excellent Uſe to find out the Contents of large Fields, eſpecially if the Land be of a dear value, as alſo when any controverſie ariſeth by the reaſon of the different admeaſurements of Surveyors of Land: For if the ſides of thoſe triangles be meaſured in the Fields, and their lengths be agreed on, all Artiſts to whom the reaſon of the rule before given is known, will agree in one and the ſame content. But yet this way of meaſuring preſuppoſeth that there is no Obſtacle, as Water, Wood, or other Impediment, to hinder the meaſuring of the ſides of thoſe Triangles in to which the Field is divided as aforeſaid.

Queſt. 53. If the Diameter of a Circle be 28.25, what is the Circumference?

Anſw. 88.749. + : For as 113 is in proportion to 355; or as 1 is to 3. 14159, ſo is the Diameter to the Circumference : Therefore multiplying always the Diameter given by the ſaid 3. 14159 the product ſhall be the Circumference required.

Queſt. 54. If the Diameter of a Circle be 28.25, what is the ſuperficial Content of that Circle?

Anſw. 626.79 +; For as 1 is in proportion to 0.785 (.78539) ſo is the ſquare of the Diameter to the ſuperficial content. Therefore multiplying always the ſaid decimal Fraction .78539 by the ſquare of the given Diameter (which ſquare is the product of the multiplication of the Diameter by it ſelf) the product ſhall be the ſuperficial content required.

Queſt. 55. If the Diameter of a Circle be 28.25: what is the ſide of a ſquare which may be inſcribed within the ſame Circle?

Anſw. 19.975 +; For the ſquare root of half the ſquare of the diameter, or the ſquare root of the double of the ſquare of the Semidiameter, ſhall be the ſide of the inſcribed ſquare ſought. Otherwiſe, as 1 is to .707106, ſo is the diameter to the ſide required. Therefore if you multiply (always) the ſaid .707106, by the diameter given, the product will be the ſide of the inſcribed ſquare required.

Queſt. 56. If the Circumference of a Circle be 88.75 what is the Diameter?

Anſw. 28.249 +; For as 355 is to 113, or as 1 is to .318309, ſo is the Circumference to the Diameter. Therefore if .318309 be multiplied always by the given circumference, the product ſhall be the Diameter required.

Queſt. 57. If the circumference of a Circle be 88.75, what is the ſuperficial content of that Circle?

Anſw. 626.801 +; For as 1 is to .079578, ſo is the ſquare of the circumference to the ſuperficial content. Therefore if .079578 be always multiplied by the ſquare of the given circumference, the product ſhall be the ſuperficial content ſought.

Queſt. 58. If the circumference of a Circle be 88.75. what is the ſide of a ſquare that may be inſcribed within the ſame Circle?

Anſw. 19.975+; for as 1 is to .225078, ſo is the Circumference to the ſide required. Therefore if .225078 be always multiplied by the Circumference given, the product will be the ſide of the inſcribed ſquare ſought.

Queſt. 59. If the ſuperficial Content of a Circle be 626.8, what is the diameter?

Anſw. 28.25+; for as 1 is to 1.27324, ſo is the content to the ſquare of the diameter. Therefore multiplying always 1.27324 by the given content, the ſquare root of that product ſhall be the diameter required.

Queſt. 60. If the ſuperficial content of a Circle be 626.8, what is the circumference?

Anſw. 88.75+; for as 1 is to 12.5664, ſo is the content to the ſquare of the circumference. Therefore if 12.5664 be always multiplied by the given content, the ſquare root of the product ſhall be the circumference required.

Queſt. 61. If the ſuperficial content of a Circle be 626.8, what is the ſide of a ſquare equal to the ſame Circle?

Anſw. 25.035+; for the ſquare root of the given content is the ſide of the ſquare required.

Queſt. 62. If the ſide of a Cube be 12 inches, how many cubical inches are contained in that Cube?

Anſw. 1728; what a Cube is may be well repreſented by a Dye, which is a little cube it ſelf being a rectangular or ſquare ſolid, that hath an equal length, breadth and depth, and is comprehended under ſix equal ſquares; now if the ſide of one of thoſe equal ſquares (which is alſo the ſide of the Cube) be 12 inches, the ſuperficial content of that ſquare will be 144 ſquare inches (for according to the preceding 49th queſtion, 12 multiplied by 12 produceth 144) which multiplied by the depth 12 inches, produceth 1728 culical inches, and ſuch is the ſolid content of that Cube whoſe ſide is 12 inches: So that by one foot of timber or ſtone in whatſoever kind of ſolid it be found, is underſtood a Cube, containing 1728 cubical or Dye-ſquare inches, and conſequently half a foot ſolid contains 864 cubick inches, and a quarter of a food ſolid contains 432 cubick inches.

In like manner, if a ſide of a Cube of ſtone be 2.53 feet, the ſolid content of that Cube will be found 16.194+ feet, for 253 being multiplied by it ſelf produceth 6.4009 ſuperficial feet, which product being multiplied by the ſaid 2.53 will produce 16.194+ ſolid feet.

Alſo if the ſide of a Cube of ſtone or wood be 6 inches, or .5 foot, the ſolid content will be found 216 cubick inches, or .125 parts of a foot ſolid (for 6 multiplied cubically produceth 216, likewiſe .5 multiplied cubically produceth .125;) whence it may be inferr'd, that 8 little cubes of ſtone or wood, each of which is half a foot or 6 Inches ſquare, are contained in a foot of ſtone or timber; for 8 times 216 produceth 1728 (being the number of cubick inches contained in a foot ſolid) likewiſe 8 times .125 produceth 1 (to wit, one entire foot ſolid.)

Queſt. 63. If the breadth of a ſquared piece of timber, ſuppoſed to be ſtreight and terminated at both ends by two equal ſquares, be 1. 55 foot, the depth alſo 1. 55 foot, and the length 17.33 feet, how many cubick feet are contained in that piece of Timber?

Anſw. 41.635 feet, that is, 41 feet and an half, and about half a quarter of a Foot. Which ſolid content is found out by this rule, *viz.* multiply the breadth 1. 55 by the depth 1. 55 the product will be 2. 4025 ſuperficial Feet, which is the content of the Baſe (that is, the Area of either of the two equal ſquares at the ends of the piece;) Laſtly, multiplying the ſaid Baſe 2.4025 by the length 17.33 the product will be 41.635 +, which is the ſolid content required.

In like manner if the breadth of a ſquared piece of Timber, ſuppoſed to be ſtreight and terminated at both ends by two equal long ſquares (which are called the Baſes) be 2.34 feet, the depth 1. 61 foot, and the length 17. 58 feet, the ſolid content will be 66. 23 + feet; for (as before) multiplying the breadth by the depth, and that product by the length, the laſt product ſhall be the ſolid content required.

Queſt. 64. If the breadth, as alſo the depth of a ſquared piece of Timber having equal ſquare Baſes, be 1.55 foot, how far ought one to meaſure along the length of that piece of Timber to make a foot ſolid?

Anſw. 416 parts of a foot, or 5 inches very near; which decimal is thus found, *viz.* Firſt find the ſuperficial content of the Baſe, which will be 2.4025 (for 1.55 multiplied by 1. 55 produceth 2.5025; Then dividing 1 (to wit 1 ſolid foot) by the Baſe 2.4025 the quotient will be .416 +

or $\frac{416}{1000}$ parts of a foot, or five inches almoſt, and ſo far ought to be meaſured along the length of the piece to make a foot ſolid. In like manner, if the breadth be 2. 34. feet, and the depth 1.61 feet, the length forward along the piece to make one ſolid foot will be found .265 parts of a foot, or three inches and almoſt $\frac{1}{5}$ part of an inch.

Queſt. 65. If a ſtreight ſquared piece of timber be terminated by unequal Baſes, whereof one contains 1.92 ſuperficial foot, the other .85 foot, and the length of that piece of Timber be 17.4 feet; what is the ſolid content, or how many cubical Feet are contained in that piece of Timber?

Anſw. 23.474 + feet (found out by one of Mr. *Oughtred*'s Rules for meaſuring a ſegment of a Pyramid in *Problem* 21. *Chapter* 19. of his *Clavis Methemat.*) The Rule is this.

Multiply the greater Baſe by the leſs, and extract the ſquare root of that product, then multiply the ſum of the two Baſes and that ſquare root by one third part of the length of the ſolid propounded, ſo ſhall the laſt product be the ſolid content required.

Example

Example.

The greater Baſe	1 . 92
The leſſer Baſe	0 . 85
The product of the multiplication of thoſe two Baſes	1 . 6320
The ſquare root of that product	1 . 2774
The ſum of that ſquare root and the two Baſes	4 . 0474
One third part of the length is	5 . 8
The product of the multiplication of the two laſt numbers is the ſolid content required	23 . 474+

Queſt. 66. A Pyramid is a ſolid comprehended under plain ſurfaces, and from a triangular, quadrangular, or any multangular Baſe, diminiſheth equally leſs and leſs till it finiſh in a point at the top; now if the ſuperficial content of the Baſe of a Pyramid be 5.756 Feet, and the height thereof 14. 25 feet (which height is the length of the perpendicular line that falleth from the top of the Pyramid to the Baſe) what is the ſolid content of that Pyramid?

Anſw. 27. 341 + feet; for if the Area of the Baſe of a Pyramid, be multiplied by one third part of the height thereof, the product ſhall be the ſolid content of the Pyramid; therefore 5. 756 x 4.75 =27. 341 feet=, the ſolidity of the Pyramid propounded.

Note, If a Pyramid be cut into two ſegments by a Plane parallel to the Baſe, one of thoſe ſegments will be a Pyramid, and the other will have two unequal Baſes, for the meaſuring of which latter ſegment; a rule hath been already given in the ſixty-fifth queſtion, the Area of each Baſe being known.

Queſt. 67. A Cone is a ſolid, which hath a Circle for its Baſe, from whence it grows equally leſs and leſs (like a round Steeple of a Church) till it finiſh in a point at the top; now if the Area of the Baſe of a Cone be 5.756 feet, and the height thereof be 14.25 feet, what is the ſolid content of that Cone?

Anſw. 27.341 feet; for if the Area of the Baſe of a Cone be multiplied by one third part of the height thereof, the product ſhall be the ſolid content of the Cone.

Note, If a Cone be cut into two ſegments by a Plane parallel to the Baſe, one of thoſe ſegments will be a Cone, and the other ſegment will have two unequal Baſes which are Circles; the ſolidity of which latter ſegment may be found out by the rule before given in the 65 queſtion, the Area of each Baſe (or Circle) being known.

Queſt. 68. A Cylinder is a ſolid which may be well repreſented by a Stone roll, ſuch as are uſed in Gardens for the rolling of Walks. Now if the circumference of a Cylinder be 4.57 feet, and the length 3.25 feet, what is the ſolid content of that Cylinder?

Anſw. 5.4 + Feet, thus found out: Firſt by the help of the given circumference 4.57, find out the ſuperficial content of that Circle (being the Baſe of the Cylinder) which content (by the preceding 57th. queſtion) will be found 1.6619 + feet; then multiplying the ſaid 1.6619 by the given length 3.25, the product will be 5.4008 which is the ſolid content required.

Queſt. 69. If the Baſe of a Cylinder be 1.6619 foot, how much in length of that Cylinder will make a foot ſoild?

Anſw. .601 parts of a foot; for 1 (to wit, 1 ſolid foot) being divided by the Baſe 1.6619, gives in the quotient the decimal .601 + for the length required.

Queſt. 70. A Globe is a perfect round body contained under one Plane; in the middle of the Globe there is a point called the Centre, from whence all ſtreight Lines drawn to the outſide are of equal length, and called Semidiameters, the double of any one of which is equal to the Diameter of the Globe; now if the Diameter of a Globe of Stone be 1.75 feet, how many Feet ſolid are contained in that Globe?

Anſw. 2.807 + feet, for as 21 is in proportion to 11, or as 1 is to .5238, ſo is the Cube of the Diameter to the ſolid content of the Globe: Therefore, multiplying always the Cube of the Diameter by the ſaid decimal .5238, the product ſhall be the ſolid content required: So the Diameter 1.75 being firſt multiplied by it ſelf, the product will be 3.0625, which multiplied by the ſaid 1.75, gives in the product 5.359375, to wit, the Cube of the diameter, which being multiplied by .5238, the product thence ariſing will be 2.807. +, which is the ſolidity of the Globe propounded.

Queſt. 71. What is the Diameter of a Globe of Stone, which contains 4 cubical or ſolid Feet?

Anſw. 1.96 + foot, for as 11 is in proportion to 21, or as 1 is to 1.9090909 ſo is 4 (the ſolid content given) to a fourth proportional, to wit, 7.636363+ whoſe cubick root is 1.96 + the diameter required.

Concerning the gauging of Veſſels.

The eaſieſt and apteſt way for practice in gaging, are thoſe which are perform'd by the help of Tables, or Gaging-rods purpoſely compos'd: Nevertheleſs, to give the Reader of this Treatiſe ſome light in this matter, I ſhall here inſert one Rule to find out the number of Gallons contained in a full Tun, Pipe, Hogſhead, Barrel, or ſuch like Veſſel according to Mr. *Wingate*'s way of reducing a Veſſel to a Cylinder.

The Rule is this;

Having found the difference of the two diameters at the Bung and Head of the Veſſel, take $\frac{7}{10}$ of that difference and add it to the leſſer diameter; then ſquare that ſum and reſerve the product; that done, if the content be required in Wine Gallons multiply the product reſerved, this decimal fraction .0034, and the length of the Veſſel, one into the other (according to the *Rule of continual Multiplication*) ſo ſhall the laſt product be the number of Wine Gallons required: but if the content be required in Ale Gallons, multiply the product before reſerved, this decimal fraction .0027, and the length of the Veſſel, one into the other continually, ſo ſhall the laſt product be the content in Ale Gallons: This Rule I ſhall firſt explain by two queſtions, and then ſhew how it is raiſed.

Queſt. 72. If the diameter at the Bung of a Veſſel be 32 inches, the diameter at the Head 28.2 inches, and the length 39 inches (which dimenſions are ſaid to agree very near with thoſe of an Engliſh Veſſel called a Pipe) what is the Content of that Veſſel in Wine Gallons?

Anſw. 126.278 Wine Gallons, that is 126 Wine Gallons and about a quart more (found out by the Rule above given, as will be manifeſt by the following Operation.

Explication.

The diameter at the Bung	32 . 0
The diameter at the Head	28 . 2
The difference	3 . 8
Which multiplied by $\frac{7}{10}$, that is	0 . 7
The product will be	2 . 66
Which added to the leſſer diameter gives the mean diameter	30 . 86
Which mean diameter being ſquared (that is, multiplied by it ſelf) produceth	952 . 3396
Which product multiplied by	0 . 0034
The product thence ariſing will be	3 . 2379+
Which multiplied by the length of the Veſſel	39 . 0
The product is the number of Wine Gallons ſought, *viz.*	126. 278+

Queſt. 73. If the Diameter at the Bung of a Barrel be 23 inches, the Diameter at the Head 19.9 inches, and the length 27.4 inches; what is the content of that Barrel in Ale Gallons?

Anſw. 36.031 Ale Gallons, that is 36 Gallons and about a quarter of a Pint more (found out by the preceding Rule.)

Explication

Explication.

The diameter at the Bung ———— 23 . 0
The diameter at the Head ———— 19 . 9
Their difference ———— 3 . 1
Which multiplied by $\frac{7}{10}$, that is ———— 0 . 7
The product will be ———— 2 . 17
Which added to the leſſer diameter gives the mean diameter ———— 22 . 07
Which mean diameter being ſquared (that is, multiplied by it ſelf) produceth ———— 487 . 0849
Which product multiplied by ———— 0 . 0027
The product thence ariſing is ———— 1 . 315+
Which multiplied by the length of the Veſſel ———— 27 . 4
The product is the number of Ale Gallons ſought, to wit ———— 36.031+

The Reaſon of the Rule.

Two Things are taken for granted in the ſaid Rule, *viz.* Firſt it is ſuppoſed that if $\frac{7}{10}$ of the difference of the two diameters at the Bung and Head, be added to the leſſer diameter, the ſum ſhall be an equated or mean diameter (near enough for practical uſe though it be not exact) *viz.* If there be a Cylinder whoſe diameter is equal to that mean diameter, and whoſe length is equal to the length of the Veſſel, that Cylinder ſhall be equal to the capacity of the Veſſel very near. Secondly the ſaid Rule preſuppoſeth that 231 cubick inches are equal to a Wine Gallon, and 282 equal to an Ale Gallon; concerning which Equalities (eſpecially the latter) Artiſts differ ſomewhat in their experiments; but according to any Equality which in that particular ſhall be agreed on, from this that follows a Rule may be framed, and Tables thence calculated for gaging a full Veſſel without conſiderable Error.

Taking then thoſe two things above mentioned for granted, we may rightly infer that if a Cylinder hath for its Baſe a Circle whoſe ſuperficial content is 231 inches, every inch in length of that Cylinder will contain 231 cubick inches, or one intire Wine Gallon; now foraſmuch as all Circles are in ſuch proportion one to the other as the ſquares of their diameters, it ſhall be as 294.11844, (to wit, the ſquare of the diameter of that Circle whoſe ſuperficial content is 231) is to 1 (to wit, the ſuperficial content 231 conſidered as the Baſe of one Wine Gallon;) or as 1 is to .0034: So is the ſquare of the equated (or any other) diameter, to the ſuperficial content of that Circle in Wine Gallons and parts of a Gallon, which content multiplied by the length of the Veſſel will produce its ſolidity or capacity in Wine Gallons. Therefore the firſt part of the preceding Rule for finding of the number of Wine Gallons contained in a full Veſſel is manifeſt: And after the ſame manner, ſuppoſing as before 282 cubick inches are equal to an Ale Gallon, the decimal .0027 preſcribed in the ſaid Rule will be found out.

Upon thoſe Grounds Mr. *Wingate* compos'd his Gaging Rod; Mr. *Oughtred* alſo in his Circles of

Proportion hath delivered another Rule for Gaging, from whence his Gaging-rod is deduced; but the particular Conſtructions of thoſe Rods, and likewiſe the making of Tables for the ſame purpoſe, being handled by ſeveral Artiſts, I ſhall not inſiſt upon them.

Now if the induſtrious and more curious Arithmetician, after he is well exercis'd in vulgar Arithmetick, deſires further Knowledge in finding out the Anſwer of ſubtil Queſtions about numbers, his beſt Guide will be the admirable *Algebraical* Art, which diſcovers Rules for the ſolving of *Problems*, as well Arithmetical as Geometrical, that are above the reach of any of the Rules of common Arithmetick, or practical Geometry, as may partly appear by the two Rules in the aforegoing 52 and 65 Queſtions, as alſo by the two following Queſtions, with which I ſhall conclude this Chapter.

Queſt. 74. To find two numbers in a given proportion, ſuppoſe the leſſer to the greater as 2 to 3 and ſuch, that if the leſſer number be added to the ſquare of the greater, alſo if the greater number be added to the ſquare of the leſſer, the two ſums ſhall be ſquare numbers whoſe roots are expreſſible by rational or true numbers (fractions being admitted for numbers.)

Anſwer, $\frac{1}{10}$ and $\frac{3}{20}$.

$a : e :: 2 : 3$

$2e = 3a$

$e = \frac{3a}{2}$

$aa + \frac{3a}{2} = bb$

$\frac{9aa}{4} + a = cc$

$\frac{5aa}{4} - \frac{a}{2} = cc - bb$

bcl $b - \frac{a}{2} = c$ $bb - ba + \frac{aa}{4} - bb = \frac{5aa}{4} - \frac{a}{2}$

$- ba = aa - \frac{a}{2}$

$ba = \frac{a}{2} - aa$

$b = \frac{1}{2} - a$

$-a + aa = aa + \frac{3a}{2}$

$-a = \frac{3a}{2}$

$= \frac{5a}{2}$

$1 = 10a$

$a = \frac{1}{10}$

$e = \frac{3}{}$

The ſquare of $\frac{3}{20}$ (the greater number) is	$\frac{9}{400}$
To which adding the leſſer number	$\frac{1}{10}$
The ſum in its leaſt terms will be	$\frac{49}{400}$
Which is a ſquare number, whoſe root is	$\frac{7}{20}$
Again, the ſquare of $\frac{1}{10}$ (the leſſer number) is	$\frac{1}{100}$
To which adding the greater number	$\frac{3}{20}$
The ſum in its leaſt terms will be	$\frac{4}{25}$
Which is a ſquare number whoſe root is	$\frac{2}{5}$

Alſo the ſaid numbers $\frac{1}{10}$ and $\frac{3}{20}$ are one to the other as 2 to 3, wherefore the queſtion is ſolved, which numbers $\frac{1}{10}$ and $\frac{3}{20}$ are found out by this following.

Theorem.

If the fraction $\frac{1}{4}$ be divided into any two parts; either of thoſe parts being increaſed with the ſquare of the other part ſhall give a fraction having a rational ſquare root.

Wherefore by dividing $\frac{1}{4}$ into the two fractions $\frac{1}{10}$ and $\frac{3}{20}$, which are in the preſcribed proportions of 2 to 3, thoſe fractions will ſatisfie the conditions in the queſtion propounded.

Likewiſe theſe two fractions $\frac{722}{10080}$ and $\frac{1083}{10080}$ will anſwer the queſtion, and are found out without extracting any root; but the manner of finding out the ſaid Theorem and laſt mentioned fractions, I have ſhewn in the 24th. queſtion of my Third Book of the Elements of *Algebra.*

Queſt. 75. To find 3 numbers, ſuch that the ſquare of any one of them being added to the other two numbers, the ſum of ſuch addition ſhall be a ſquare number, whoſe root is a rational number.

Anſw. 1, $\frac{8}{3}$, and $\frac{16}{3}$.

The Proof.

Firſt, the ſquare of the firſt number 1 is	1
To which adding the ſecond and third numbers $\frac{8}{3}$ and $\frac{16}{3}$, the ſum will be	9
Which is a ſquare number whoſe root is	3
Secondly, the ſquare of the ſecond number $\frac{8}{3}$ is	$\frac{64}{9}$
To which adding the firſt and third numbers 1 and $\frac{16}{3}$, the ſum in its leaſt terms will be	$\frac{121}{9}$
Which is a ſquare number whoſe root is	$\frac{11}{3}$
Thridly, the ſquare of the third number $\frac{16}{3}$ is	$\frac{256}{9}$
To which adding the firſt and ſecond numbers 1 and $\frac{8}{3}$, the ſum in its leaſt terms will be	$\frac{289}{9}$
Which is a ſquare number whoſe root is	$\frac{17}{3}$

Wherefore it is manifeſt that the three numbers 1, $\frac{8}{3}$ and $\frac{16}{3}$ will ſatisfie the conditions in the queſtion, which may be ſolved alſo by other numbers, but the manner of finding them out I have ſhewn in the 32d. Queſtion of my 3d. Book of the Elements of *Algebra.*

CHAP XI.

Of Sports and Paſtimes.

Probl. I.

To diſcover a number which any one ſhall have in his mind, without requiring him to reveal any part of that or any number whatſoever.

AFter any one hath thought upon a number at pleaſure, bid him double it, and to that double bid him add any ſuch even number which you pleaſe to aſſign, then from the ſum of that addition let him reject one half, and reſerve the other half: Laſtly, from this half bid him to ſubtract the number which he firſt thought upon; then may you boldly tell him what number remaineth in his mind after that ſubtraction is made, for it will always be half the number which you aſſigned him to add.

For Example ſuppoſe he thought upon 6, the double thereof is 12, to which bid him add ſome even number at your pleaſure, ſuppoſe 4, ſo will the ſum be 16, whereof the half is 8, from which if he ſubtract 6 (the number firſt thought on) the remainder is 2 (to wit, half the number 4, which was by you aſſigned to be added;) which remainder you diſcover, notwithſtanding all the operation was performed in his mind, without his making known of any number whatſoever. Note, that the adding of an even number as aforeſaid is not of neceſſity, but only to avoid a fraction which will ariſe by taking the half of an odd number.

The

The Reaſon of the Rule.

If to the double of any number (which number for diſtinction ſake I call the firſt) a ſecond number be added, the half of the ſum muſt neceſſarily conſiſt of the ſaid firſt number, and half the ſecond; therefore if from the ſaid half ſum the firſt number be ſubtracted, the remainder muſt of neceſſity be half of the ſecond number which was added.

Probl. II.

Two numbers, the one even and the other odd, being propounded unto two Perſons, to the end they may (out of your ſight) ſeverally chuſe one of thoſe numbers; to diſcover which of theſe numbers each perſon ſhall have choſen.

Suppoſe you have propounded unto *Peter* and *John* two numbers, the one even and the other odd as 10 and 9, and that each of thoſe Perſons is to chuſe one of the ſaid numbers unknown to you. Now to diſcover which number each perſon ſhall have choſen, you muſt take two numbers, the one even and the other odd, as 2 and 3; then bid *Peter* multiply that number which he ſhall have choſen by 2; and cauſe *John* to multiply that number which he ſhall have choſen by 3; that done, bid them add the two products together, and let them make known the ſum to you, or elſe demand of them whether the ſaid ſum be even or odd, or by any other way more ſecret endeavour to diſcover it, by bidding them to take the half of the ſaid ſum, for by knowing whether the ſaid ſum be even or odd, you do obtain the principal end to be aimed at, becauſe if the ſaid ſum be an even number, then infallibly he that multiplied his number by your odd number (to wit, by 3) did chuſe the even number (to wit, 10;) but if the ſaid ſum happen to be an odd number, then he whom you cauſed to multiply his number by your odd number (to wit, by 3) did infallibly chuſe the odd number (to wit, 9.)

For Example, if *Peter* had made choice of 10, and *John* 9, ſuppoſe you willed *Peter* to multiply his number 10 by 2, and *John*, to multiply his number 9 by 3; the Products will be 20 and 27, whereof the ſum is 47, which being an odd number, you may thence conclude that *John* whom you cauſed to multiply his number by 3, did chuſe the odd number 9, and therefore *Peter* did chuſe 10. But if you had willed *John* to have multiplied his number 9 by 2, and *Peter* to have multiplied his number 10 by 3, the products would have been 18 and 30, whereof the ſum is 48, which is an even number, from whence you may infer, that he that multiplied his number by 3 did chuſe the even number, and therefore *Peter* choſe 10, and *John* 9.

Demonſtration.

The reaſon of the ſaid Rule is very eaſie, and dependeth principally upon the 28th. and 29th. Propoſitions of the 9th. Book of *Euclid*; for one may infer from the 21ſt. of the ſame Book, that an even number multiplied by any number whatſoever produceth an even number, but an odd number is of a different nature, for if it be multiplied by an even number, the Product is an even number (by the ſaid 28 Propoſition;) and if it be multiplied by an odd number, the product is odd (by the ſaid 29th. Propoſition.) Therefore if in making this Sport it happeneth that the even number be multiplied by your odd number, both the products ſhall be even, and conſequently the ſum ſhall be infallibly an even number) by the ſaid 21 propoſition.) But if it happen that you cauſe the odd number to be multiplied by your odd number, that product will be odd, and the other product even, therefore the ſum of theſe two products ſhall be an odd number (as *Clavius* hath demonſtrated upon the 23d. of the 9th. of *Euclid.*

Probl. 3.

A certain number of diſtinct Things being propounded, to diſpoſe them in ſuch an order, that caſting away always the ninth, or the tenth, or any other that ſhall be aſſigned, unto a certain number, thoſe remaining may be ſuch as were firſt intended to be left.

This Problem is uſually propounded in this manner, *viz.* Fifteen *Chriſtians* and Fifteen *Turks* being at Sea in one and the ſame Ship in a terrible Storm, and the Pilot declaring a neceſſity of caſting the one half of thoſe Perſons into the Sea, that the reſt might be ſaved; they all agreed that the Perſons to be caſt away ſhould be ſet out by Lot after this manner, *viz.* the thirty Perſons ſhould be placed in a round form like a Ring, and then beginning to count at one of the Paſſengers, and proceeding circularly, every Ninth Perſon ſhould be caſt into the Sea, until of the thirty Perſons there remained only fifteen. The queſtion is, how thoſe thirty Perſons ought to be placed, that the Lot might infallibly fall upon the fifteen *Turks*, and not upon any of the fifteen *Chriſtians*? For the more eaſie remembring of the Rule to reſolve this queſtion, I ſhall preſuppoſe the five Vowels, *a*, *e*, *i*, *o*, *u*, to ſignifie five numbers, to wit, (*a*) one (*e*) two, (*i*) three, (*o*) four, and (*u*) five; then will the Rule it ſelf be briefly comprehended in in theſe two following verſes.

From numbers, aid, and art
Never will Fame depart.

In which Verſes you are principally to obſerve the vowels, with their correſpondent numbers before aſſigned, and then beginning with the *Chriſtians*, the vowel *o* (in *from*) ſignifieth that four *Chriſtians* ate to be placed together; next unto them, the vowel *u* (in *num*) ſignifieth that five *Turks* are to be placed; in like manner *e* (in *bers*) denoteth 2 *Chriſtians*; *a* (in *aid*) 1 *Turk*; *i* (in *aid*) 3 *Chriſtians*; *a* (in *and*) 1 *Turk*; *a* (in *art*) 1 *Chriſtian*; *e* (in *ne.*) 2 *Turks*; *e* (in *ver*) 2 *Chriſtians*; *i* (in *will*) 3 *Turks*; *a* (in *fame*) 1 *Chriſtian*; *e* (in *fame*) 2 *Turks*; *e* (in *de*) 2 *Chriſtians*; *a* (in *part*) 1 *Turk*.

The Invention of the ſaid Rule, and ſuch like, dependeth upon the ſubſequent demonſtration, *viz.* if the number of Perſons be thirty, let thirty Figures or Cyphers be placed circularly, or elſe in a right Line as you ſee,

oooooooooooooooooooooooooooooo

That done, begin to count from the firſt, and mark the ninth (or what other ſhall be aſſigned) by putting a point or croſs over it; then count forward from that which you have marked, and place nother point over the next ninth; and continue o do the ſame, beginning again when you ſhall be t the end (if the Cyphers are placed in a right Line) and paſſing over thoſe, which you ſhall have lready marked, until you have marked the number required, as in the Example propounded, unil you have marked 15, for then all the Cyphers marked ſhall be thoſe which muſt be caſt away, nd the others thoſe which ſhall remain. Hence t is evident, that if you obſerve how thoſe Cyphers which are marked, are diſpoſed amongſt hoſe which are not marked, you will eaſily make Rule for any number whatſoever.

By this Invention (as ſome do conjecture) the amous Hiſtorian *Joſephus* the *Jew*, preſerved his Life very ſubtilty in the Cave, to which himſelf and Forty of his Country Men had fled from the furious and conquering *Romans* at the Siege of *Jotapata*: For his ſaid Country Men having moſt wickedly reſolved to kill one another, rather than ield to their Enemies, he at length (when no Arguments that he could uſe would diſſwade them from ſo horrid an Act) prevailed with them to excute their tragical Deſign by Lot; and ſo by the elp of the aforeſaid Artifice, as we may ſuppoſe, imſelf with one other Perſon only remaining live, after the reſt were inhumanly murthered, ney agreed to put an end to the Lot, and therey ſave their Lives This Story you may ſee at rge in the Fourteenth Chapter of the Third Book of the Hiſtory of *Joſephus* of the Wars of ne *Jews*.

Probl

Probl. 4.

Many numbers which proceed from 1 *or unity in a progreſſion, according to the natural order of numbers, (ſuch as theſe,* 1, 2, 3, 4, 5, 6, &c.) *being placed in a round form like a Ring ; to diſcover which of thoſe numbers any one ſhall have thought upon.*

Let any multitude of numbers in the aforeſaid progreſſion, ſuppoſe theſe 10, to wit, 1. 2. 3. 4. 5. 6. 7. 8. 9. 10 to be written upon 10 Ivory counters (or for want thereof upon ten ſmall pieces of Paper) which may be repreſented by theſe 10 Letters, A. B. C. D. E. F. G. H. K. L. *viz.* ſuppoſe 1 to be written upon the counter A, 2 upon B, 3 upon C. *&c.* Then having placed thoſe Counters circularly as you ſee (with their blank faces uppermoſt, and the Figures underneath, that the ſubtilty of the Sport may the better be concealed) let any one think upon any number of unites which doth not exceed 10 ; that done bid him touch one of thoſe Counters at pleaſure, and to the number on the backſide of the counter touched (which you cannot be ignorant of, having noted well the place of 1 or



A) add ſecretly in your mind, the juſt number of all the counters, and reſerve the ſum; then bid him imagine in his mind the counter touched to be the number which he thought, and from that counter to count backwards, untill he ſhall have made up the aforeſaid ſum, which you reſerved, ſo will his computation infallibly end upon the counter upon which the number thought upon is written.

For Example, ſuppoſe that he thought 7 or G, and that he touched B, to wit, 2. add to 2 the number of all the counters, to wit, 10, ſo the ſum will be 12; then bid him to count unto 12 beginning at B going backwards, and eſteeming B to be the number thought, to wit 7, ſo will 8 fall upon A, 9 upon L, 10 upon K, 11 upon H, and laſtly, 12 upon the counter G, which being turned up will ſhew 7 the number thought.

The reaſon of this Rule is not difficult to be apprehended, two principles being preſuppoſed, the one is this, to wit, many counters or things whatſoever being diſpoſed orderly one after the other, in one continu'd line, whether it be right or circular, if you value or name the firſt counter to be ſome number of unites at pleaſure, and continue to count forward according to the natural order of numbers until another number be named which falleth upon the laſt counter; or if you imagine or name the laſt counter, to be the ſame number of unites as before you put upon the firſt, and continue to count backwards unto the firſt counter; I ſay, that the ſame number will be named at the end of both thoſe computations: For Example, in theſe 9 Letters A.B.C.D.E.F.G.H.K. if the Letter A be eſteemed to be 4, and from thence you count forwards unto K, according to the natural order of numbers the Letter K will fall upon the number 12. In like manner, if you eſteem K to be 4, and count backwards from K to A, the Letter A will likewiſe fall upon 12.

4.	5.	6.	7.	8.	9.	10.	11.	12
A.	B.	C.	D.	E.	F.	G.	H.	K.
12.	11.	10.	9.	8.	7.	6.	5.	4.

The other principal is this, to wit, many counters being diſpoſed in a round manner like a Ring, if you eſteem any one of thoſe counters to be ſome number at pleaſure, and then from that counter if you count circularly, until you end upon the counter where you began, the number laſt named will be equal to the ſum of the number of all the counters, and of the number which you put upon the firſt counter; for Example, If D be one of 10 Letters placed in a circumference, and that imagining D to be 7, you begin with it, and count round the whole circumference, according to the natural progreſſion of numbers, till



you end with D where you began; the number 17 which is compoſed of 10 and 7 will neceſſarily fall upon D; for 9 (which is the number of Letters in the circumference beſides D) being added to 7 (which was firſt put upon D) makes 16, to which 1 being added (becauſe D doth end as well as begin the circumference) the ſum is 17.

Now theſe two Principles being preſuppoſed, it will not be difficult to apprehend the reaſon of the aforeſaid Rule in all caſes that can happen; for imagine that one hath thought upon 7, or the counter G, then that counter which he ſhall touch muſt either be the ſame counter G or ſome other that proceedeth or followeth G.

Firſt, therefore ſuppoſing the counter or number touched to be the ſame with the number thought, the truth of the Rule will be then evident, for by the Rule given, he ſhall begin to count from the ſame G unto 17, putting 7 upon G, therefore by the ſecond preſuppoſition the number 17 will fall upon G.

Secondly, imagine that he touched a counter or number following G the number thought, as L or 10, then according to the Rule adding 10 (the multitude of all the counters placed circularly) unto 10, or L (the counter touched) bid him count backwards unto 20 by beginning at L, and eſteem L to be 7. Now becauſe by beginning to count at G which is 7, and proceeding to count forward, the number 10 will fall upon L; therefore by the firſt preſuppoſed principle, if we eſteem L to be 7 and count backwards, the number 10 will infallibly fall upon G, and then the number 20 ſhall alſo fall upon the ſame G by the ſecond preſuppoſed principle.

Laſtly, imagine he touched ſome number or counter which precedeth 7 the number ſought, as B or 2; then adding 10 to 2, you are to bid him count unto 12, he having firſt imagined B to be the number thought 7, and going backwards to A, L, K, *&c.* Now becauſe by proceeding to count at B, which is 2, and beginning to count forward to C, D, *&c.* the number 7 falleth upon G; therefore if one imagine that G is 2, and from thence count backwards towards F, E, *&c.* the number 7 will fall upon B (by the firſt preſuppoſed principle;) therefore when one aſſumeth B to be 7, and counteth towards A, L, *&c.* to any aſſigned number, it is in effect as much as when one imagineth G to be 2, and counteth towards F, E. *&c.* unto the ſaid aſſigned number, for each of thoſe computations will end in the ſame point; but it is manifeſt (by the ſecond preſuppoſed principle) that eſteeming G to be 2, and counting towards F, E, D, *&c.* round the whole circumference, the number 12 will fall upon the ſame G. And becauſe G being ſuppoſed to be 2, and counting on the ſame coaſt as before, the number 7 falls upon B; therefore if the computation be continued on the ſame coaſt from B 7, unto 12, the number 12 will fall upon the ſame G. So that the practice of this ſport in all its Caſes is fully demonſtrated.

Note, that to the number of the counter touched you may not only add the number of all the counters once (as the rule directs) but twice, thrice or more times: For Example, B being touched, you may cauſe him to count unto 12, or unto 22, or to 32, 42, *&c* the reaſon whereof is evident from the ſecond preſuppoſed principle.

Probl.

Probl. 5.

Many numbers being ſhewed by pairs, to wit, two by two, unto any one, that he may think upon any one of thoſe pairs at pleaſure; to diſcover the pair that was thought upon.

Let 20 numbers, ſuppoſe theſe, 1. 2. 3. 4. 5. 6. 7. 8. 9. 10. 11. 12. 13. 14. 15. 16. 17. 18. 19. 20. be written upon Ivory counters (or for want thereof upon ſmall pieces of Paper) to wit, 1 upon one counter, 2 upon another, 3 upon a third, *&c.* Then diſpoſe then into pairs as you ſee, *viz.* Suppoſe 1 and 2 to be one pair, 3 and 4 to be another pair, *&c.* and of theſe pairs let any one think upon which pair he pleaſeth. That done, you are to diſtribute the ſaid 20 numbers in ranks, into the form of a long ſquare, until there be 5 numbers in length, and 4 in breadth, after this manner, *viz.*

1.	2
3.	4
5	6
7.	8
9.	10
11.	12
13.	14
15.	16
17.	18
19.	20

Lay the three firſt number, 1, 2 and 3 in a rank (as you ſee in the ſecond figure) from A towards B; then place 4 underneath 1, and 5 after 3 (in the ſaid rank AB.) Again place 6 under 4, and 7 after 5 (in the ſaid rank AB.) Then place 8 under 6, alſo 9. 10. 11. on the right hand of 4 in the rank CD. Again place 12 under 9, and 13 on the right hand of 11 in the rank CD. and 14 under 12. Moreover place 15. 16. 17 on the right hand of 12 in the rank EF. Laſtly, place 18. 19. 20. on the right hand of 14 in the rank GH, ſo will all the numbers be ranked as you ſee in the Table. That done, you are to demand of him that thought upon two numbers as aforeſaid, in what rank or ranks the ſaid numbers do happen to be found, *viz.*

A	1	2	3	5	7	B
C	4	9	10	11	13	D
E	6	12	15	16	17	F
G	8	14	18	19	20	H

in which of the ranks AB, CD, EF, GH, or in which two of the ſaid ranks: Now if he anſwer that the two numbers which he firſt thought upon are in the firſt rank AB. then 1 and 2 ſhall be the numbers thought upon; if in the ſecond CD, then 9 and 10 ſhall be the numbers thought; if in the third rank EF, then 15 and 16 ſhall be the numbers thought: If they are in the fourth rank GH, then 19 and 20 ſhall be the numbers thought; but if he ſhall ſay that the numbers thought are in different ranks, then you are heedfully to mark the ſaid numbers 1 and 2, 9 and 10, 15 and 16, 19 and 20, which may be called the keys of the ſport, in regard they ſerve not only to diſcover the two numbers thought, when they are both in one and the ſame rank (as aforeſaid) but alſo when they are in two different ranks, for in this latter caſe as ſoon as it hath been declared to you in which two ranks the two numbers thought are placed, you muſt take the key of the higheſt of thoſe two ranks, and deſcending in a down right line from the firſt number of that key unto the lower of the ſaid two ranks, you ſhall there find one of the two numbers thought, and upon the right hand of the ſecond number of the ſaid key, at the ſame diſtance ſidewiſe from the ſecond number of the key, as one of the numbers thought was diſtant from the firſt number of the key, you ſhall find the other number thought.

For Example, ſuppoſe the two numbers thought are 7 and 8, and that it ſhall be declared unto you that they are in the firſt and fourth ranks; take then the key of the higheſt of theſe two ranks, to wit, of the firſt, which is 1 and 2, and deſcending down right from 1 unto the fourth rank, you ſhall there find 8 one of the numbers thought; then ſeek ſidewiſe on the right hand of 2. the ſecond number of the key, a number as far ſeparated from 2, as 8 is diſtant from 1, and you will find 7 the other number thought.

Again, ſuppoſe he ſaid that the numbers thought are in the ſecond and third ranks; take then the key of the ſecond rank which is 9 and 10. and deſcending downright from 9 to the third rank, you ſhall there find 12 which is one of the numbers thought; then ſeek ſidewiſe on the right hand of 10 (the ſecond number of the key) a number as far diſtant from 10 as 12 is from 9, and you ſhall find 11 which is the other number thought.

The reaſon of this will be apparent from a ſerious conſideration of the placing of the numbers according to the Rules before given, for it is thereby evident that of the numbers coupled two by two, there can never be found more than one pair in one and the ſame rank, and of all the other pairs one number is always found in one rank, and the other number in another rank.

Note alſo, that this ſport may be practiſed with divers perſons at once, and not only with 20 numbers, but with any ſuch multitude of numbers which is produced by the multiplication of any two numbers which differ by 1, or unity; as 30, which is the product of 5 multiplied by 6, and 42 which is the product of the multiptication of 6 and 7. That which is chiefly to be regarded is the placing of the numbers in ranks according to the directions before given: And for the more eaſie comprehending of that Order, I have in the following Table ranked 30 numbers in their due places, which being compared with the former Table, and well viewed, will be a clearer Illuſtration than can be expreſt by many words.

1	2	3	5	7	9
4	11	12	13	15	17
6	14	19	20	21	23
8	16	22	25	26	27
10	18	24	28	29	30

Probl. 6.

hree jealous Husbands with their Wives, being ready to paſs by Night over a River, do find at the River ſide a Boat which can carry but two Perſons at once, and for want of a Boatman they are neceſſitated to row themſelves over the River at ſeveral times: the queſtion is how theſe 6 Perſons ſhall paſs 2 by 2, ſo that none of the 3 Wives may be found in the Company of 1 or of 2 Men unleſs her Husband be preſent.

They muſt paſs in this manner, *viz.* Firſt two Women paſs, then one of them bringeth back the Boat and repaſſeth with the third Women; that done, one of the three Women bringeth back the Boat, and ſitting down upon the Ground with her Husband permitteth the other two Men to paſs over to find their Wives; then one of the ſaid Men with his Wife bringeth back the Boat, and placing her upon the Ground he taketh the other Man, and repaſſeth with him: Laſtly, the Woman which is found with the three Men entreth into the Boat, and at twice goeth to fetch over the other two Women.

Probl. 7.

Two merry Companions are to have equal ſhares of 8 Gallons of Wine, which are in a Veſſel containing exactly 8 Gallons, now to make this equal Partition they have only two other empty Veſſels, whereof one containeth 5 Gallons, and the other 3; the queſtion is, how they ſhall exactly divide the Wine by the help of thoſe three Veſſels.

Firſt, from the Veſſel which containeth 8 gallons and is full of Wine, let 5 Gallons be poured into the empty Veſſel of 5, and from this Veſſel ſo filled let 3 be poured into the empty Veſſel of three, ſo there will remain 2 Gallons within the Veſſel of 5. Then let the three Gallons which are within the Veſſel of 3 be poured into the Veſſel of 8, which will now have 6 Gallons within it, that done let the 2 Gallons which are in the Veſſel of 5, be put into the empty Veſſel of 3, then of the 6 Gallons of Wine which are within the Veſſel of 8 fill again the five, and from thoſe 5 pour out 1 Gallon into the Veſſel of 3, which wanted only 1 Gallon to fill it, ſo there will remain exactly 4 Gallons within the Veſſel of 5 and 4 within the other two Veſſels. This queſtion may be reſolved in another way, but I leave that as an Exerciſe to the Wit of the ingenious Reader.

Now albeit at firſt ſight it may be thought by ſome, that the two laſt mentioned *Problems* cannot be reſolved by any certain Rule, but only by many Trials, yet by infallible Argumentation and Diſcourſe, the ſolution of thoſe Queſtions may be found out, or elſe the impoſſibility of them, if by chance they ſhould have been propounded, impoſſible; as the moſt ingenious *Gaſper Bachet* hath manifeſted in a little Book in the *French* Tongue, Entituled, *Problemes plaiſans & delectables qui ſe font par les nombres*, from which Book I have extracted the Contents of this Chapter.

A Supplement

A
SUPPLEMENT.

CONTAINING

Several Eaſy Contractions, in the Neceſſary and Uſeful Parts of **Arithmetick**, conducing to a Ready and Expeditious way of Accounting.

WITH

Decimal Tables of *Flemiſh* **Exchanges**, at the moſt Uſual and Probable Rates of Exchange.

AS ALSO

Tables of **Intereſt**, ſerving at any Rate, for any Time.

To which is added,

The **Menſuration** of Superficies and Solids, in Various and Practical Inſtances.

BY

GEORGE SHELLEY, Writing-Maſter, at the *Hand* and *Pen* in *Warwick-Lane*, near St. *Paul's*

LONDON,
Printed for *J. Philips*, *J. Taylor*, and *J. Knapton*,
in St. *Paul's Church-Yard*, M DCC IV.

TO THE
READER.

T*IS very well known, That the*
Ingenious Mr. Wingate *hath*
excellently Explained the
rinciples of that moſt Uſeful Science,
rithmetick; *which together with the*
comparable Additions of Mr. Kerſy,
e Theoretical Part is ſo largely in-
ſted on, that it is in a manner inca-
ble of any further Improvement.

But thô a Theory may be ſo ſettled as
have no Alteration, yet the Practices
educed, are often changed and much
prov'd.

To the R E A D E R.

In this Edition of Mr. Wingate' Arithmetick, *I was prevail'd upon, no only to Correct it, but alſo to Add as Supplement, ſome Practical, Compendious, and* Eaſy *Rules, not inſerted b the Authors; or whatever elſe migh tend to the making it more generall Uſeful. And which I have endeavoure chiefly in the moſt neceſſary Rules* Arithmetick.

The Tables of Simple Intereſt *and* Exchanges, *were never before Publiſhed, and are ſuch as I dare recommen for Practice, beyond any I know extant.*

G. SHELLEY

SECTION I.

MULTIPLICATION.

HOW to *Multiply* by any Number under 20, ſo as to bring the whole Operation into one Line.

Rule (1.) *Multiply each Figure in the Multiplicand by the nite Figure in the Multiplier, adding unto each its back Figure; s in the following Examples,*

47947
13
623311

27942
17
475014

vhich are done thus, *viz.* 3 times 7 is 21, ſet down 1 and carry 2; hen 3 times 4 is 12, and 2 is 14, and 7 (which is the back Figure to 4) is 21, 1 and carry 2; 3 times 9 is 27, and 2 is 29, nd 4 (the back Figure) is 33, 3 and carry 3; 3 times 7 is 1, and 3 is 24, and 9 (the back Figure) is 33, 3 and carry 3; 3 times 4 is 12, and 3 is 15, and 7 is 22, 2 and carry 2, which added to 4 makes 6.

In like manner, the whole Operation of the Multiplication of Numbers between 20 and 30, are brought into one Line, by taking in the Double of the back Figure, and between 30 nd 40, the Treble of it, *&c.*

By which Rule, in *Multiplying* by Numbers compounded of thoſe under 20, the *Product* is readily found thus;

By 112, either *Multiply* by 2 and 11, or by 12 and 15 s,

94267
112
188534
1036937
10557904

94267
112
1131204
94267
10557904

Alſo in *Multiplying* by 1614, and 1715; in the firſt I *Multiply* by 14 in one Line, and by 16 in the other: And in the latter by 15 in one Line, and 17 alſo in the other, as above directed; obſerving always, To place the firſt Figure of a Line under its own *Multiplier*; as,

```
   479479        479479
     1614          1715
  6712706       7192185
7671664       8151143
773879106     822306485
```

otherwiſe for any Number under 20, a way not ſo troubleſome to the Memory.

Rule (2.) *Multiply by the unite Figure, ſetting that whole Product one Figure back on the right hand, which added to the Multiplicand, gives the Product.* viz.

```
 47947      27942
    13         17
143841     195594
623311     475014
```

But for 112, *Multiply* by 12, ſetting the *Product* two Figures back, which likewiſe added to the *Multiplicand*, gives the *Product*; as,

```
   94267
     112
 1131204
10557904
```

Note, If the *Multiplicand* hath leſs Significant Figures in it, then the *Multiplier*, work as if it were the *Multiplier*; as in Examples,

```
   13000         1107
     874         3758
11362000        26306
               413380
              4160106
```

II. How to *Multiply* by the component *Parts* of a Number instead of the *whole*.

Tha

That is to *Multiply* 42991 by 7, and that *Product* by 8, is the ſame as if it were *Multiplied* by 56, the *Product* of 7 by 8. Example——

42991
7
300937
8
2407496

Note, If any *odd Numbers* are given that cannot be produced by the *Multiplication* of any two Figures, then find out two Figures whoſe *Product* comes neareſt, either under or above it, to which add what is wanting, or from which Subtract what is over the given Number; as,

479 by 47	927 by 29
6	4
2874	3708
8	7
22992 Subtracted	25956 Added
479	927
22513	26883

III. To *Multiply* Numbers of ſeveral Denominations.

(1.) *By whole Numbers.*

l. *s.* *d.*	*C.* *q.* *l.*
47 : 18 : 11	9 : 3 : 15
8	6
383 : 11 : 04	59 : 1 : 06

(2.) *By the parts inſtead of the whole*, viz.

Ya. *q.* *n.*	*C.* *q.* *l.*
47 : 3 : 3 by 45	21 : 3 : 08 by 36
5	6
239 : 2 : 3	130 : 3 : 20
9	6
2157 : 0 : 3	785 : 2 : 08

(3.) *By mixt Numbers.*

```
 C.   q.   l.                         l.   s.   d.
 9    3    15 by 11 1/2               39 : 18 : 11   9 : 1/4
           11                                        9
108 : 2 : 25                          359 : 10 : 03
  4 : 3 : 21 : 1/2                      9 : 19 : 08 : 3/4
113 : 2 : 18 : 1/2                    369 : 09 : 11 : 3/4
```

```
            s.   d.
     47 . 18 . 11 . by 27 3/4
                 4
    191 . 15 .  8
                7
   1342 . 09 .  8
     11 . 19 .  8 . 3/4
   1330 . 09 . 11 . 1/4
```

In the firſt of theſe laſt examples, I *Multiply* by 11 adding one half of the *Multiplicand* for the ½. In the 2d. I *Multiply* by 9, adding a fourth for ¼. In the 3d. I *Multiply* by 4 and that *Product* by 7, gives 28 times the ſum, from which *Subtracting* ¼ of the firſt ſum, the *Remainder* is the *Product* of the given ſum, *Multiply'd* by 27 ¾.

(4) *By Numbers of Several Denominations.*

(1) By Aliquot Parts. (2) *By Croſs Multiplication*

```
        F.    i.    p.                         F.    i.
        47 . 08 . 06                           47 . 08
by       9 . 04 . 04                            9 . 04
       429 . 04 . 06                          423 . 00
 1/3    15 . 10 . 10                           06 . 00
 1 1/2   1 . 03 . 10                           15 . 06
       446 . 07 . 02     Anſwer                     02
                                              444 . 10 . 8
```

Which are done thus *viz.*

In the firſt, the ſeveral *Denominations* in the *Multiplicand*, are *Multiply'd* by the Integers in the *Multiplier*, unto which *Product*

duct add the one third of the *Multiplicand*, for 4 Inches (being one $\frac{1}{3}$ of one foot) and the $\frac{1}{12}$ of that $\frac{1}{3}$ for the 4 parts (being the $\frac{1}{12}$ of 4 Inches) gives the Anſwer.

In the ſecond called *Croſs Multiplication*, the *Feet* are *Multipli'd* by the *Feet*, then (Croſs-wiſe) *viz.* the 9 *Feet* in the *Multiplier* by the 8 *Inches* in the *Multiplicand*, gives 72 *Inches* or 6 *Feet*, and the 47 *Feet* in the *Multiplicand* being *Multiply'd* by 4 *Inches* in the *Multiplier*, gives 15 *Foot*, 8 *Inches*. Laſtly the *Inches* in both the *Factors*, being *Multiply'd* one into the other, make 32 parts or 2 *Inches*, 8 parts the *Total* of which is the *Product* of 47 *Feet*, 8 *Inches* by 9 *Feet* 4 *Inches*.

Section II.

DIVISION.

I. TO *Divide* whole Numbers by the *Component* parts of a Number.

(1) Examples having no *Remainders*.

Divide 22992 by 48 that is by 6 and that Quotient by 8.

6) 22992
8) 3832
479 Quotient

Divide 25956 by 28, or by 4,& that Quotient by 7.

4) 25956
7) 6489
927 Quot.

(2) Examples *having Remainders.*

Divide 47953 by 49

7) 47953
7) 6850 . 3
978 . 4

The Quotient 978 . $\frac{31}{49}$

Divide 47991 into 36 parts.

4) 47991
9) 11997 . 3
1333 . 0

$$1333 \cdot \frac{3}{36} \Big| \frac{1}{12}$$

Anſwer 1333 . $\frac{3}{36}$ | $\frac{1}{12}$

If 21 pieces of Bagg *Holland* contain 471 *Ells* how much is contain'd in one piece

3 | 471
7 | 157
22 . 3

Anſwer Ell. 22 . $\frac{9}{21}$ | $\frac{3}{7}$

The true remainders of which *Diviſions* are found by the following Rule,

Multiply the firſt Diviſor by the laſt remainder, adding thereto the firſt remainder, (if any be) gives a Numerator, whoſe Denominator is the whole Number compoſing the parts which Fraction ſo ariſing is a true Remainder, and the very ſame that would be found in Dividing by the whole.

II. Of *Dividing* Numbers of *Several Denominations*, by the *Parts* of the *Whole.*

l. s. d.
(1) Ex. If 19 . 14 . 10 be *Divided* among 16 Men what is each Mans Share.

4 | 19 . 14 . 10
4 | 4 . 18 . 08 . $\frac{1}{2}$
Anſwer 1 . 04 . 08 . $\frac{1}{8}$

(2) Ex. What is the weight of one *Bale*, when 25 weigh 93 . 1 . 01

C. q. l.
5) 93 . 1 . 01
5) 18 . 2 . 17 oz.
3 . 2 . 25 . 12 . $\frac{4}{5}$

III. To *Divide* Numbers by *Broken* or *Mixt Numbers.*

Rule, *Reduce both the Dividend and the Diviſor, into the ſame Denominator.*

(1) Ex. What comes 1 Yard of Flowred Damask to, when 13$\frac{1}{3}$ Yards coſt 21 *l.* 15 *s.* 11 *d.* to do this, reduce the Yards by *Multiplying* by 3 into thirds, which *Product* will be

40,

40, likewiſe Reduce the Sum into thirds, is 65*l.* 7*s.* 9*d.* which being to be *Divided* by 40, work according to the foregoing Rule taking $\frac{1}{8}$ of $\frac{1}{5}$ for the anſwer.

21 . 15 . 11   13.
3   3
5) 65 . 07 . 09   40
8) 13 . 01 . 06 . $\frac{3}{5}$
Anſwer 1 . 12 . 08 . $\frac{1}{5}$

(2) Ex. What comes one piece of Cloth to when 17$\frac{1}{2}$ pieces coſt

2
35

l. *s.* *d.*
57 . 07 . 09
2
5 114 . 15 . 06
7 22 . 19 . 01 . $\frac{1}{5}$
Anſwer 3 . 05 . 07 . $\frac{1}{35}$

3. Ex. Divide 8792 . $\frac{1}{2}$ by 27 . $\frac{1}{2}$

8792 . $\frac{1}{2}$ 2 .
2   55
5 17585   The Quotient is 319 . $\frac{8}{11}$
11 3517
319 . $\frac{8}{11}$

IV. How to *Divide* Numbers of *Several Denominations* by thoſe of *Several Denominations.*

Exam. (1.) How many pieces of Gold, each valu'd at 2. 11. 8. are contained in 478. 19. 10. Anſwer 185 pieces, as appears by the following operation.

2 . 11 . 8)

```
2 . 11 . 8) 478 . 19 . 10 (185
            2 . 11 . 08
            1 . 08 . 4
                    10
           21 . 03 . 4
           20 . 13 . 4
            0 . 10 . 0
                    10
           13 . 00 . 0
           12 . 18 . 4
              .01 . 8
              .19 . 10 Added
               21 . 6 Remainder
```

To do which, Firſt ſeek how many times 2 . 11 . 8 is contained in *l.* 4, which is once, then ſetting one in the Quotient, Multiply the Diviſor by it, the Product of which ſetting under, and Subtracting from 4 *l.* reſt 1 *l.* 8 . 4, which Multiply by 10, taking in 7 *l.* the next Figure of the Dividend, then taking a new Quotient Figure ,you repeat the ſame method as above, until finiſht. Obſerving at the laſt, to add the odd Money in the Dividend, to the Money Remaining, as in the *Example.*

Note, That in Adding, Multiplying, and Dividing of Money, work the ſecond place in the Shillings, as Angels.

Of

Section III.

Of REDUCTION.

I. *In* Money.

Pounds into Guineas, Multiply by 40, which Product divide by 43.
Marks into Pounds, Subſtract $\frac{1}{3}$
Pounds into Marks, Add $\frac{1}{2}$

II. *In* Avoirdupois.

To Reduce Hundreds, *&c.* into Pounds, Multiply by 112 in two lines, as Taught in *Multiplication*, taking in the Pounds contain'd in the odd Weight.

By this Weight Silk is Weigh'd, in which there is a great Pound of 24 *Oz.* and by which is Weigh'd Raw, Long, Short, China, Morea Silk, *&c.*

And a ſmall one of 16 *Oz.* by which Ferret, Fillozel, Sleeve Silk, Fine and Courſe, *&c.* are Weigh'd.

To Reduce one into the other.

Note, Great Pounds of 24 *Oz.* into Pounds of 16 *Oz.* Add $\frac{1}{2}$.

Small Pounds of 16 *Oz.* into Pounds of 24, Subtract $\frac{1}{3}$.

III. Liquid Meaſure.

Of Beer
Firkins into *Hogſheads* ——————— Take $\frac{1}{6}$
Hogſheads into *Barrels* ——————— Add $\frac{1}{2}$
Kilderkins into *Hogſheads* ——————— Take $\frac{1}{3}$
Barrels into *Tunns* ——————— Take $\frac{1}{7}$

54 *Gallons* one *Hogſhead* of Beer.

Of Wine	*Hogſheads* into *Tierce*	Add $\frac{1}{2}$
	Tierce into *Hogſheads*	Sub. $\frac{1}{3}$
	Tierce into *Pipes*	Take $\frac{1}{3}$
	Pipes into *Hogſheads*	Double it

63 *Gallons* one *Hogſhead* of Wine.

IV. Cloth Meaſure.

Of which there are 2 Sorts moſt in uſe among us, a *Yard* and an *Ell*, both Subdivided into 4 *Quarters*, and each *Quarter* into 4 *Nails*. But the *Nail* of an *Ell* is $\frac{1}{4}$ greater than the *Nail* of a *Yard*; whence it is, That 20 *Nails* or 5 *Quarters* of a *Yard* are contain'd in one *Ell Engliſh*.

Yards into *Ells Engliſh*	Subſtract $\frac{1}{5}$ part
Ells Engliſh into *Yards*	Add $\frac{1}{4}$ part
Yards into *Flemiſh*	Add $\frac{1}{3}$ part
Ells Flemiſh into *Yards*	Subſtract $\frac{1}{4}$ part
Ells Engliſh into *Flemiſh*	Add 0 and Divide by 6
Ells Flemiſh into *Engliſh*	Multiply by 6, and cut off the laſt Figure.

Of Tare *and* Tret, *&c.*

Call'd Beyond Sea, *The Courteſies of London*, becauſe not Practiſed in any other place; concerning which, obſerve, That

Groſs is the Weight of the Commodity, and that which contains it without any Deduction.

Tare is an Allowance for that which contains the Commodity, whether it be *Bag*, *Barrel*, *Sack*, *Frail*, &c. and is either,

I. At ſo much *per Bag*, *Barrel*, *Frail*, &c.
II. At ſo much *per Cent*: or,
III. By ſo much of the *Groſs Weight*, call'd, *Invoice-Tare*.

I. When the *Tare* is ſo much *per Bag*, as *Almonds*; per *Barrel*, as Oyl in Candy *Barrels*; per *Frael*, as Raiſins, *&c.*

Rule, *Multiply the Number of the ſaid Bags, Barrels, Frails, Hogſheads,* &c. *by the Allowance of Tare, which Product Subtract from the Groſs, and (when no Tret is mention'd) the Remainder is Neat.*

Ex. What is the *Neat Weight* of 38 Hogſheads of *Tobacco*, weighing *Groſs* 102 . 2 . 17, *Tare* per Hogſhead 70 *l.*

```
                                         38
 102  2 . 17                            2660
 112
 ----
  1297
 11497 Groſs
  2660 Tare
  ----
  8837 Neat
```

Or which is more ſhort, tho' not according to *Rule*, *Multiply* the Hogſheds by 5, *Dividing* the *Product* by 8, will give the *Tare* of the whole in *Hundreds*, *Quarters*, and *Pounds*.

```
  38        C.    q.   l.
   5       102 . 2 . 17 Groſs
8)190       23 . 3 . 00 Tare
  23 3/4    78 . 3 . 17 Neat    Equal to 8837 l.
```

II. When the *Tare* is at ſo much *per Cent*, and that,

(1.) If an *Aliquot part, or parts* of the *Hundred Weight*;

as *Figs*, *Almonds*, *Argol*, &c. 14 }
Caroteels, and *Buts of Currants*, &c. 16 } *per Cent.*
Oyl in uncertain *Caſks*, &c. 18 }

Rule, *Divide the whole Groſs by the ſaid part or parts that the Tare is of an Hundred, gives the Tare of the whole, for ſuch a part of the Groſs muſt be the Tare of the whole, as the given Tare is of an Hundred.*

Ex. What is the *Neat Weight* of 15 Barrels of *Argol*, *Groſs* C. 51 . 2 . 18. *Tare* 14 *per Cent*, which being the 8th of an Hundred, take the 8th of the *Groſs* for *Tare.* C.

```
        C.   q.   l.
   8)  51 . 2 . 18        Groſs
        6 . 1 . 23   4    Tare
       45 . 0 . 22 : 42   Neat
```

Example (2.) How many *Neat Hundreds*, in 18 Quarter Roles of *Currents* each *Groſs* 8 . 3 . 10 *Tare* 21 *l. per Cent.*

```
                 C.   q.   l.
                 8 .  3 . 10 each
                             3
                26 .  2 . 02
  l.                        16
        1/8)   159 .  0 . 12 oz. Groſs
  14    1/2)    19 .  3 . 15   8
   7             9 .  3 . 21 . 12
                29 .  3 . 09 . 04 Tare
               129 .  1 . 02 . 12 Neat
```

Tare					
For 2	Divide the *Groſs* by	7	and that quote by	8	gives the *Tare*.
4		4		7	
7		4		4	
8		2		7	

2: But if the *Tare* is not readily found to be an *Aliquot part*, or parts of a *Hundred.*

Rule, *Multiply the Pounds Groſs by the Allowance of Tare, Dividing the Product by* 112, *the Quotient is the Tare of the whole.*

Exam. What is the *Neat Produce* of 12 Barrels of *Pot-aſhes*, each *Groſs* 203 *l. Tare* 10 *l. per Cent.*

203

	203		2435	Groſs
	12		217 . ½	Tare
112)	24360	(217	2218 . ½	Neat
	196			
	840			
	56			

III. If the Allowance for *Tare* is not by the *Hundred*, but according to the *Groſs* Weight, as *Sumack* in *Baggs*, *Bales* or *Fangots* of *Aleppo*, *Cyprus*, and *Smyrna* Silk, &c. ſuch *Tare* is called, *Invoice-Tare*; of which the *Book of Rates*, in the Table of Allowances for *Tare*, ſays, *viz.*

Cyprus, *and* Smyrna *Silk.*

Bales	about, or above 300 *l.*	allowed for *Tare*	16
	from 300 to 200		14
	from 200 downwards		12

Tobacco *from* Virginia.

			L.
Hogſheads	5 *C.* and upwards	allow'd *Tare,*	100
	From 4 to 5,		90
	From 3 to 4,		80
	under 3 Hundred		70

Exam. What is the *Neat Weight* of 4 Hogſheads of *Virginia* Tobacco, weighing *Groſs* and Numbred? *viz.*

No.	*C.* *q.* *l.*		*l.*
27	2 . 2 . 19	*Tare* found as above	70
9	5 . 0 . 15		100
4	3 . 2 . 11		80
19	4 . 2 . 19		90
	16 . 0 . 08	*Groſs*	112) 340 (3
	3 . 0 . 04	*Tare*	4
	13 . 0 . 04	*Neat* and ſo of Silk, &c.	

Alſo Sugar *from* India.

In Casks and Caniſters, } Tare $\frac{1}{6}$
In Cheſts, and Casks, St. *Thome.* } Tare $\frac{1}{5}$

Tret, Is an Allowance of 4 *l. per* 104, on ſome ſort of Goods, (to the Buyer by the Merchant) for the *Duſt* and *Waſte* in them (which when allow'd) after you have Subtracted the Tare from the Groſs, the Remainder is *Suttle.*

Rule. *Which Suttle take out, and divide by* 26 (*becauſe the allowance of* 4 *l. per* 104, *is* $\frac{1}{26}$ *part*) *and the Quote is the* Tret, *which* Tret *Subtract from the Suttle, and the Neat Remains.*

Ex. How many *l. Neat* in 12 *C.* 3 *q.* 12 *l. Groſs, Tare* 2 *l. per Cent. Tret* 4 *l. per* 104?

```
26) 1415 (54                C.   q.   l.
    115                     12 . 3 . 12
     11                     112
                         8)1440  Groſs
                         7) 180
                             25 + Tare
                           1415  Suttle
                             54  Tret
                           1361  Neat
```

Otherwiſe to find and deduct Tret,

Rule. *Add* 2 *Cyphers to the Suttle Pounds, which Number divide by* 104, *and the Quote is the Neat Pounds.*

Ex. What is the Neat Weight of 4 *Par. of* Spaniſh *Tobacco, containing?* viz.

No	C.	q.	l.
4	1	0	04
7	0	3	25
9	1	0	05
3	0	2	26

Tare 14 *per Cent.*
Tret 4 *per* 104 *l.*

3 · 3 · 0

```
         3 . 3 . 04
         112
      8) 424  Groſs
          53  Tare
  104)37100 (356 † Neat
         590
          700
           76
```

Cloff, Is another Allowance of 2 *l.* weight, to Citizens of *London*, on every Draught above 3 Hundred Weight, on ſome Goods, as *Galls*, *Madder*, *Sumack*, *Argol*, &c.

Neat, Is the pure Weight of the Commodity, when all given Allowances are deducted.

As for other Allowances not ſo common, as *Break*, *Damage*, *&c.* the firſt being generally at ſo much *per Cask*, *Bale*, *Bagg*, *&c.* the other ſo much in the whole, can have no difficulty.

Ex. What is the *Neat Weight* of 7 *Hogſheads* Groſs 28 *C.* 1 *q.* 19 *l. Tare* 16 *per Cent. Break* 8 *per Hogſhead Damage* 98 ½ *l.*

```
      C.   q.   l.
  7)  28 . 1 . 19        Groſs
       4 . 0 . 06 . 11   Tare
      24 . 1 . 12 .  5
      —— . 2 . 00 .  0   Break
      23 . 3 . 12 .  5
           3 . 14 .  8   Damage
      22 . 3 . 25 . 13   Neat
```

In Oyl *the* Tare *is of two Sorts.*

(1.) On *Candy* Barrels————————at 29 *l. per Barrel.*
(2.) On uncertain Casks————————at 18 *l. per Cent.*

The *Neat Pounds* of which are found by the foregoing *Rules.* But to reduce thoſe *Pounds* into *Gallons*, Rule. *Multiply thoſe Pounds Neat by* 2, *the Product of which Divide by* 3, *and that Quote by* 5, *give Neat Gallons,* 7 *l.* ½, *making a Gallon.*

Ex. Oyl, *Groſs* 127 *C.* 3 *q.* 10 *l. Tare* 18 *l. per Cent,* how many *Neat Gallons?*

C. q. l.
127 . 3 . 7
112
1615
7) 14315 Groſs
8) 2045
255
2300 Tare
12015 Neat Pounds
2
3) 24030
5) 8010
1602 Gallons

Section IV.

Of PRACTICE.

THE variety of this Rule may be reduc'd to theſe two Heads, *viz.*

(1.) In reſpect to the *Rate* or *Price* of *Goods.*
(2.) The *Given Number*, or *Quantity* of ſuch *Goods.*

Of the *Price*, when
- a Fraction,
- Farthings,
- * An Aliquot part of a *s*, } whether *d.*
- * Not an Aliquot part } or *d.* & *far.*
- Between one *s.* and two
- * An Aliquot part of a *l.* } whether *s.*
- * Not an Aliquot part } or *s.* and *d.*
- * Between one Pound and two, having a Fraction annext.

Of the *Given Number*, when
- * It hath odd W. or M. in it.
- It hath a Fraction annext.
- It is not of the ſame name with that on which the *price* is ſet.
- It is of leſs *Denomination* than it.

Mr. *Kerſy* having in the 2*d.* *Chapter* of his *Appendix* Treated of thoſe Heads, to which an *Aſterisk* is prefixt, I ſhall therefore omit what he hath ſaid.

A Table of ſuch Pence as are an Aliquot part of a Pound.

d.			
2	is	$\frac{1}{120}$	of a Pound.
3		$\frac{1}{80}$	
4		$\frac{1}{60}$	
6		$\frac{1}{40}$	
8		$\frac{1}{30}$	

I. *Of the* Given Price.

1. When it is a *Fraction*,

Rule. *Multiply the Given Number by the Numerator, the Product of which Divide by the Demoninator, gives the Anſwer in the ſame Denomination with the Integer, of which the Price is a Fraction.*

Ex. What comes the *Whitening* of 479 Ells of *Oſnabrigs* to, at $\frac{5}{8}$ of a *Penny* an *Ell?*

479
5

8) 2395

12) 299 . $\frac{3}{8}$

Anſ. 24 . 11 . $\frac{3}{8}$

II. When the Price is *Farthings* only,

Rule. *Reduce the* Farthings *into 3 Pences, which Divided by 80, produce Pounds.*

Ex. What come 6472 *Yards* of Tape to, at 1 Farthing, or 3 Farthings *per* Yard.

12) 6472 at $\frac{1}{4}$ *per* Yard

8|0) 53|9 . 4

6 . 14 . 10

4) 6472 at $\frac{3}{4}$ *per Yard*

8|0) 161|8

20 . 04 . 6

To value the *Remainder* in *Dividing* by any Number conſiſting of a *Digit* and a *Cypher*, as 80 . 60 . 40 *&c.*

Rule. *What Pounds remain, turn into Shillings (taking in the double of the Figure cut off) which Shillings being divided by the Digit in the Diviſor, gives the Odd Shillings; what Shillings remain from thence, turn likewiſe into Pence, which Dividing alſo by the Digit, gives the Odd Pence; and ſo in the Farthings.* Otherwiſe, *What Remains in the Pounds, reckon as Tens, which add to the Unite cut off, the Sum of which Divide by half the Digit, gives Shillings; what Remains from thence, turn into Pence, which Dividing alſo by the half, gives Pence, &c.*

III. When the Price is an Aliquot part of a Shilling, the *Rule* is, *To Divide the Given Number by the ſame part, and the Quote is the Anſwer in Shillings.* But ſince thoſe Parts of a Shilling are alſo the Aliquot parts of a Pound,

Rule, *Divide the Given Number by the ſaid Part of a Pound, the Quote is the Anſwer in Pounds, the Remainder being valued as above.*

Ex. What come 472 *Oz.* to, at 4 *d.* or 8 *d. per Oz.*

6|0)	47|2 at 4 *d. per Oz.*	3|0)	47 2 at 8 *d. per Oz.*
Anſw.	7 . 17 . 4	Anſw.	15 . 14 . 8

IV. If the *Price* is not an *Aliquot Part* of a *Shilling*, as 5, 7, 9, 10, and 11 *Pence*,

Rule. *Take the greateſt even Part of a Pound, and for what is wanting, take Parts out of that Part, which added to the other, gives the* Facit.

Ex. What come 645 *l.* of *Sugar* to, at 7 *d.* or 9 *d. per Pound?*

4|0)	64|5 at 7 *d. per Pound*		3 0)	64|5 at 9 *d. per Pound*
½)	16 . 02 . 6		⅛)	21 . 10
	2 . 13 . 9			2 . 13 . 9
	18 . 16 . 3	Facit	An. *l.*	24 . 03 . 9

V. When any *Farthings* are added to any of the fore-mention'd *Prices*, the

Rule is the ſame with the foregoing.

Ex. What come 735 *Dozen* of *Buttons* to, at 7 *d.* ½ or 6 *per Dozen?*

4|0) 73|5 at 6 *d.* $\frac{3}{4}$ *per Dozen* 4|0) 73|5 at 7 *d.* $\frac{1}{2}$ *per Oz.*
$\frac{1}{8}$) 18 . 07 . 06 $\frac{1}{4}$ 18 . 07 . 06
2 . 05 . 11 . $\frac{1}{4}$ 4 . 11 . 10 : $\frac{1}{2}$
20 . 13 . 05 . $\frac{1}{4}$ 22 . 19 . 04 . $\frac{1}{2}$

VI. If it were here required at ſuch a *Profit* in the *Shilling*, to know what *Profit per Cent.*

Rule. *Divide* 100 *l. by the Parts that the ſaid Profit is of a Shilling, the total of which is the Anſwer.*

Ex. What *Profit per Cent. when I Gain* 4 *d.* $\frac{1}{2}$ or 7 *d. per Shilling?*

4 $\frac{1}{3}$ 100 4 $\frac{1}{3}$ 100 at 7
$\frac{1}{2}$ $\frac{1}{8}$ 33 : 06 : 08 3 $\frac{1}{4}$ 33 : 06 : 08
4 : 03 : 04 25 : 00 : 00
37 : 10 : 00 58 : 06 : 08

Otherwiſe 4 *d.* $\frac{1}{2}$ being the $\frac{3}{8}$, and 7 *d.* the $\frac{7}{12}$ of a *s.* Multiply a 100 by the Numerator, and Divide that Product by the Denominator, according to the firſt Rule, gives the Facit, as

100 100
3 7
8) 300 12) 700
Anſwer 37 . 10 Facit 58 . 06 : 08

VII. When the *Price* is any Number of *Pence*, under two *Shillings*, as, 14, 15, 16, 18, 21, 22, *Pence.*

Rule. *Reduce the ſaid Prices either into* 2 *d.* 3 *d.* 4 *d.* 6. *d.* or 8 *pences and thoſe into Pounds at one Operation, by the third Rule.*

Gallons *Buſhels*
2751 at 14 *d. per Gal.* 375 at 15 *d. per Buſh.*
7 5
12|0) 1925|7 2 Pences 8|0) 187|5 3 Pences
Facit 160 . 09 . 06 Facit 23 . 08 . 09

Dozen

Dozen	*Pair*
297 at 16 *d. per Dozen*	736 at 18 *d. per Pair*
2	3
3|0) 59|4	4|0) 220|8
Anſw. 19 . 16	Facit 55 . 04

L.	*Ells* at 22 *d.* per Ell.
715 at 21 *d. per l.*	917
7	11
8|0) 500|5	12|0) 1008|7
Facit 62 . 11 . 3	Anſw. 84 . 01 . 02

For 13, 17, and 19 pence.

Rule. *Multiply the Given Number by the ſaid Pence, (as Taught in Multiplication) which Product dividing by* 12 *and* 20, *gives the Facit.*

Examples.

463 *l.* of *Copper, at* 17 *d. l.*	479 *Ells of Long Cloth, at* 19 *d.*
463	479
17	19
12) 7871	12) 9101
655 . 11	2|0) 75|8 . 5
Anſw. 32 . 15 . 11	37 . 18 . 5

VIII. If any of the laſt foregoing *Prices* had *Farthings* annext,

Rule. *Then take Parts for the Pence and Farthings above a Shilling, out of a Shilling, which added to the Given Number, the total is the Anſwer in Shillings*; as

Ex. What do I ſpend *per Annum,* at 19 *d.* $\frac{1}{2}$. or 18 *d.* $\frac{3}{4}$ *per Day?*

365 *Days* at 19 *d.* $\frac{1}{2}$ each.

6 $\frac{1}{2}$ 365
1 $\frac{1}{2}$ $\frac{1}{4}$ 182 . 06
45 . 07 . $\frac{1}{2}$
2|0) 59|3 . 01 . $\frac{1}{2}$
29 . 13 . 01 . $\frac{1}{2}$

365 Days at 18 *d.* $\frac{3}{4}$ *per Day.*

6 $\frac{1}{2}$ 365
$\frac{3}{4}$ $\frac{1}{8}$ 182 . 06
22 . 09 . $\frac{3}{4}$
2|0) 57|0 . 03 . $\frac{3}{4}$
28 . 10 . 03 . $\frac{3}{4}$

IX. If the *Price* is any *Even Number* of *Shillings*, as, 2, 4 6, 8, 12, 14, 16, 18, 22, *&c.*

Rule. *Multiply the Given Number by the half of the Price, ſetting the Double of the Unite Figure for Shillings.*

Ex. What come 379 *Yards* of *Broad-Cloth* to, at 18 *s.* or 22 *s. per Yard?*

379 at 18 *s.*
9
341 . 2

379 at 22 *s.*
11
416 . 18

See Mr. Kerſy's *5th & 6th Rule of the ſecond Chapter.*

X. When the *Price* is an *Odd Number* of *Shillings*,

Rule. *Multiply the Given Number by the ſaid Shillings, Dividing the Product by* 20.

Ex. What come 796 *Small Groſs* to, at 13 *s.* or 17 *s. per Groſs?*

796 at 13 *s.*
13
2|0) 1034|8
517 . 08

796 at 17 *s.*
17
2|0) 1353|2
676 . 12

Or rather according to Mr. *Kerſey*'s 8th Head, in the Rule of *Practice.*

XI. When the *Price* is *Shillings* and *Pence*, and ſuch *Shillings* and *Pence* are the ſame Figure,

Rule, *Multiply the Given Quantity by the Shillings, taking always the* $\frac{1}{12}$ *of the Product for the Pence, the total of which dividing by* 20, *gives the Anſwer in Pounds.*

Ex. What come 731 *Ells* of *Holland* to, at 7 *s.* 7 *d.* or 11 *s.* 11 *d. per Ell?*

731 *Ells* at 7 *s.* 7 *d.*	731 *Ells* at 11 *s.* 11 *d.*
7	11
$\frac{1}{12}$) 5117	$\frac{1}{12}$) 8041
426 . 05	670 . 01
2|0) 5541|3 . 05	2|0) 8711|1 . 01
277 . 03 . 05	435 . 11 . 01

XII. When the *Price* is between one *Pound* and two,

Rule, *Work after the manner of the 8th Rule,* i. e. *only for the Money above a Pound, adding thereto the Given Quantity.*

Ex. What come 224 *l.* of *Cocheneal* to, at 27 *s.* 6 *d. per l.*

s. *l.*	But the Price being $\frac{11}{8}$ of a *l.* work, *viz.*
5 $\frac{1}{4}$) 224	224
2 . 6 $\frac{1}{2}$) 56	8) 2464
28	308 Anſwer
308	

XIII. When the *Price* hath a *Fraction* annext,

Rule, *Work for the Pence according to the foregoing Rules, and for the Fraction, either value it as directed in the firſt Rule: Or if an Aliquot part of the Money foregoing, take ſuch part for it.*

Ex. (1.) What come 479 *l.* of *Tobacco* to, at 22 *d.* $\frac{5}{8}$ *per l.*

479	
11	479
12|0) 526|9	3
43 . 18 . 2	8) 1437
14 . 11 . $\frac{5}{8}$	12) 179 . $\frac{5}{8}$
Anſwer 44 . 13 . 01 . $\frac{5}{8}$	14 . 11 . $\frac{5}{8}$

Ex. (2.) How much *Sterling* for 7439 French Crowns Exchange, at 55 *Pence* $\frac{5}{8}$ *per Crown?* 7439

s.	d.	$\frac{1}{8}$	7439
2	6	$\frac{1}{12}$	929 . 17 . 06
1	8	$\frac{1}{4}$	619 . 18 . 04
	5	$\frac{1}{8}$	154 . 19 . 07
	$\frac{5}{8}$		19 . 07 . 05 . $\frac{3}{8}$
Anſwer			1724 . 02 . 10 $\frac{3}{8}$ Sterling

II. *Of the* Given Quantity.

I. When it hath *Odd Weight* or *Meaſure* added,

Rule, *Value the whole Number as before Directed, then take Parts for ſuch Odd Weight or Meaſure out of an Integer; Dividing the Given Price by ſuch Parts, the total of which added to the value of the whole Number gives the amount.*

Ex. What come 294 *C.* 2 *q.* 18 *l.* to, at 10 *s.* 4 *d.* *Hundred?*

	294		s. d.
	5	$\frac{1}{2}$	10 . 4
$\frac{1}{20}$)	147 . 00	$\frac{1}{7}$	5 . 2
	4 . 18	$\frac{1}{8}$	1 . 5 . $\frac{3}{2}$
	06 . 09 . $\frac{1}{2}$		0 . 2 . 0
	152 . 04 . 09 . $\frac{1}{2}$		6 . 9 . $\frac{1}{2}$

II. When the *Given Number* is not of the ſame *Name* with the *Integer* on which the *Price* is,

Rule, *It muſt be reduc'd into the ſame according to the Rules of Reduction.*

Ex. What come 49 Thouſand 10 *l.* 11 *oz.* of Silk to, at
to, at 17 *s.* 4 *d. per Hundred?* 2 *s.* 8 *d. per Oz.*

490	10 . 11
9	16
441 . 00	3)171
16 . 06 . 8	57. Anſw. 22 *l.* 16 *s.*
424 . 13 . 04	22 , 8

III. When it hath a Fraction annext,

Rule. *Value the whole Number according to the Rules aforegoing;* *and for the* Fraction, *Multiply the Price by the Numerator, and that Product divided by the Denominator, gives the value of the* Fraction, *which added to the value of the whole Number, gives the Anſwer.*

Ex. What come 372 *Ells* $\frac{3}{5}$ of *Bag Holland* to, at 5 *s.* 10 *d.*

$\frac{1}{4}$ 372
$\frac{1}{6}$ 93
15 . 10
3 . 6
108 . 13 . 6

5 . 10
3
5) 17 : 06
3 . 06

IV. When the *Given Quantity* is of *Leſs Denomination* than that on which the *Price* is Given,

Rule. *Take parts for the Leſs Name out of the Greater, dividing the Price by ſuch parts.*

Ex. At 7 *l.* 10 *s.* per *Tun*, what per *Barrel?*

7) 7 . 10
Anſw. 1 . 01 : 5 . $\frac{1}{7}$

How much a *Gallon*, at 38 *s.* 3 *d.* per *Hogſhead?*

9) 38 . 3
7) 4 . 3
Anſw. 0 . 7 . $\frac{2}{7}$

What come a *Cheſhire Cheeſe* to weight 44 *l.* at 37 *s.* 9 *d.* *Hundred?*

$\frac{1}{4}$ 37 . 9
$\frac{1}{7}$ 9 . 5 . $\frac{1}{4}$
4 . 8 . $\frac{1}{3}$
Anſw. 14 . 9 . $\frac{3}{4}$

At 19 *l.* 7 *s.* 8 *d.* *Hundred*, how much a *Pound?*

		l.	*s.*	*d.*	
16	$\frac{1}{7}$	19	07	08	
2	$\frac{1}{8}$	2	15	04	$\frac{1}{2}$
1	$\frac{1}{2}$	0	06	11	0
	Anſw.	0	03	05	$\frac{1}{2}$

The Quantities *of Particular Goods in Whole-ſale Trade, found in the* Book of Rates, *with Conciſe and Ready ways of Caſting up the ſame.*

I. Goods Sold by Six-Score to the Hundred.

Lamb-Skins, Barlings, Balks, Great, Middle, and Small; Clap-Boards, Pipe-Boards, Bow-Staves, Capravens, Deal-Boards, Spars *of all Sorts*, Cod, Cole, Ling; *and all Sorts of* Stock Fiſh, *with many Sorts of Linen, viz.* Hamborough, Sleecia, Iriſh, Muſcovy, Weſtphalia, Hannover, *&c.*

To Caſt up Goods of this Quantity,

Rule, *Take half the Pence of the Price of one, which gives you the Value of* 120 *in Pounds, Shillings*, &c.

$\frac{120}{240}$ being the ½ of a *l. Sterling.*

Ex. What come 120 *Ells* of *Canvas* to, at 5 *d.* ½ per *Ell*?

$$2)\ \frac{5\ .\ \frac{1}{2}}{2\ .\ 15}$$ Anſwer 2 *l.* 15 *s.*

Ex. (2.) At 7 . 15 *s.* per *Hundred* of *Deals*, how much each?

$$\frac{\quad\quad\cdot 2}{15\ .\ 10}$$ Anſwer 15 *d.* ½.

N. B. That you eſteem every *Penny* that Remains (in the Firſt Example) as 20 *s.* and every 20 *s.* as one *Penny*, (in the Second.) As

In Example, the Firſt there Remains one *Penny*, which reckoning 20 *s.* and accordingly the *Half-penny* in the Given Price for 10 *s.* both which making 30 *s.* which Dividing by 2, gives 15 *s.* and ſo the Anſwer is found to be 2 *l.* 15 *s.*

II. Goods Sold by the Thouſand, are

Paſte-Boards *for* Books, Tennis-Balls, Limons, Oranges, Teazels, Flanders Paving *and* Pan-Tyles, Lantern-Horns, Barrel-

Hoops, *and* Boards, Quills, Lamperns, Squirrel-Skins, Ox-Bones, Yards *of* Liſt, *&c.*

To Caſt up Goods of this Quantity,

Rule, *Multiply the Pence that one coſts by 50, and the Product divided by 12, gives the Value of a Thouſand.*

$\frac{1000}{240}$ being the $\frac{50}{12}$ or $\frac{25}{6}$ of 20 *s.*

Ex. (1.) What come 1000 of *Flanders Tyles* to, at 3 *d.* each?

$$\begin{array}{r} 3 \\ 50 \\ \hline 12)\,150 \\ \hline \text{Anſwer } l.\ 12\ .\ 10 \\ \hline \end{array}$$

Ex. (2.) If a 1000 of Squirrel-Skins coſt 11 *l.* 05 *s.* what per Skin?

$$\begin{array}{r} 11\ .\ 05 \\ 12 \\ \hline 5|0)\,1315\ .\ 00 \\ \hline 2\ .\ \frac{1}{2}\ \text{per Skin.} \end{array}$$

III. Goods Sold by the *Great Groſs*, containing 12 *Small Groſs*, or 144 *Dozen*, viz.

Buttons *of* Metal, Glaſs, Thread, Silk, *or* Hair, Beads *of* Bone, Box, Glaſs *or* Wood, Cap-Hooks, Cheſs-men, Tobacco-Pipes, Combs *of* Light *or* Box-Wood, Thread *or* Silk Points, Playing-Cards, *&c.*

Goods of this Quantity are Caſt up, *viz.*

Rule, *Multiply the Pence of the Price of one Dozen by 3, and Divide the Product by 5, gives the Value of a Great Groſs in Pounds,* &c.

$\frac{144}{240}$ being the $\frac{3}{5}$ of a Pound Sterling.

Ex. (1.) What come a *Great Groſs* of Silk Buttons to, at 8 *d.* ½ per *Dozen?*

$$\begin{array}{rl} 8\ .\ \frac{1}{2} & \\ 3 & \\ \hline 5)\ 25\ .\ \frac{1}{2} & \text{Anſwer } 5\ l.\ 2\ s. \\ \hline 5\ .\ 2 & \\ \hline \end{array}$$

Ex. (2.) What come 1 Dozen to, when a Great Groſs coſt 7 *l.* 8 *s.*

$$\begin{array}{rl} 7\ .\ 8 & \\ 5 & \\ \hline 3)\ 37\ .\ 0 & \text{Anſwer } 12\ d.\ \frac{1}{4}. \\ \hline 12\ .\ \frac{1}{4} & \\ \hline \end{array}$$

IV. Goods Sold by the *Small Groſs* of 12 *doz.* viz.

Tobacco *and* Pepper Boxes, Neſt *and* Tinder-Boxes, Buckles *of ſome Sorts*, Waſh-Balls, Ink-horns, Bodkins, *ſeveral Sorts of* Knives, Comb-Caſes, *&c.*

Goods of this Quantity may be Caſt up by the laſt preceeding Rule, ſeeing there are as many particulars in a Small Groſs, as Dozens in a Great one.

V. Goods Sold by *Five-Score* to the *Hundred.*

Annotto, Capers, Saflower, Thrums, Ginger, Cloves, Indico, Croſs-Bow-Thread, Pack-Thread, Kids *or* Goats-Hair, Quick-Silver, Cotton-Wool, *Engliſh* Hard Wax, Braſs *and* Latin Work, *as* Chafing-Diſhes, Andiorns, Laver-Cocks, *&c.*

To caſt up which Goods,

Rule. *Multiply the Pence of the Price of one l. Weight by* 5, *dividing the Product by* 12.

$\frac{100}{240}$ being the $\frac{5}{12}$ of a *l.*

Ex. (1.) What come 100 Weight of *China Roots* to, at 15 *d* per *l.*

$$\begin{array}{rl} 15 & \\ 5 & \\ \hline 12\ \ 75 & \text{Anſwer } 6\ l.\ 5\ s. \\ \hline 6\ .\ 5 & \end{array}$$

Ex. 2. What comes 1 Pound to, when 100 Pound coſt 14 *l.* 15 *s.*

l. *s.*
14 . 15
12
5) 177 . 00
Anſwer 35 . $\frac{2}{5}$ or 2 *s.* 11 *d.* $\frac{2}{5}$

To Value Goods ſold by the *common Hundred* of 112 *Pound* to the *Hundred.*

Rule. *Multiply the Pence that one Pound coſts by* 7, *and the Product divide by* 15, *or multiply by* 14 *and divide by* 30 *gives the Value* $\frac{112}{240}$, *being the* $\frac{14}{30}$ *or* $\frac{7}{15}$ *of a Pound.*

Ex. 1. What comes a *Hundred weight* to, at 8 *d.* $\frac{3}{4}$ per *Pound?*

8 . $\frac{3}{4}$
7
15) 61 . $\frac{1}{4}$ *d.*
4 . 1 . 8.

Ex. 2. At 9 *l.* per *Hundred* what comes 1 *Pound?*

9
15
7) 135
19 . $\frac{1}{4}$

To caſt up Goods of this Quantity in mind, ſee Mr. *Kerſey*'s 18th *Rule* of the 2d *Chapter.*

Now if it be further required to know what any Number (not exceeding 5 Score) of the aforeſaid *Quantities* in wholeſale Trade (or any other) amount to at any *Rate*, the moſt practical and eaſieſt Way is by the following Rule.

Rule. *Multiply the Price of one by the Parts of the given Quantity (as taught in Multiplication.)*

Examples.

What comes 12 *C.* ½ at 7 *l.* 11 *s.* 8 *d.* per *Hundred?*

```
   7 . 11 . 8
          12 . ½
 ------------
  91 . 00 . 0
   3 . 15 . 10
 ------------
  94 . 15 . 10
```

At 4 *s.* 7 *d.* ½. per *Groſs,* what comes 46 *Groſs?*

```
          4 . 7 . ½
                  5
 ------------------
         23 . 01 . ½
                  9
 ------------------
 10 . 08 . 01 . ½
      04 . 07 . ½
 ------------------
 10 . 12 . 09 . 0
```

What comes 2900 to, at 2 *l.* 14 *s.* 7 *d.* per *Hundred?*

```
   2 . 14 . 07
             3
 -------------
   8 . 03 . 09
            10
 -------------
  81 . 17 . 06
   2 . 14 . 07
 -------------
  79 . 02 . 11
```

33 *Hundred* at 19 *s.* 7 *d* per *Hundred.*

```
       19 . 07
            11
 -------------
  10 . 15 . 05
             3
 -------------
  32 . 06 . 03
```

If inſtead of knowing the *Value* or *Worth* of *Goods* at a certain *Price,* It be required to know what *Quantity* of *Goods* can be Bought for a certain Sum of *Money.*

If the *Price* be even *Shillings.*

Rule. *Add a Cypher to the Pounds, and divide by ½ the Price.*

Ex. 1. How many *lb.* *oz.* and *drams,* can I buy for 479 *l.* at 6 *s.* per *lb?*

```
          3) 4790    oz.  dra.
 Anſwer, lb. 1596 . 10 . 10 . ⅔.
```

Ex. 2. At 8 *s.* per *Yard,* how many *Yards, Quarters* and *Nails* can I buy for 315 *l?*

```
 4) 3150               Yards  Quarters.
    ----
    787 . ½      Anſw. 787  .  2
```

At 14 *s.* . 3 *d.* per Yard how much per Ell? Add $\frac{1}{4}$ part.

4) 14 . 3
3 . 6 . $\frac{3}{4}$
Anſw. 17 . 9 . $\frac{3}{4}$

What coſt 1 Pound when $\frac{2}{3}$ coſt 16 *s* . 8 *d*? Add $\frac{1}{2}$ the price.

16 . 8
8 . 4
Anſw. 25 . 0

Vhat comes 47 Pound to, when $\frac{3}{4}$ coſt 6 *d.* *i. e.* 8 *d.* per Pound?

3|0) 4|7
Anſw. 1 . 11 . 4

How much can I buy for 11 *d.* when 24 *lb.* 11 *oz.* coſt 11 *s*?

1$\frac{1}{2}$) 24 . 11 *dr.*
Anſwer 2 . 00 . 14 $\frac{2}{3}$.

If 22|0 Yards coſt 57 *l.* what is that per Yard?

11) 57 *d.*
Anſwer 5 . 2 . 1$\frac{2}{1}$.

Here I divide 57 *l.* by 11, ſeeing there is the like reaſon etween 11 and 57, as there would have been between 220 nd 1140, the Product of 57 by 20.

SECTION. V.

Of *Simple* INTEREST.

TO find the *Intereſt* of any *Sum* at any *Rate*, for any *Time* given.

I. By Months.

In this the *Year* is taken as containing 12 *Months* of 30 *Days* each; which wants $_{1}\frac{5}{2}$ + of a Day, and therefore the preciſe *Intereſt* cannot be found by this method.

Yet becauſe 'tis ſo much in uſe among Traders, ſo little differing from truth, and ſo ready in operation, I ſhall give three ſeveral ways of performing it.

The Firſt and uſual *Way*.

Rule, *Multiply the Principle by the Rate of Intereſt* per Cent per Annum, *and the Product (caſting off two Figures on the right Hand) is the Intereſt for a Year: From whence 'tis found for any Time required.*

Ex. 1. What is the *Intereſt* of 479 *l.* 18 *s.* for 2 *Years*, at 4 *l.* ½. per *Cent.* per *Annum?*

```
    479 . 18
         4 ½
   ---------
   1919 . 12
    239 . 19
   ---------
  21|59 . 11
    |20
  -------
  11|91
    |12                  l . s . d.
  -------
  10|92                 21 . 11 . 10 . ¾
    | 4                                2
  -------          ----------------------
   3|68          Anſw. 43 . 03 . 09 . ½
```

Ex. 2. At 6 per *Cent.* per *Annum,* what is the *Intereſt* of 571 *l.* 15 *s.* for 8 *Months?*

571 . 15
6
34|30 . 10
|20
6|10
|12
1|20

6 ½
2 3

l. s. d.
34 : 06 . 01
17 . 03 . 00 . ½
5 . 14 . 04 . 0
Anſw. 22 . 17 . 04 . ½

Note, By this *Rule, Broakage, Commiſſion, Proviſion, Storage, Inſurance,* &c. are wrought, having no reſpect to *Time.*

Ex. 1. Unto what comes the *Commiſſion* of 259 *l.* 17 *s,* 11 *d.* ½ at 2 ½ per *Cent.*

259 . 17 . 11 . ½
2
½ 519 . 15 . 11 . 0
129 . 18 . 11 . ¾
6|49 . 14 . 10 . ¾
|20
9|94
|12
11|38
| 4
1|55

Ex. 2. At 1 ¼ per *Cent.* what comes the *Broakage* of 198 *l.* 11 *s.* 7 *d.*?

4| 198 . 11 . 7
49 . 12 . 10 . ¾
2|48 . 4 . 5 . ¾
|20
9|64
|12
7|73
| 4
2|95

Anſw. 2 *l.* 9 *s.* 7 *d.* ½

But Queſtions of this nature are more conciſely and readil wrought by this

Rule, *Divide the given Sum by the Aliquot part that the Rat of the Commiſſion or Broakage,* &c. *is of* 100 *l.* As for In ſtance, in the laſt Examples.

Ex. 1. The *Commiſſion* being $2\frac{1}{2}$ per *Cent.* which is the 40t part of 100 *l.* Therefore

$$\begin{array}{l} 4|0)\ \underline{25|9\ l.\ 17\ s.\ 11\ d.\ \frac{1}{2}.} \\ \textit{Anſw.}\ 6\ .\ 09\ .\ 11\ .\ \frac{1}{4}. \end{array}$$

Ex. 2. The *Broakage* being $1\frac{1}{4}$ per *Cent.* which is the 80t part of 100 *l.* Therefore

$$\begin{array}{l} 8|0)\ \underline{19|8\ l.\ 11\ s.\ 7\ d.} \\ \textit{Anſw.}\ 2\ .\ 09\ .\ 07\ .\ \frac{1}{2}. \end{array}$$

The Second way.

Rule, *Multiply the Principle, by the Intereſt of* 100 l. *for th Time required, and the Product (ſtriking off two Figures on th Right-hand) is the Anſwer.*

Ex. 1. What is the Intereſt of 479 *l.* 18 *s.* for two Year at $4\frac{1}{2}$ per *Cent.* per *Ann.?*

The *Intereſt* of 100 *l.* for two Years, at $4\frac{1}{2}$ per *Cent.* is 9 *l.* Therefore

$$\begin{array}{r|l} \multicolumn{2}{r}{479\ .\ 18} \\ \multicolumn{2}{r}{9} \\ \hline 43 & 19\ .\ 02 \\ & 20 \\ \hline 3 & 82 \\ & 12 \\ \hline 9 & 84 \\ & 4 \\ \hline 3 & 36 \end{array}$$

Anſw. 43 *l.* 03 *s.* 09 *d.* $\frac{3}{4}$.

Ex. 2.

Ex. What is the *Intereſt* of 571 *l.* 15 *s.* for 8 *Months*, at 6 per *Cent.* per *Annum?*

The *Intereſt* of 100 *l.* for 8 *Months*, at 6 per *Cent.* is 4 *l.* Therefore

$$\begin{array}{r|l} & 571\,l.\ 15\,s. \\ & \qquad\ \ 4 \\ \hline 22 & 87\ .\ 00 \\ & 20 \\ \hline 17 & 40 \\ & 12 \\ \hline 4 & 80 \\ & \ \ 4 \\ \hline 3 & 20 \end{array}$$

Anſw. 22 *l.* 17 *s.* 04 *d.* ¾.

Note, If it be for 3 Months, at 5 per *Cent.* per *Annum*, then you need only *add* ¼ *of the Principle to it ſelf, working as before directed*, &c.

The Third way.

Rule, *Since the Intereſt of any Principle,*

at $\left\{\begin{array}{c}5\,l.\\6\\4\\3\end{array}\right\}$ per *Cent.* for $\left\{\begin{array}{c}12\\10\\15\\20\end{array}\right\}$ Months.

is juſt ſo many Shillings as it hath Pounds. Therefore

(1) What is 379 *l.* 15 *s.* for 12 Months, at 5 per *Cent?*
(2) What is 416 *l.* 12 *s.* 6 *d.* for 10 Months, at 6 per *Cent?*

2|0) 37|9 . 15
Anſw. 18.19 . 9

2|0) 41|6 . 12 . 6 *d.*
Anſw. 20.16 . 7½.

(3) What is 427 *l.* 13 *s.* 9 *d.* for 20 Months, at 3 per *Cent?*
(4) What is 927 *l.* 11 *s.* 3 *d.* for 15 Months at 4 per *Cent?*

2|0) 42|7 . 13 . 9
Anſw. 21 . 7 . 8½.

2|0) 92|7 . 11 . 3
Anſw. 46 . 7 . 6½.

Hence, To find the *Intereſt* of a given *Principle*, for any other *Time*.

Rule, *Take parts, for the Time required, out of the Time affixed.*

Thus, If the *Intereſt* of 379 *l.* 15 *s.* for 12 Months, at 5 per *Cent.* is ſo many *Shillings*, then for 6 Months it is the ½ of ſo many *Shillings*, or ſo many *Six-pences*; and for 4 *Months* the ⅓ of it, or ſo many Groats, as the *Principle* hath *Pounds*.

Ex. 1. What is the *Intereſt* of 279 *l.* 11 *s.* for 4 Months, at 5 per *Cent?*

2|0) 27|9 . 11
3) 13 . 19 . 6½.
Anſw. 4 . 13 . 2

Ex. 2. What is the *Intereſt* of 197 *l.* 11 *s.* for 2 Months, at 6 per *Cent?*

2|0) 19|7 . 11
5) 9 . 17 . 6½.
Anſw. 1 . 19 . 6

Ex. 3. What is 491 *l.* 17 *s.* 10 *d.* for 5 Months, at 4 per *Cent?*

2|0) 49|1 . 17 . 10
3) 24 . 11 . 10¾.
Anſw. 8 . 3 . 11½.

Ex. 4. What is 279 *l.* 11 *s.* 8 *d.* for 7½ Months, at 4 per *Cent.*

2|0) 279 . 11 . 8
2) 13 . 19 . 7
Anſw. 6 . 19 . 9½.

For what odd Money is adjoyn'd to the *Principle*, value every 20 *s.* as 12 *d.* and leſſer Sums in proportion.

SECTION VI.

II. Intereſt for Days.

To find the exact number of *Days*, between any two given *Times*, without any *Book* or *Table*.

This is done, by carrying in Memory that old Diſtich, viz.

Thirty Days hath *September*, *April*, *June* and *November*.
February hath 28 alone; each of the reſt hath *Thirty one*.

And according to this reckoning the Year contains 365 Days; therefore, for Inſtance,

To find the number of *Days* between the 7th of *June* 1701, and the 15th of *September* 1703. Say thus,

365
2

From the 7th of *June* 1701 to the 7th of *June* 1703 is 730 or two Years.

From the 7th of *June* 1703 to the end of the ſame Month is 23

Then in { *July* —— 31
Auguſt —— 31 }

From which, to the 15th of *September*, is —— 15

So that from the 7th of *June* 1701, to the 15th of *September* 1703, is } Days 830

Note, *That* 1 *Day muſt be added to every* 4 *Years contain'd in the Time*. For every fourth Year, being *Leap-year*, *February* hath then 29 Days.

After this manner may any number of Days, even to ½ or ¼ of a Day, be exactly and readily found. Now

To find the *Intereſt* of any *Sum*, at any *Rate*, for any number of Days, obſerve theſe two General ways.

Firſt Rule, *Multiply the Pence of the Principle by the number of Days, and the Product by the Rate of Intereſt for a Dividend, which divide by* 36500, *the Quotient is the Anſwer in Pence.*

Otherwiſe, *Take the Product of the Pence of the Principle, Multiply'd by the Days, for a Dividend, and the Quotient of* 36500, *divided by the Rate of Intereſt, for a Diviſor, and the Quotient of that Diviſion is the Anſwer.*

Ex. I.

Ex. 1. What comes the Intereſt of 192 *l.* 14 *s.* 10. for 830 Days,at 5 per *Cent.* per *Annum*?

```
                       20
                     ----
                     3854
                       12
                    -----
                    46258
                      830
                  -------
                  1387740
                  370064
                 --------
                 38394140
                        5
                 --------
365|00) 1919707|00 (5259
        947          43|8 . 3
        2170         21.18 . 3 Anſw.
        ----
         3457
         17200
```

*Ex.*2. What comes the Intereſt of 311 *l.* 13 *s.* 4 *d.* for 315 Days at 6 per *Cent.* per *Annum?*

```
                   20
                 ----
                 6233
                   12
                -----
                74800
                  315
              -------
              1122000
             224400
             --------
6083) 23562000 (3873
        53130     32|2 . 9
        44660     --------
                  16 . 2 . 9
        20790
        -----
         2541
```

The ſecond, which I received from my Friend Mr. *Ralph Snow*, is.

Rule, *Multiply the Pence in the given Principle, by the number of Days, divide the Product always by* 7300, *and the Quotient is the Intereſt, in Pence, of that Principle, for that Time, at* 5 *per* Cent. *per Annum.*

Thus the Intereſt of 192 *l.* 14 *s.* 10 *d.* for 830 *d.* at 5 per *Cent.*

20
3854
12
46258
830
1387740
370064
73|00) 383941|40 (5259 Intereſt in Pence at 5 per *Cent.*
189
434
691
34

Whence, To find the *Intereſt* of the ſame *Principle*, for the ſame *Time*, at the *Rate* of 3, 4, 6, 7, &c. per *Cent.*

Add ſo many Fifths as the Rate of Intereſt exceeds 5; *Or Subtract ſo many Fifths as it is leſs than* 5.

The Fifths of any Number is readily found, by *Multiplying double the Numerator, cutting off the unite Figure of the Product.*

Becauſe ſo many *Fifths* of any Number is the ſame with twice ſo many *Tenths* of that Number.

Now 5259 *d.* being the Intereſt of 192 *l.* 14 *s.* 10 *d.* for 830 Days, at 5 per *Cent.* (as in the laſt Precedent) let it be required, to know the Intereſt of the ſame Principle, for the ſame time, at any other Rate, ſuppoſe 3, 4 $\frac{1}{2}$, or 6 per *Cent.* per *Annum*, then, *viz.*

5259
4
Ex. 1. $\frac{4}{10}$, or $\frac{2}{5}$, being ſubtracted reſts 2103 . 6
The Intereſt likewiſe in Pence, at 3 per *Cent.* 3155 . 4

Ex. 2. One Tenth ſubtracted, reſts the Intereſt at $4\frac{1}{2}$ per *Cent.*

```
 5259
  525 . 9
 ---------
 4733 . 1
```

Ex. 3. $\frac{2}{10}$, or $\frac{1}{5}$, being added, gives the Intereſt at 6 per *Cent.*
And the like for any other Rate whatever.

```
 5259
 1051 . 8
 ---------
 6311 .
```

Of Exchequer Notes and Tallies.

Chequer Notes of five and twenty Pound, did formerly bear five Farthings, but now a Penny per Day Intereſt.

Tallies have been at 6, but now at 5 per *Cent.* per *Ann.* the Intereſt payable every 3 Months.

Example. Unto what comes the Intereſt of two 25 *l.* Notes from the 27 of *November* 1702 to this Day, being the 4th of *January* $170\frac{3}{4}$?

The Days found, according to the direction given in the beginning of this Section, are 403, which at 2 *d.* per Day Intereſt, is 3 *l.* 7 *s.* 2 *d.* as, *viz.*

```
12|0) 40|3
      -----
     3 . 7 . 2
```

Ex. 1. What is the quarterly Intereſt of a Tally, *viz.* Numb. 2704, ſtruck *June* 11, 1702 for 579 *l.* at 6 per *Cent?*

$\frac{1}{2}$) 579

```
½) 579
     289 . 10
   ----------
   8 68 . 10
       |20
   ----|---
    13 |70
       |12
   ----|---
     8 |40
       | 4
   ----|---
     1 |60        Anſw. 8 l. 13 s. 8 d. ¼
```

Ex. 2. What comes the Intereſt, until the 10th of *January* 170$\frac{3}{4}$, of two Tallies, the Number, Principle and Time, of whoſe Orders are, *viz.*

N°. 3081, from *Auguſt* 5. 1702, for 914 *l.* 5 *s.* at 6 per *Cent.* per *Annum*?

```
                           20
                        -----
                        18285
                           12
                      -------
                       219420
                          523 Days.
                      -------
                       658260
Anſw. 78 l. 12 s.     438840
                    1097100
              ----------------------
              73) 1147566|60 (15720
                    417           3144
                     525    12)18864
                      146    ---------
                       --      157|2
                        6     -------
                              78 . 12
```

N°. 5792, ſtruck *May* 22. 1703, for 379 *l.* 10 *s.* at 5 per *Cent.* per *Annum.*

```
                 20
               ----
               7590
                 12
              -----
              91080
                233
             ------
             273240
            273240
           182160
          ---------
      73) 212516|40 (2907
            662            24 2 . 3
               516         --------
                (5         12.2.3
```

TABLES

OF

Simple Intereſt,

Serving at any Rate for any Time.

SECTION VII.

Days	At 3 per *Cent.*	At 4 per *Cent.*	At 5 per *Cent.*
1	.000082192	.000109589	.000136986
2	.000164383	.000219178	.000273972
3	.000246575	.000328767	.000410959
4	.000328767	.000438356	.000547945
5	.000410959	.000547945	.000684931
6	.000493150	.000657534	.000821918
7	.000575342	.000767123	.000958904
8	.000657534	.000876712	.001095890
9	.000739726	.000986301	.001232877
10	.000821918	.001095890	.001369863
11	.000904109	.001205479	.001506849
12	.000986301	.001315068	.001643835
13	.001068493	.001424657	.001780822
14	.001150685	.001534246	.001917808
15	.001232876	.001643835	.002054794
16	.001315068	.001753424	.002191781
17	.001397260	.001863013	.002328767
18	.001479452	.001972602	.002465753
19	.001561644	.002082192	.002602740
20	.001643835	.002191781	.002739726
21	.001726027	.002301370	.002876712
22	.001808219	.002410959	.003013698
23	.001890412	.002520548	.003150685
24	.001972602	.002630137	.003287671
25	.002054794	.002739726	.003424657
26	.002136986	.002849315	.003561644
27	.002219178	.002958904	.003698630
28	.002301369	.003068493	.003835616
29	.002383561	.003178082	.003972603
30	.002465753	.003287671	.004109589
31	.002547945	.003397260	.004246575
32	.002630137	.003506849	.004383561
33	.002712328	.003616438	.004520548
34	.002794520	.003726027	.004657534
35	.002876712	.003835616	.004794520
36	.002958904	.003945205	.004931507

Days	At 6 per *Cent.*	At 7 per *Cent.*	At 8 per *Cent.*
1	.000164383	.000191781	.000219178
2	.000328767	.000383561	.000438356
3	.000493150	.000575342	.000657534
4	.000657534	.000767123	.000876712
5	.000821918	.000958904	.001095890
6	.000986301	.001150685	.001315068
7	.001150685	.001342466	.001534246
8	.001315068	.001534246	.001753424
9	.001479452	.001726027	.001972603
10	.001643835	.001917808	.002191781
11	.001808219	.002109589	.002410959
12	.001972602	.002301370	.002630137
13	.002136986	.002493150	.002849315
14	.002301370	.002684931	.003068493
15	.002465753	.002876712	.003287671
16	.002630137	.003068493	.003506849
17	.002794520	.003260274	.003726027
18	.002958904	.003452054	.003945205
19	.003123287	.003643835	.004164383
20	.003287671	.003835616	.004383561
21	.003452055	.004027397	.004602739
22	.003616438	.004219178	.004821917
23	.003780822	.004410959	.005041095
24	.003945205	.004602739	.005260273
25	.004109589	.004794520	.005479452
26	.004273972	.004986301	.005698630
27	.004438356	.005178082	.005917808
28	.004602739	.005369863	.006136986
29	.004767123	.005561643	.006356164
30	.004931506	.005753424	.006575342
31	.005095890	.005945205	.006794520
32	.005260273	.006136986	.007013698
33	.005424657	.006328767	.007232876
34	.005589041	.006520547	.007452054
35	.005753424	.006712328	.007671233
36	.005917808	.006904109	.007890411

Days	At 3 per *Cent.*	At 4 per *Cent.*	At 5 per *Cent.*
37	.003041096	.004054794	.005068493
38	.003123287	.004164383	.005205479
39	.003205479	.004273972	.005342466
40	.003287671	.004383561	.005479452
41	.003369863	.004493150	.005616438
42	.003452054	.004602739	.005753424
43	.003534246	.004712328	.005890411
44	.003616438	.004821917	.006027397
45	.003698629	.004931506	.006164383
46	.003780821	.005041095	.006301369
47	.003863013	.005150684	.006438356
48	.003945205	.005260273	.006575342
49	.004027396	.005369862	.006712329
50	.004109589	.005479452	.006849315
51	.004191780	.005589041	.006986301
52	.004273972	.005698629	.007123287
53	.004356164	.005808218	.007260274
54	.004438355	.005917807	.007397260
55	.004520547	.006027396	.007534246
56	.004602739	.006136985	.007671233
57	.004684931	.006246575	.007808219
58	.004767123	006356164	.007945205
59	.004849314	.006465753	.008082192
60	.004931506	.006575342	.008219178
61	.005013698	.006684931	.008356164
62	.005095890	.006794520	.008493150
63	.005178082	.006904109	.008630137
64	.005260273	.007013698	.008767123
65	.005342465	007123287	.008904109
66	.005424657	.007232876	.009041096
67	.005506849	.007342465	.009178082
68	.005589041	.007452054	.009315068
69	.005671232	.007561643	.009452055
70	.005753424	.007671232	.009589041
71	.005835616	.007780821	.009726027

At

Days	At 6 per *Cent.*	At 7 per *Cent.*	At 8 per *Cent.*
37	.006082191	.007095890	.008109589
38	.006246575	.007287671	.008328767
39	.006410959	.007479452	.008547945
40	.006575342	.007671233	.008767123
41	.006739726	.007863013	.008986301
42	.006904109	.008054794	.009205479
43	.007068493	.008246575	.009424657
44	.007232876	.008438356	.009643835
45	.007397259	.008630137	.009863013
46	.007561643	.008821917	.010082191
47	.007726027	.009013698	.010301369
48	.007890410	.009205479	.010520547
49	.008054794	.009397260	.010739725
50	.008219178	.009589041	.010958904
51	.008383561	.009780821	.011178082
52	.008547945	.009972602	.011397259
53	.008712328	.010164383	.011616438
54	.008876712	.010356164	.011835616
55	.009041095	.010547945	.012054794
56	.009205479	.010739725	.012273972
57	.009369863	.010931506	.012493150
58	.009534246	.011123287	.012712328
59	.009698629	.011315068	.012931506
60	.009863013	.011506849	.013150684
61	.010027397	.011698629	.013369862
62	.010191780	.011890410	.013589040
63	.010356164	.012082191	.013808219
64	.010520547	.012273972	.014027397
65	.010684931	.012465753	.014246575
66	.010849314	.012657533	.014465753
67	.011013698	.012849314	.014684931
68	.011178081	.013041095	.014904109
69	.011342465	.013232876	.015123287
70	.011506849	.013424657	.015342465
71	.011671232	.013616437	.015561643

Days	At 3 per *Cent.*	At 4 per *Cent.*	At 5 per *Cent.*
72	.005917808	.007890410	.009863013
73	.006000000	.008000000	.010000000
74	.006082192	.008109589	.010136986
75	.006164383	.008219178	.010273972
76	.006246575	.008328767	.010410959
77	.006328767	.008438356	.010547945
78	.006410959	.008547945	.010684931
79	.006493150	.008657534	.010821918
80	.006575342	.008767123	.010958904
81	.006657534	.008876712	.011095890
82	.006739726	.008986301	.011232877
83	.006821918	.009095890	.011369863
84	.006904109	.009205479	.011506849
85	.006986301	.009315068	.011643835
86	.007068493	.009424657	.011780822
87	.007150685	.009534246	.011917808
88	.007232876	.009643835	.012054794
89	.007315068	.009753424	.012191781
90	.007397260	.009863013	.012328767
91	.007479452	.009972602	.012465753
---	--Months.--	--Months.--	---Months.--
3	.007500000	.010000000	.012500000
6	.015000000	.020000000	.025000000
9	.022500000	.030000000	.037500000
---	-Years, &c.-	-Years, &c.-	-Years, &c.-
1	.030000000	.040000000	.050000000
1¼	.037500000	.050000000	.062500000
1½	.045000000	.060000000	.075000000
1¾	.052500000	.070000000	.087500000
2	.060000000	.080000000	.100000000
2¼	.067500000	.090000000	.112500000
2½	.075000000	.100000000	.125000000
2¾	.082500000	.110000000	.137500000
3	.090000000	.120000000	.150000000
3¼	.097500000	.130000000	.162500000
3½	.105000000	.140000000	.175000000
3¾	.112500000	.150000000	.187500000

A

Days	At 6 per *Cent.*	At 7 per *Cent.*	At 8 per *Cent.*
72	.011835616	.013808218	.015780821
73	.012000000	.014000000	.016000000
74	.012164383	.014191781	.016219178
75	.012328767	.014383561	.016438356
76	.012493150	.014575342	.016657534
77	.012657534	.014767123	.016876712
78	.012821918	.014958904	.017095890
79	.012986301	.015150685	.017315068
80	.013150685	.015342466	.017534246
81	.013315068	.015534246	.017753424
82	.013479452	.015726027	.017972603
83	.013643835	.015917808	.018191781
84	.013808219	.016109589	.018410959
85	.013972602	.016301370	.018630137
86	.014136986	.016493150	.018849315
87	.014301370	.016684931	.019068493
88	.014465753	.016876712	.019287671
89	.014630137	.017068493	.019506849
90	.014794520	.017260274	.019726027
91	.014958904	.017452054	.019945205
---	---Months---	---Months---	---Months---
3	.015000000	.017500000	.020000000
6	.030000000	.035000000	.040000000
9	.045000000	.052500000	.060000000
---	-Years, &c.-	-Years, &c.-	-Years, &c.-
1	.060000000	.070000000	.080000000
1¼	.075000000	.087500000	.100000000
1½	.090000000	.105000000	.120000000
1¾	.105000000	.122500000	.140000000
2	.120000000	.140000000	.160000000
2¼	.135000000	.157500000	.180000000
2½	.150000000	.175000000	.200000000
2¾	.165000000	.192500000	.220000000
3	.180000000	.210000000	.240000000
3¼	.195000000	.227500000	.250000000
3½	.210000000	.245000000	.280000000
3¾	.225000000	.262500000	.300000000

Explication.

The firſt Colume of every Page being Time, againſt it ſtands the Intereſt of one Pound for that Time, at the Rate per *Cent.* ſpecify'd on the top of the Colume, and at the ſame View you have the Intereſt of 10 *l.* 100 *l.* 1000 *l.* and all in one Line. Let the Intereſt of 500 *l.* for 73 Days at 5 per *Cent.* be required; turn to 73 Days and againſt that, under the Rate of 5 per *Cent.* it appears that

l.				*l.*	*s.*	*d.*	
1	is	.010	or			2	$\frac{1}{4}$
10	is	.10	or		02	0	0
100	is	1 .00	or	1	00	0	0
		1 .11	or	1	02	2	$\frac{1}{4}$

So that one Pound after that manner being found to be the Intereſt of an 100 *l.* at 5 per *Cent.* per *Annum* for 73 Days, 5 times that Mony, to wit, 5 *l.* muſt therefore be the Intereſt of 500 *l.* Again, If the Intereſt of 111 *l.* at the ſame Rate and for the ſame Time were required; take the total of the above Sums for an Anſwer, being 1 .11 or 1 *l.* 2 *s.* 2 *d.* $\frac{1}{4}$. So that let the given Principle be what number of Pounds, Shillings and Pence it will, it may be either taken in Parts as 111. *l* above, or by multiplying the Intereſt of one Pound by the given Principle; ſo that no Sum at any of the inſerted Rates or Time can require more than one Multiplication. And altho' this is Calculated only for 3 Years $\frac{3}{4}$. beyond which Time ſimple Intereſt ſeldom happens, yet the Intereſt from thence for any longer Time may as eaſily and readily be found, *viz.*

Ex. 1. Unto what amounts the Intereſt of 79 *l.* 15 *s.* for 6 Years ½ and 73 Days, at 6 per *Cent* per *Annum.*

Years.
.180 ⎫ the Intereſt of ⎧ 3
.210 ⎬ one Pound for ⎨ 3 ½
.012 ⎭ 73 Days.
.402 the Intereſt of one Pound at the given Rate and Time, which multiply'd by the Principle, produces the Anſwer.

79.75
.402
15950
31900
32,05950 *Anſ.* 32 *l.* 01 *s.* 2 *d.* +

Likewiſe may the Intereſt of any Sum at any other Rate per *Cent.* whatſoever, be readily found according to the Directions in the laſt Head in the Rule of Intereſt for Days.

Ex. 2. What comes the Intereſt of 432 *l.* 10 *s.* for 1 Year ¼. at 4 *l.* ½. per *Cent.* per *Annum.*

Firſt take the Intereſt of one Pound for the given Time at 5 per *Cent.* per *Annum*, which is .0625
From which Subtracting one Tenth 625
gives the Intereſt of one Pound at 4 ½ per *Cent.* .5625
Multiply'd by the given Principle 432,5
28125
11250
16875
22500
Anſwer 24 *l.* 06 *s.* 05 *d.* ¼ 24,328125

SECTION VIII.

EXCHANGE.

1. IN *London*, and throughout *England*, Accompts are kept in Pounds, Shillings and Pence Sterling, and are caſt up as *Flemiſh* and *French* Money are, by 12 and 20, and it Exchanges with all places by Pence, *i. e.* giving ſo many pence Sterling for the Pieces on which the Exchange is made, except

1. *Portugal*, and with it in Shillings and Pence, on their *Milrea*.
2. *Antwerp*, *Hamborough*, &c. Countries bordering upon *Flanders* and *Holland*, and with them by the Pound of 20 *s.* Sterling.
3. *Ireland*, and with it by the hundred Pound.

II. The principle Places where the courſe of Exchange is ſetled, and in what Denomination they Exchange with *London*, you may find in the 3d Chapter of Mr. *Kerſy*'s Appendix; but the places to which there is a more conſtant courſe of Exchange from *London*, are

In *Italy*,

Genoa, *Legorn*, *Venice*, &c.

Genoa and *Legorn* exchange with *London* by the Dollar, or piece of Eight, *Venice* by the Ducat.

Genoa and *Legorn* keep their Accompts in Livres, Solz and Deniers, 5 Livres a piece of Eight at *Genoa*, and 6 at *Legorn*.

The Par of the Dollar, or piece of Eight, with *London* is 54 *d.* Sterling.

In *Spain.*

Madrid, Cales, Bilbo, &c.
Exchange alſo by the piece of Eight.
The Par being the ſame as *Genoa* and *Legorn.*

Portugal.

Lisbon and *Oporto,* &c.
Exchange on the *Milrea.*
The Par about 6 *s.* 8 *d.* ½ Sterling.

In *France.*

Paris, Lyons and *Roven*
Exchange by the Crown.
But keep their Accompts in Livres, Solz and Deniers,

12 Denier	makes	1 Sol.
20 Solz		1 Livre
3 Livres		1 Crown

The Par of the *French* Crown with *London* is 4 *s.* 6 *d.* Sterling.

In the *Netherlands.*

Antwerp, Bruſſels, Amſterdam, Rotterdam, Hamborough, &c.
Exchange with *London* by Shillings and Pence *Flemiſh.* Accompts are in theſe places kept in *Flemiſh* Pounds, Shillings and Pence, by ſome in Guilders, Stivers and Pennicks. The *Flemiſh* Pounds, Shillings and Pence, are divided as our Money, *viz.* one Pound into 20 *s.* and one Shilling into 12 *d.* But,

16 Pennicks,	one	Stivre	The par with *Lon.*	0 *s.* 1 *d.* $\frac{1}{5}$
6 Stivres,		Flemiſh Shilling		0 : 7 : $\frac{1}{5}$
20 Stivres,		Guilder		2 : 0 : 0
6 Guilders,		Flemiſh Pound		12 : 0 : 0
30 Stivres,		Common Dol,		3 : 0 : 0
50 Stivres,		Specie Dollar		5 : 0 : 0
63 Stivres,		Duccatoon		6 : 3 : $\frac{2}{5}$

The Par between the *Flemiſh* Pound and Pound Sterling, is 33 *s.* 4 *d.* But the courſe of Exchange may, and doth vary ſome Shillings.

In the Arithmetical part, or caſting up of Exchanges, I ſhall only inſtance in thoſe with *France* and *Flanders*; for as *London* exchanges with *France* by the Pence, ſo it doth with all places in *Spain* and *Italy*; and as it exchanges with *Flanders*, at *Bruſſels* and *Antwerp*, ſo it doth with *Germany* and *Holland*, at *Hamborough* and *Amſterdam*, &c. by the Pound Sterling, for their *Flemiſh* Shillings and Pence.

Of French *Exchange.*

The moſt common ways of caſting theſe up are two, *viz.*

Rules, *Either multiply the given Crowns by the Pence of Exchange, taking parts for the Farthings*; *(if any be)* otherwiſe,

Take the aliquot Parts for the Pence of the Exchange out of a Pound, dividing the given Crowns by thoſe Parts, and the Total of which is the Anſwer.

Ex. How much Sterling muſt I pay here to receive in *France* 479 Crowns, Exchange at 52 *d.* Sterling per Crown.

4 *s.*	$\frac{1}{5}$	479 Crowns
4	$\frac{1}{12}$	95 : 16
		7 : 19 : 8
		103 : 15 : 8

479 Crowns
52
958
2395
12) 24908
2\|0) 207\|5 : 8
103\|15 : 8

So that all Exchanges made with *London*, upon the Ducat, Dollar, Duccatoon, Florin, Milrea, *&c.* are wrought after the ſame manner, and are no more than ſo many ſeveral Queſtions in the Rule of Three, or of Practice ; for if the above Crowns were Ells, *&c.* you muſt pay as much for them at ſuch a price, as Sterling, for ſo many Crowns at ſuch Exchange. If the Exchange is at ſo many Pence and a Fraction of a Penny, ſee my ſecond Example of the 13th Rule in Practice.

Of Flemiſh *Exchange.*

Firſt, To reduce Sterling into *Flemiſh*.

Rule, *Take the Aliquot parts for what the Rate of the Exchange is above a Pound, dividing the given Sterling by the ſaid Parts, the total of which, with the given Sterling, is the* Flemiſh *Money.*

Ex. 1. One in *Antwerp* delivering Money by Exchange for *London* at 35 *s.* 6 *d. Flemiſh* per pound Sterling, how much muſt he pay there to receive here 597 *l.* Sterling?

10 *s.*	$\frac{1}{2}$	597
5	$\frac{1}{2}$	298 : 10
6	$\frac{1}{10}$	149 : 05
		14 : 18 : 6
	Anſwer l.	1059 : 13 : 6 *Flemiſh.*

Ex. 2. A Merchant at *Rotterdam* hath a Bill drawn on him for the value of 673 *l.* 16 *s.* 8 *d.* Sterling, Exchange at 33 *s.* 4 *d.* *Flemiſh* per pound Sterling, how much muſt he pay there?

10 s.	$\frac{1}{2}$	673 : 16 : 8
3 : 4	$\frac{1}{6}$	336 : 18 : 4
		112 : 06 : 1 $\frac{1}{4}$
Anſwer 1123 *l.* 1 *s.* 1 *d.* $\frac{1}{3}$ *Flemiſh*		1123 : 01 : 1 $\frac{1}{4}$

Secondly, To reduce Sterling into *Guilders.*

Rule, *Reduce the Sterling into* Flemiſh, *by the foregoing Rule, which* Flemiſh *Money multiply by* 6.

Ex. If 397 *l.* 15 *s.* Sterling, Exchange at 34 *s.* 8 *d.* *Flemiſh* per pound Sterling, were remitted to *Amſterdam*, how many Guilders, Stivers, &c. may be received at the ſaid Rate of Exchange?

	10 s.	$\frac{1}{2}$	397 : 15
	4	$\frac{1}{5}$	198 : 17 : 06
	8	$\frac{1}{6}$	79 : 11 : 00
			13 : 05 : 02
			689 : 08 : 08
Guilders.	*Stivers*		6
Anſw. 4136 :	12		4136 : 12 : 00

Thirdly, To Reduce *Flemiſh* into Sterling.

Rule, *Reduce the given* Flemiſh *Sum, either into Pence, two Pences, three Pences, Groats or Six-pences,* &c. *which make a Dividend, then Reduce the Rate of* Exchange *into the ſame, which take for a Diviſor, and the Quote of that Diviſion will be the Anſwer in Pounds Sterling, the Remainder being valued according to Mr.* Wingate's *5th Rule of the 7th Chapter.*

Ex. 1. A Merchant at *Bruſſels* delivers 579 *l.* 10 *s.* *Flemiſh*, Exchange at 35 *s.* 6 *d.* to receive at *London* one Pound Sterling, How much Sterling muſt he receive?

35 : 6 579 *l.* 10 *s.*
2 40
71 71) 23180 (326 : 09 : 06 †
88
460
Anſw. 326 : 09 : 06 34

Ex. 2. How much Sterling muſt be here paid to receive in *Hamborough* 836 *l.* 8 *s.* 4 *d.* *Flemiſh*, Exchange at 36 *s.* 8 *d.* *Flemiſh* per pound Sterling?

36 : 8 836 *l.* 8 *s.* 4 *d.*
3 60
11|0 5018|5 Groats
Anſwer 456 *l.* 4 *s.* 6 *d.* 1 $\frac{[illegible]}{1}$

Fourthly, To reduce Guilders, Stivers and Pennicks into Sterling.

Rule, *Reduce the ſaid Guilders, Stivers and Pennicks into Pence, two Pences, or Groats, which divide by the given Rate of Exchange (when reduced into the ſame Name) and the Quotient will be Pounds Sterling.*

Note, That 40 *Flemiſh* Pence makes one Guilder.

Ex. 1. A Merchant at *Rotterdam* delivering 1489 *Guild.* 10 *Stiv.* 8 *Pen.* for *London*, Exchange at 35 *s.* 7 *d.* I demand how much Sterling he may draw the Bill for?

1489 : 10 : 8
40
427) 59581 (139 : 10 : 8 +
1688
4071
228 *Anſw.* 139 : 10 : 8

Guilders. Stiv.
Ex. 2. How much Sterling is equivalent to 4372 : 10, Exchange 36 *s.* 4 *d.* *Flemiſh* per pound Sterling?

3
109 Groats 109) 43725 (401 : 2 : 11 +
125
16

Or more ſhort.

When the Pence of the Exchange is ſo many, as that 40 (which is the *Flemiſh* Pence in one Guilder) is an aliquot part thereof.

Rule, *Divide the Guilders, Stivres,* &c. *by that Aliquot part, and the Quote is the Sterling Money in Pounds, Shillings and Pence; but to reduce Sterling into Guilders, Stivers,* &c. *multiply by the aforeſaid part.*

Ex. 1. How much Sterling, at 36 *s.* 8 *d.* *Flemiſh* per pound Sterling muſt be paid for 1587 : 10 *Stivers?*

11) 1587 : 10
144 : 06 : 04 1 $\frac{4}{1}$ *Anſw.* 144 : 06 : 04 1 $\frac{4}{1}$.

Ex. 2. Exchange at 33 *s.* 4 *d. Flemiſh* per pound Sterling, how many Guilders, Stivers, *&c.* may I draw for 179 *l.* 17 *s.* 10 *d.* Sterling?

179 : 17 : 10
Guild. Stiv. Penn. 10
Anſw. 1798 : 18 : 04 1798 : 18 : 05 +

Here note, That when the Rate of Exchange is at Par, then the Proportion between their Money and ours, is the ſame as between their Cloth Meaſure and ours; for as five of their Ells make three of ours, ſo, at ſuch Exchange, five of their Flemiſh *Pounds make three Pound Sterling, or* 500 *l.* Flemiſh 300 *l. Sterling.*

Of Gain or Loſs by Exchange.

Note, That when Exchange is made with *Italy*, *France*, *Spain*, &c. with which, as aforeſaid, *London* Exchanges for ſo many Pence for their Pieces. The Gain thereby is ſo much the more, by how much the Courſe of Exchange runs low; becauſe it is evident That I can receive more Ducats for 500 *l.* when the Exchange is at 53 *d.* than at 56 per Piece.

But when I Remit for *Flanders*, or *Holland*, with which *London* Exchanges on the pound Sterling, the higher the Exchange is above Par, the more we have the advantage of Gain; for according to reaſon, I may receive more *Flemiſh* Money for 1000 *l.* Sterling, when the Exchange is at 36 *s.* 8 *d.* than at 35 *s.* 4 *d.*; for at 35 *s.* 4 *d.* I gain but 2 *s.* per Pound, but at 36 *s.* 8 *d. Flemiſh*, I gain by Exchange 3 *s.* 4 *d.* for every pound Sterling I remit.

Decimal Tables

FOR

FLEMISH Exchanges,

At the moſt uſual

Rates of Exchange.

A DECIMAL TABLE

Rate of Exchan.		The Value of Pounds Sterling in Pounds *Flemiſh*.	The Value of Pounds *Flemiſh* in Pounds Sterling.
s.	d.		
32	---	1 .6	.625....
	1	1 .6041666	.6233766
	2	1 .6083333	.6217616
	3	1 .6125...	.6201550
	4	1 .6166666	.6185567
	5	1 .6208333	.6169665
	6	1 .625....	.6153846
	7	1 .6291666	.6138107
	8	1 .6333333	.6122449
	9	1 .6375...	.6106870
	10	1 .6416666	.6091370
s.	11	1 .6458333	.6075949
33	---	1 .65.....	.6060606
	1	1 .6541666	.6045343
	2	1 .6583333	.6030150
	3	1 .6625...	.6015037
	4	1 .6666666	.6......
	5	1 .6708333	.5985037
	6	1 .675....	.5970149
	7	1 .6791666	.5955335
	8	1 .6833333	.5940594
	9	1 .6875...	.5925925
	10	1 .6916666	.5911330
	11	1 .6958333	.5896806
34	----	1 .7......	.5882352
	1	1 .7041666	.5867970
	2	1 .7083333	.5853658
	3	1 .7125...	.5839416
	4	1 .7166666	.5825242

Rat

Rate of Excha.	The Value of Pounds Sterling in Guilders.	The Value of Guilders in Pounds Sterling.
s.		
32	9.6	.10416666
1	9.625	.10389610
2	9.65	.10362694
3	9.675	.10335916
4	9.7	.10309278
5	9.725	.10282775
6	9.75	.10256410
7	9.775	.10230178
8	9.8	.10204081
9	9.825	.10178116
10	9.85	.10152283
11	9.875	.10126581
33	9.9	.10101010
1	9.925	.10075571
2	9.95	.10050250
3	9.975	.10025061
4	10.0	.1
5	10.025	.09975061
6	10.05	.09950248
7	10.075	.09925558
8	10.1	.09900990
9	10.125	.09876541
10	10.150	.09852216
11	10.175	.09828011
34	10.2	.09803921
1	10.225	.09779951
2	10.25	.09756096
3	10.275	.09732360
4	10.3	.09708736

Rate of Exchan.		The Value of Pounds Sterling in Pounds *Flemiſh*.	The Value of Pounds *Flemiſh* in Pounds Sterling.
s.	d.		
34	5	1 .7208333	.5811138
	6	1 .725....	.5797101
	7	1 .7291666	.5783132
	8	1 .7333333	.5769230
	9	1 .7375...	.5755395
	10	1 .7416666	.5741621
	11	1 .7458333	.5727923
35----		1 .75.....	.5714285
	1	1 .7541666	.5700712
	2	1 .7583333	.5687209
	3	1 .7625...	.5673758
	4	1 .7666666	.5660377
	5	1 .7708333	.5647058
	6	1 .775....	.5633802
	7	1 .7791666	.5620609
	8	1 .7833333	.5607476
	9	1 .7875...	.5594405
	10	1 .7916666	.5581395
	11	1 .7958333	.5568445
36----		1 .8......	.5555555
	1	1 .8041666	.5542725
	2	1 .8083333	.5529954
	3	1 .8125...	.5517241
	4	1 .8166666	.5504587
	5	1 .8208333	.5491990
	6	1 .825....	.5479452
	7	1 .8291666	.5466970
	8	1 .8333333	.5454545
	9	1 .8375...	.5442176
	10	1 .8416666	.5429864
	11	1 .8458333	.5417607

Rate of Exchan.	The value of pounds Sterling in Guilders.	The value of Guilders in pounds Sterl.
s. *d.*		
34 5	10.325....	.09685231
6	10.35.....	.09661835
7	10.375....	.09638553
8	10.4......	.09615383
9	10.425....	.09592325
10	10.45.....	.09569368
11	10.475....	.09546538
35----	10.5......	.09523808
1	10.525....	.09501186
2	10.55.....	.09478681
3	10.575....	.09456263
4	10.6......	.09433961
5	10.625....	.09411763
6	10.65.....	.09389671
7	10.675....	.09367681
8	10.7......	.09345793
9	10.725....	.09324008
10	10.75.....	.09302325
11	10.775....	.09280741
36----	10.8......	.09259259
1	10.825....	.09237875
2	10.85.....	.09216590
3	10.875....	.09195402
4	10.9......	.09174311
5	10.925....	.09153316
6	10.95.....	.09132420
7	10.975....	.09111616
8	11.0......	.09090909
9	11.025....	.09070293
10	11.05.....	.09049773
11	11.075....	.09029345

Rate of Exch.	The value of Pounds Sterl. in Pounds *Fl.*	The value of Pounds *Fle.* in Pounds ſter.
s. d.		
37---	1 .85	.5405405
1	1 .8541666	.5393258
2	1 .8583333	.5381166
3	1 .8625 . . .	.5369127
4	1 .8666666	.5357143
5	1 .8708333	.5345211
6	1 .875	.5333333
7	1 .8791666	.5321508
8	1 .8833333	.5309734
9	1 .8875 . . .	.5298013
10	1 .8916666	.5286343
11	1 .8958333	.5274725
38---	1 .9	.5263157
1	1 .9041666	.5251641
2	1 .9083333	.5240174
3	1 .9125 . . .	.5228758
4	1 .9166666	.5217391

Rate

Rate of Exch.	The value of Pounds Sterl. in Guilders.	The value of Guilders in Pounds Sterl.
s. d.		
37---	11.1......	.09009009
1	11.125....	.08988763
2	11.15.....	.08968610
3	11.175....	.08948545
4	11.2......	.08928571
5	11.225....	.08908685
6	11.25.....	.08888888
7	11.275....	.08869180
8	11.3......	.08849556
9	11.325....	.08830021
10	11.35.....	.08810571
11	11.375....	.08791208
38---	11.4......	.08771928
1	11.425....	.08752735
2	11.45.....	.08733624
3	11.475....	.08714596
4	11.5......	.08695651

The Explication.

The firſt Column ſhews the Rate of the Exchange, from 32 *s.* to 38 *s.* 4 *d.* and is extended no farther; for a regulation being made of the Coin, it's not probable it will riſe higher, or fall lower than thoſe Limitations.

2. The ſecond Column on the Left-hand, ſhews the value of Pounds Sterling, expreſt in Pounds *Flemiſh*, whereby, at the firſt view, you diſcern what is the amount of 1, 10, 100, 1000, *&c.* to ten Millions of Sterling, in Pounds *Flemiſh*, at any of the inſerted Rates of Exchange: For Example, If the Exchange is at 35 *s.* 2 *d.* One Pound Sterling being Equivalent to one Pound fifteen Shillings and two Pence at one view; therefore it appears, that

l.	*l.*	*s.*	*d.*	
10 *l.* Sterling makes	17	11	8	*Flem.*
100	175	16	8	
1000	1758	6	8 &c.	

all which *Flemiſh* Moneys are found in the Line againſt the ſaid given Rate of Exchange: But if the Sterling Money, for which the value in pounds *Flemiſh* is required, conſiſts in odd Pounds, or have Shillings, *&c.* annexed; then multiply the *Flemiſh* Money, at the given Exchange, by the *Sum* in Sterling, and the Product anſwers the Queſtion. Thus 67 *l.* 10 *s.* Sterling, multiply'd by 1, 75833, the *Flemiſh* Money at 35 : 2, makes 118 *l.* 13 *s.* 9 *d.* *Flemiſh.*

3. The Third Column on the Left Page ſhews the value, or amount of pounds *Flemiſh* in Sterling, after the ſame manner as the ſecond ſhews the value or amount of pounds Sterling in *Flemiſh* Money:

For Inſtance.

At the aforeſaid Exchange of 35 *s.* 2 *d.* I ſee that one Pound
Flemiſh being Equivalent to .5687209, or *l.* 11*s.* 4*d.* ½ Ster-
Therefore 10 *Flem.* makes 5.687209, or 5 : 13 : 9 : ling
100 *l.* 56.87209, or 56 : 17 : 6 : very
and 1000 568.7209, or 568 : 14 : 5 : near

But if your *Flemiſh* Money, the value of which is required in Sterling, conſiſts of odd Pounds, or have Shillings, *&c.* annex'd, Multiply the ſame by the Sterling, as before directed, and the Product of ſuch Multiplication anſwers the Queſtion; ſo the Exchange being at 35 *s.* 2 *d.* 137 *l.* 10 *s.* *Flemiſh*, multiplied by .5687209 Sterling, produces 78 *l.* .19912375, or 78 *l.* 4 *s.* + Sterling.

4. The Fifth Column, or the ſecond on the Right-hand Page, ſhews the value or amount of Pounds Sterling in Guilders, or what number of Guilders I ought to receive, or pay, for any Sum of Sterling, at any of the inſerted Rates of Exchange;

For Example.

At the aforeſaid Exchange of 35 *s.* 2 *d.* it appears, that 10 Guilders, 10 Stivers, and 8 Pennicks, are equivalent to one pound Sterling, as likewiſe, that

Guild.	*Stivers*		*l.*	
105	: 10	are to	10	Sterling.
1055		to	100	
10550		to	1000	

If the Sterling Money is in odd Pounds, or have Shillings, &c. annexed, you are to multiply the Guilders correſponding to the Rate of Exchange by your Sterling Sum, as above directed; ſo 132 *l.* 10 *s.* Sterling, Exchange at 35 *s.* 2 *d.* will be found to make 1397 Guilders, 17 Stivers, 8 Pennicks.

V. The laſt Column ſhews the value of Guilders in pounds Sterling, or how much Sterling I ought to receive, or pay, for any number of Guilders, at any of the ſpecify'd Rates of Exchange.

So at the aforeſaid Rate of Exchange, of 35 *s.* 2 *d.* one Guilder being worth .09478681, or 1 *s.* 10 *d.* ¼ Sterling;

Therefore

		l.	*s.*	*d.*	
10 make	.9478681, or		18	11 : ½	Sterling very near.
100	9.478681, or	9	09	6 : ¾	
1000	94.78681, or	94	15	8 : ¾	

If your Guilders be in odd Numbers, or have Stivers annex'd, multiply them alſo by the Value in Sterling correſponding to the Rate of Exchange, and the Product will anſwer your Demand.

Guild. Stiv. Penn.
So 1397 : 17 : 8, Exchange at 35 *s.* 2 *d.* being multiply'd by .09478681, amounts to 132 *l.* 10 *s.* Sterling, as aboveſaid.

☞ *Note*, The reaſon of the Tables being Calculated ſo high, is, That you may be more exact in long Sums, altho' you'l ſeldom have occaſion to make uſe of more than five or ſix Figures. And the Dots plac'd to the Right-hand are incerted for the more eaſie, and ſafe, Obſervation of large even Sums. But if the caſting up of odd Sums ſhould be thought as troubleſome to be effected by theſe Tables, as by the common Method of doing them (though in all Caſes perform'd by one ſingle Multiplication) yet Even Sums moſt commonly happening in Drawing and Remitting by Exchange, the uſefulneſs of theſe Tables will be ſufficiently Manifeſt.

SECTION X.

Of MEASURING.

THE various kinds of which, are Three, *viz.*

I. *Lineal*, call'd (by Work-men) *Running Meaſure*, in that it reſpects *Length* only, the parts of which are

12 *Inches*,	1 *Foot*,
3 *Feet*,	1 *Yard*,
16½ *Feet*,	1 *Rodd.*

II. *Superficial*, or *Square Meaſure*, in that it reſpects *Length* and *Breadth*; the parts are, *viz.*

144 *Inches*,	1 *Foot*,
72 *Inches*,	½ *a Foot*,
36 *Inches*,	¼ *of a Foot*,
48 *Inches*,	⅓ *of a Foot*,
18 *Inches*,	½ *a-quarter of a Foot*,
272 ¼ *Feet*,	1 *Rodd*,
136 *Feet*,	½ *a Rodd*,
Inches 1296, or 9 *Feet*,	1 *Yard.*

Note, That in dividing by 272¼, which reduces *Superficial Feet* into *Square Rodds*, Work-men take no notice of the ¼ of the Foot, dividing only by 272, which gives the Content a little too much.

III. *Solid*, or *Cube Meaſure*, reſpecting *Length*, *Breadth* and *Thickneſs*; whoſe parts are,

Inches.		
1728,	}	1 *Foot*,
1296,	}	¾ *of a Foot*,
864,	}	½ *of a Foot*,
432, &c.	}	¼ *of a Foot*.
27 *Feet*,	}	1 *Yard*.

Of Superficial Meaſure.



A General Rule.

Multiply the Length by the Breadth, and the Product is the Superficial Content.

Demonſtration.

For if a *Line*, A B, being divided into any number of *equal parts*, be moved conſtantly at *right Angles* with another Line, B C, divided into any other number of ſuch *equal Parts*, as are in, A B, it will, when it is advanc'd vanc'd to one part of, B C, deſcribe, thereby as many *equal Squares* as there are *Parts* in it; and if mov'd on further, to the next *Part*, it will deſcribe as many more *Squares*, and ſo the ſame number of *Squares* will be as often deſcribed as there are *Parts* in the Line, B C, and conſequently there muſt be as many *Squares* in the *Surface*, made by theſe *Lines*, as there are *Units* in the *Product* of the *Parts* of the two *Sides*.



Now the *Square* of any thing is found three ſeveral ways, viz.

1. *By Decimals,*
2. *By aliquot Parts,*
3. *By croſs Multiplication.*

All which I ſhall promiſcuouſly uſe in the following Examples. But the latter of the three is us'd by moſt Workmen, except *Glaziers*, whoſe Foot being *Decimally* divided, caſt up the Content of their Work by *Decimals*.

How to Multiply by the two laſt, ſee my 4th Head in Multiplication.

Note, *The Content of Work, is given either by the Foot, the Yard, the Square, or the Rodd,* viz.

I. By the Foot.

Glazing and *Maſoning.*

Dimenſions are taken in *Feet* and *Inches*, the *Content* given *per* Foot Square.

In Glazing.

Example, How many Foot of *Glazing* is there in that *Pane of Glaſs* which is 4 *Foot* 5. and 2 *Foot* 42.

```
   2,42
    4,5
  -----
   1210
   968
  -----
  10,890    Anſw. 10 f. 890
```

In Maſoning.

Example, Suppoſe a *Yard pav'd* with *Free-ſtone*, viz. *Length* 22 *Feet* 4 *Inch. Breadth*, 19 *Feet* 7 *Inch.* how many *Square Feet?*

```
   22 . 4
   19 . 7
  424 . 4
    7 . 5 . 4
    5 . 7
  437 . 4 . 4        Feet In. Parts.
               Facit. 437 . 4 . 4
```

But if only the *Breadth* was given, and it were required how much in *Length* will make a *foot Square*, ſee Mr. *Kerſey*'s 51 Queſtion of his 10th Chapter.

II. By the Yard.

Painting, Joynery, Plaiſtering, &c.

Dimenſions are taken in *Feet* and *Inches*; but the Content given in *Square Yards*, found by dividing the *Superficial Feet* by 9.

In Painting.

Example, The *Height* of a *Room* being 12 . 4 (*Feet. Inches*), and 84 *feet* 11 *Incehs* about, how many *Square Yards?*

```
    84 . 11 about
    12 .  4 high
  1019 .  0
    28 .  3 . 8
 9)1047 .  3 . 8              Yd. feet. inch. part.
   116 . 3 . 3 . 8   Anſw. 116 . 3 . 3 . 8
```

Joynery.

Example. How many *Yards* of *Wainſcot*, doth that *Room* take, up whoſe *Heigth* is 12 *feet* 3 *Inches*, and *Compaſs* 104 *feet* 6 *Inches?*

feet. *inches.*
104 . 6
12 . 3
1248 . 0
06
26 . 1
9) 1280 . 1 . 6
142 . 2 . 1 , 6

yar. *feet.* *inch.* *part.*
Anſ. 142 . 2 . 1 . 6

Plaiſtering.

Example. How many *Yards* in a *Cieling* 47 *feet*, 4 *Inches*, 7 *Parts* Long, and 18 *feet* Broad?

47 . 4 . 7 long
6 broad
284 . 3 . 6
3
9) 852 . 10 . 6
94 . 6 . 10 . 6

Yar. *feet*, *inch.* *parts.*
Facit. 94 . 6 . 10 . 6

Under which head you may further note, that if you multiply the $\frac{1}{3}$ part of the *length*, by the $\frac{1}{3}$ of the *breadth*, the

Product will be the Content in *Square Yards*, it being the ſame to take $\frac{1}{3}$ of *Lineal* as $\frac{1}{9}$ of *Superficial Meaſure.*

III. By the Square.

Partitioning, Flooring, Roofing, Tyling.

Dimenſions are taken in *Feet* and *Inches*, but the Content or Value is by the *Square* of 10 *Feet*, done by cutting off two *Figures* of the *Superficial Feet* towards the Right-hand.

Partitioning.

Example. How many *Squares* are contained in that Work that is 199 *feet* 10 *Inches*, in *Length*, and 10 *feet* 7 *inche* *high* ?

```
  199 . 10
   10 .  7
 ---------
 1998 .  4
   66 .  7 . 4
   49 . 11 . 6
 -------------
21,14 . 10 . 10     Square feet, Inch. parts.
           Anſwer  21 . 14 . 10 . 10
```

From which ſometimes the Content of *Doors* and othe *Vacancies* are deducted,

Floor

Flooring.

Example. A *Floor* being 49 *feet*, 7 *inch.* 4 *part.* long, 26 *feet*, 6 *inch.* broad, how many *Squares?*

49 . 7 . 4
26 . 6 . 0
1289 . 10 . 8
24 . 9 . 8
Anſw. 13,14 . 8 . 4

Roofing.

Example. Suppoſe an Houſe 18 *feet* 4 *inches* in *Front*, and 37 *feet* 10 *inches* in *Depth*, how many *Square* in the Roof?

37 . 10
3
113 . 6
6
$\frac{1}{3}$) 681 . 0
12 . 7 . 4
$\frac{1}{2}$) 698 . 7 . 4
346 . 9 . 8 added
10,40 . 5 . 0

Square feet, inch.
Anſw. 10 . 40 . 5

Here note, after you have Multiply'd the *Depth* by the *ront*, and ſo found the Content of the *Ground-Plat* in *Feet*, alf of which, being added, gives the Content of a itch *Roof*, which reduce into *Squares.*

IV. By the Rodd, as in

Brick-Work.

Dimenſions are taken in Feet and Inches, and the Content given per the *Square* of a *Rodd* of 16 ½ *Foot.*

The Content of this Work is found in the ſame manner as all the other, to wit, by Multiplying the *Length* by the *Breadth,* and only differs from the former in two things.

1. When you have ſo found the *ſuperficial* Content in Feet, *divide it by* 272, *the ſquare Feet in one Rodd, the Quote is Rodds, the Content being always ſo given.*

2. That when 'tis more or leſs than a *Brick* and ½ thick it muſt be reduced into that thickneſs, being always meaſured by the ſquare Rodd of a *Brick* and ½. To do which obſerve,

Rule. *Multiply the Product of the Length and Height of the Wall by as many half Bricks as it contains in Thickneſs, which Product divided by* 3, *gives the Content of the Wall,* 1 *Brick and* ½ *thick, which ſuperficial Feet divide by* 272, *the Quote is ſquare Rodds reduced to* 1 *Brick and* ½, *or more ſhort by the following Table.*

	Brick.		
For	1 Subſtract	⅓	Reduces the ſaid Thickneſſes into 1 Brick and ½.
	2 Add	⅓	
	3 Multiply by	2	
	4½ Multiply by	3	
	6 Multiply by	4	

Ex. 1. Suppoſe a *Wall* of a *Garden* 213 *feet* about, and 13 *feet* 4 *inches* high, and 1 *Brick* and ½ thick, how many *Rodd* ?

213
13 . 4
2769 . 0
71 . 0 *Rodds, feet.*
272) 2840 . 0 (10 . 120
. 120

Ex. 2. How many *Rodd* of *Brick*-work in that *Wall* which is 40 *feet.* 7 *inch.* long, and 11 *feet* high, being 1 *Brick* thick ?

feet. inches.
40 . 7
11 . 0
3) 446 . 5
148 . 9 . 8
Rodd. feet. inch. part. 272) 297 . 7 . 4
Anſw. 1 . 25 . 7 . 4 1 . 25 . 7 . 4

Ex. 3. If a *Wall* be 254 *feet* about, and 12 *feet,* 7 *inches* high, and 3 Bricks thick, how many *Rodd?*

254
12 . 7
3048 . 0
127 . 0
21 . 2
Rodds, feet, inch. 3196 . 2
Anſw. 23 : 136 : 4 2 *Rodds, feet, inch.*
272) 6392 : 4 (23 : 136 : 4
. 952
136

Of *ſolid Meaſure.*

Which reſpecting *Length, Breadth* and *Thickneſs.*

Rule, *Multiply the length by the breadth, and that Product by the Depth.*

Otherwiſe, *Multiply the Area of the Baſe by the height, the Product gives the ſolid Content, which if in Feet, divide by* 27; *the ſolid Feet in one Yard, reduces them into ſolid Yards, if in Inches by* 1728, *gives ſolid feet,* &c.

Demonſtration.

For a *Surface* A B C D being imagined to move, ſuppoſe upwards or downwards, ſo that its *Points* A, B, C, D, deſcribe the *Lines* A H, B G, C F, D E, and conſequently, each *Line* therein deſcribe other *Surfaces,* &c. there will be generated by this Motion, a *Magnitude* called a Solid, having three *Dimenſions,* viz. *Length, Breadth* and *Thickneſs*; Therefore that *Surface* A B C D, being divided into little *Square Area's,* and moving the heighth A H, divided by a like meaſure for *Length,* may be conceiv'd to produce as many little *Cubes,* as is the number of the little *Area's* in the *Surface* A B C D, multiply'd by the number of Diviſions in the heighth A H.





Timber and *Stone.*

The Content given by the *Solid Foot.*

Ex. 1. How many *Solid Feet* is contain'd in that piece of *Timber*, whoſe Length is 17 *Feet* ½, Breadth 23 *Inches*, and Depth 2 *Feet* 7 *Inches?*

Anſw. 86 *feet*, 7 *inches*, 9 *parts* +.

Ex. 2. A Stone
Length 5 : 9
Breadth 3 : 11
Depth 2 : 8
How many ſolid Feet.

```
 3 . 11
 5 . 9
19 . 07
 1 . 11 . 6
 0 . 11 . 9
22 . 6 . 3
  2 . 8 . 0
45 . 0 . 6              feet, inches, parts
 11 . 3 . 1 . 6   Anſ. 60 : 00 : 8
  3 . 9 . 0 . 6
60 . 0 . 8 . 0
```

Digging.

The Content given by the *Solid Yard.*

Ex. 3. A Vault dug 9 *feet* Deep, 4½ Long, and 3 *feet*) *Inches* Broad, how many ſolid Yards?

```
          4 . 6 Length
          9 . 0 Depth
         40 . 6
          3 . 9 Breadth
        121 . 6
         20 . 3
         10 . 1 . 6
    27) 151 . 10 . 6     ya. feet, inch. par.
                    Anſw. 5 . 16 . 10 . 6
```

Ex. 4. How many Yards of digging will there be in that Cellar that is 25 *feet* 4 *inches* long, 15 . 8 Broad, and 7 *feet* ½ deep?

Anſwer 110 *yards*, 6 *feet*, 8 *inch.*

For the Menſuration of Round and Square Pillars, Pyramids, Cones, Globes, Cylinders, &c. ſee Mr. *Kerſey*'s 10th Chapter of his Appendix.

Soli Deo Gloria.

ADVERTISEMENTS.

There will be ſhortly publiſh'd

A *New Copy-Book*, containing great variety of Penmanſhip, in thirty Plates, having the Alphabet of Capital and Small Letters, Miniums, and Joyning of the moſt uſual Hands; together with an Alphabet of large *Roman* Capitals, with very little and free turn of ſtroak: The whole Book being adorn'd with near one hundred new Figures and Fancies, made more open, looſe and ſtrikable, than any hitherto publiſh'd, and that from the Original Papers of the late Incomparable Mr. *John Seddon*, who was juſtly thought, by reaſon of his Lively and Boundleſs Fancy, as great a Maſter in the Ornamental part of Penmanſhip as every our Nation produc'd. Done by *George Shelley* Writing Maſter, and Author of this Supplement. And Publiſh'd by Mr. *Thomas Read*, Clark of St. *Giles*'s in the Fields.

All ſorts of Mathematical Inſtruments, either for *Land* or *Sea*, are accurately made in Silver, Braſs, Ivory or Wood, as alſo all ſorts of Weather-Glaſſes, by *Richard Glynne* Mathematical Inſtrument-maker, next Door to the *Latin* Coffee-houſe in *Ave-maria*-lane, near St. *Paul*'s, *London.*

Thomas Banks
His Booke

14.1.2
14.1.2
2824

23
1 9 9
3. 1
7
2. 1 7
1 9 9
2 2. 0 7
2 2 0 7
1 5 4 4 9
0
4 4 1 4
4 1 4
8 4 9
8 7. 0 0 0 2 7
9 4 7
4. 0 9 5 9 8.
7 4 1 6 9
1 5 1 2 9 2 3
1 7 1 5
2 7 4
5 2 6 0
2 0 5
9 0
6. 0 3 1 0

$\frac{1}{3}$ 6

$\frac{2}{3}$ $\frac{1}{12}$ $\frac{1}{20}$

$\frac{3}{6}$

$\frac{3}{6}$

$\frac{2}{2}$ $\frac{2}{3}$

2)72 $\frac{11}{18}$

Sum

James

$\frac{2}{3}$ $\frac{1}{20}$

60

$\frac{1}{15}$

Thomas Bancks

28
28
13
141 65
0
8
7
27
2 0 6 2 0
2 2 1 1 1 9 9
9 2 2 0 6 2
3 1
3 7
2 1 7
1 9 9
4 1 6
4 1 6
6
2 4 9
4 1 6
1 6 6 4
1 7 3 0